

Airway Clearance Techniques for the Intubated Adult: A Scoping Review

Gabriella Cork¹ Clare Wade^{2,3} Alison Gordon⁴ Anna Vaughan-France⁵ Amelia Palmer⁶ Katy Walker⁷ Una Jones⁸

1. University of Liverpool
2. Northumbria University
3. The Newcastle upon Tyne Hospitals NHS Foundation Trust
4. University College London Hospitals NHS Foundation Trust
5. University Hospital Southampton NHS Foundation Trust
6. University Hospitals Sussex NHS Foundation Trust
7. The Walton Centre NHS Foundation Trust
8. Cardiff University

Correspondence: Gabriella Cork gcork@liverpool.ac.uk 0151 795 8103

Key Words: Intensive Care Unit; Physiotherapy; Sputum Clearance; Retained Secretions; Mechanical Ventilation

Objective

The aim of this scoping review was to understand the extent and type of evidence available in relation to airway clearance techniques in the intubated adult.

Introduction

This review was commissioned by the Association of Chartered Physiotherapists in Respiratory Care (ACPRC) special interest group as a method of summarising the available evidence on this topic on behalf of its members. Airway clearance in the intubated adult is a key objective of respiratory physiotherapists and although evidence-based guidelines exist in this area, there is no recent summary regarding the extent of the literature which could inform future research and clinical practice.

Inclusion criteria

Studies which investigated adults who were intubated either via an endo-tracheal or tracheostomy tube met the criteria. All study designs, including reviews, case reports and animal studies, which reported any physiotherapy-related airway clearance techniques were included.

Methods

The following databases were searched; SCOPUS, PubMed, PEDro, CINAHL Plus, and Clinical Trials Registry. The search was completed in December 2021 and limited to full text papers published since 2011. Following the key word search strategy, each title and abstract was screened for relevance to the scoping review aim and the study design was identified. Population, Intervention, Comparator and Outcome (PICO) data extraction was completed for all included papers in order to identify themes. The number and type of evidence retrieved, as well as key themes and outcomes were summarised.

Results:

The scoping review identified 138 suitable papers for inclusion. Of these, 11 were systematic reviews and 39 were randomised clinical trials, representing a moderately large evidence-base on this topic. Also included were other experimental, observational and qualitative studies, narrative reviews and animal and bench studies. Key interventions were identified including multi-modal chest physiotherapy, hyperinflation and manual chest compression techniques. Reported outcome measures were mainly short-term, such as sputum yield and oxygenation, whilst longer-term outcome measures such as ICU length of stay and ventilator-associated pneumonia (VAP) rates were reported less frequently. Outcome measures related to physiological stability were also reported by some studies.

Findings of the review were that airway clearance techniques for the intubated adult appear to be safe. There is a moderate body of evidence regarding their efficacy for short-term outcomes such as sputum yield, oxygenation and respiratory mechanics. There is limited evidence regarding their efficacy for longer-term outcomes.

Conclusion:

This scoping review summarises the extent of available evidence regarding airway clearance for intubated adults. Future research should focus on the effects of airway clearance techniques on longer-term outcome measures such as VAP rates and extubation outcome.

Introduction

The ACPRC Editorial Board is comprised of respiratory physiotherapy clinicians and academics who have volunteered through their ACPRC membership to be part of the Editorial Board. The purpose of the board is to lead scoping, commissioning, co-ordination and delivery of all new ACPRC guidance documents and resources, in order to facilitate knowledge sharing and drive improvements in the quality of care for respiratory patients.

The Editorial Board discussed potential areas for investigation which had been suggested by its membership and agreed that the area of airway clearance for the intubated adult should be prioritised. The lead author (GC), as a member of the Editorial Board, was nominated to lead the scoping review and other ACPRC members who were practising respiratory physiotherapist clinicians volunteered to assist with the process.

Airway clearance for the intubated adult is an important responsibility for respiratory physiotherapists in the Intensive Care Unit (ICU) (Twose et al 2019) and involves the mobilisation and subsequent removal of respiratory secretions via the endotracheal or tracheostomy tube. Intubation and the associated mechanical ventilation, prolonged recumbency and sedation result in reduced cough efficacy, reduced mucociliary transport and atelectasis which can in turn lead to retained secretions and ventilator-associated pneumonia (Piccin *et al*, 2011; Beuret *et al*, 2009; Malbouisson *et al*, 2001; Konrad *et al*, 1995; Konrad *et al*, 1994). Physiotherapeutic techniques to assist with the removal of sputum from the intubated patient such as manual chest compression, hyperinflation and positioning are frequently used by physiotherapists (Connolly *et al*, 2020; Skinner *et al*, 2015). However,

evidence investigating the efficacy of such techniques has been deemed overall of poor quality with conflicting findings (Stiller, 2013).

Recent Faculty of Intensive Care Medicine (FICM) guidelines recommend “targeted airway clearance interventions” for invasively ventilated patients but do not stipulate which airway clearance interventions should be utilised (FICM, 2019, p. 51). Furthermore, the same publication recommends that individual physiotherapy services should develop their own evidence-based guidelines for the use of airway clearance techniques. A major purpose of this scoping review was to determine whether there is sufficient evidence available on this topic to inform collaborative clinical guidelines.

A preliminary search of SCOPUS and the Cochrane Database of Systematic Reviews was conducted and whilst recent reviews in this area exist, they have focused on individual techniques such as manual therapy (Borges *et al*, 2017) or hyperinflation (Anderson *et al*, 2015; Paulus *et al*, 2012), on specific populations such as those with community-acquired pneumonia (van der Lee *et al*, 2020) and traumatic brain injury (Newman *et al*, 2018), or on specific outcomes such as ventilation-associated pneumonia (VAP) rates (Wang *et al*, 2019). One systematic review (Stiller, 2013) did have a wider focus and included a variety of physiotherapeutic techniques and outcomes relevant to airway clearance, however a number of new experimental studies have subsequently been published. These have not yet been captured by guidelines, recommendations or systematic review and may further contribute to the knowledge base in this topic.

The aim of this scoping review is to understand the extent and type of evidence in relation to airway clearance in the intubated adult in order to inform future recommendations for respiratory physiotherapy clinicians and researchers.

Review question

What is the extent of the current evidence-base in relation to airway clearance in the intubated adult?

Eligibility criteria

Participants

Adults who were intubated either via endotracheal or tracheostomy tube at the time of the investigation. Paediatric studies were excluded.

Concept

Airway clearance techniques that are performed by physiotherapists as summarised by Berry and Marti (2014). Additional airway clearance techniques that are more commonly performed by medical staff such as bronchoscopy or that are pharmacological in nature were excluded. Techniques such as automated lateral bed rotation, humidification or endotracheal suctioning alone were excluded as these were deemed to be primarily routine, nurse-delivered interventions.

Context

Airway clearance techniques for the intubated adult are usually performed in the ICU, however studies were not excluded if they investigated intubated adults in other clinical

settings such as weaning units or post-op recovery areas. This scoping review was planned and initial searches conducted prior to the global Covid-19 pandemic, therefore studies relating to the treatment of Covid-19 were excluded as they were deemed by the co-authors to be beyond the initial remit and purpose of the review.

Types of Sources

This scoping review considered both experimental and quasi-experimental study designs including randomized controlled clinical trials, non-randomized controlled trials and before and after studies. In addition, analytical observational studies including prospective and retrospective cohort studies, case-control studies and analytical cross-sectional studies were considered for inclusion. This review also included descriptive observational study designs including case series, individual case reports and descriptive cross-sectional studies as well as animal and bench studies. Qualitative studies on this topic were additionally summarised alongside systematic reviews and meta-analyses that met the inclusion criteria.

Opinion papers (including editorials) as well as conference abstracts were excluded.

Methods

The scoping review was conducted in accordance with the JBI methodology for scoping reviews (Peters *et al*, 2020).

Search strategy

An initial limited search of SCOPUS was undertaken to identify articles on the topic. The text words contained in the titles and abstracts of relevant articles, the index terms used to

describe the articles, and a collaborative, iterative process by the co-authors were used to develop a full search strategy (see Appendix 1 for full SCOPUS search strategy). The search strategy, including all identified keywords and index terms, was adapted for each of the included databases. The reference lists of all review papers were subsequently screened for additional studies.

Studies published in any language where a full English version was available were included. Studies published from 2011 onwards were included as the most comprehensive systematic review was published in 2013, and the vast majority of its 85 included papers were published pre-2011 (Stiller, 2013).

The databases searched were SCOPUS, PubMed, CINAHL Plus, and PEDro. Google Scholar search engine was additionally employed using the same search terms, limited to the initial 500 papers due to default sort by relevance. The Clinical Trials Registry was also searched for unpublished studies that were completed within the previous three years which might reasonably be in the process of being published at the time of the review. The final search was completed in December 2021.

Source of Evidence selection

Following the search, titles and abstracts were screened by co-authors (GC, AG, AVF, KW, AP) for assessment against the inclusion criteria. Potentially relevant sources were retrieved in full and uploaded into EndNote X9, 2018 (Clarivate Analytics, PA, USA) and duplicates removed. The full text of selected citations was assessed in detail against the inclusion criteria by the lead author (GC) and cross-checked independently by CW. Reasons for the exclusion of evidence following full text review were recorded and reported (Figure 1).

Data Extraction and Synthesis

Data regarding study design, population, intervention, comparator and outcome (PICO) was extracted from papers included in the scoping review by the co-authors using a data extraction tool developed by UJ. The data extracted included key findings relevant to the review question. Themes were identified during the data extraction process and the papers grouped both by type of evidence and sub-themes within the overall airway clearance topic. For papers which included more than one type of airway clearance, the intervention was classified as “multi-modal chest physiotherapy”. Key outcome measures were also identified. Due to the breadth of the scoping review, quality assessment of the experimental papers was not undertaken beyond classifying them according to their study design.

Results

******INSERT FIGURE 1 NEAR HERE******

The scoping review retrieved 138 relevant papers (see Figure 1). Of these, 11 were systematic reviews (summarised in Appendix 1) and 39 were randomised clinical trials (summarised in Appendix 2), see Figure 2 for full break-down of papers by evidence-type and Appendices 3-8 for summaries of all other included papers.

****** INSERT FIGURE 2 NEAR HERE******

The most common reported intervention was multi-modal chest physiotherapy with 45 publications exploring this topic. Common airway clearance techniques such as hyperinflation and manual chest compressions were also extensively studied. Figure 3 gives a full break-down of the papers included in the scoping review according to their theme.

****** INSERT FIGURE 3 NEAR HERE******

The key interventions, outcome measures and findings from comparative studies included in this scoping review are summarised in Table 1. The most commonly reported outcome measures were sputum yield and oxygenation.

****** INSERT TABLE 1 NEAR HERE******

A summary of the non-experimental research included in this scoping review is provided in Table 2.

****** INSERT TABLE 2 NEAR HERE******

Discussion

The papers retrieved by this deliberately wide-ranging scoping review were diverse and as well as clinical efficacy papers, included assessment of the requirements for airway clearance techniques (ACTs), physiological effects of ACTs, opinions of caregivers, service delivery and clinical recommendations.

Non-Experimental Research

Clinician opinion regarding airway clearance techniques using surveys of current practice has been the subject of a number of recent studies (Bhat *et al*, 2014; Lottering and van Aswegen, 2016; Matilde *et al*, 2018; van der Lee *et al*, 2017; Rose *et al*, 2016; Hayes *et al*, 2011; Stilma *et al*, 2021). These surveys have highlighted that a number of varying airway clearance and cough augmentation techniques are used by critical care clinicians. Studies reported heterogeneity of intensity and combination of ACTs in addition to variation in clinical practice. Rationale for commencing airway clearance techniques is similar across studies, including to aid in sputum clearance and promotion of improved alveolar recruitment and ventilation. Some studies highlight that lack of knowledge, training, and expertise may contribute to reduced adoption of techniques such as mechanical in-exsufflation (Rose *et al*, 2016) and ventilator hyperinflation (Hayes *et al*, 2011). Such studies recognise the lack of clinical guidance in this area.

Qualitative studies with physiotherapists and wider critical care clinicians highlight the importance of teamwork, clinical reasoning, clinical experience and communication as key in the selection and effective implementation of airway clearance interventions for mechanically ventilated adults (Connolly *et al*, 2020; van der Lee *et al*, 2020).

A recent focus of non-experimental research has been the production of clinical guidelines using a Delphi technique to achieve expert consensus, specifically to identify core clinical competencies for practitioners implementing airway clearance techniques (Skinner *et al*, 2016; Twose *et al*, 2019) and best practice for the treatment of community-acquired pneumonia (Van der Lee *et al*, 2019). Expert consensus panels recognise that physiotherapy competence in airway clearance interventions such as hyperinflation techniques, manual chest wall techniques, positioning, normal saline instillation and suction are a minimum standard of practice for physiotherapists working in ICU in their respective countries (Skinner *et al*, 2016; Twose *et al*, 2019). This scoping review did not retrieve any recent clinical guidelines to aid in the selection, implementation or evaluation of airway clearance techniques for non-specific intubated adults.

Experimental Research: Short-term outcomes

Papers focusing on clinical efficacy investigated interventions that enhanced inspiratory volume and/or expiratory flow. The effects of such airway clearance techniques were primarily reported on short-term outcomes such as oxygenation, sputum yield, respiratory mechanics (for example, dynamic compliance) and peak expiratory flow (PEF; either absolute PEF or peak inspiratory-expiratory flow (PIF:PEF) ratio). Although sputum yield is the most direct outcome measure for the efficacy of airway clearance techniques, oxygenation and respiratory mechanics are reported frequently. Techniques which appear to enhance sputum yield include adjuncts such as high-frequency chest wall oscillation (HFCWO) and oscillatory positive expiratory pressure (OPEP) devices (Chicayban *et al*, 2011; Jones *et al*, 2013; Kluayhomthong *et al*, 2019; Kuyruklyildiz *et al*, 2016), head down positioning (Shetty *et al*, 2020), and multi-modal chest physiotherapy (Van der Lee *et al*, 2020; Naue *et al*, 2014). The

evidence for the effect of hyperinflation, manual techniques and cough augmentation on sputum yield is mixed, with contradictory findings. Several studies have reported no difference in effect between manual hyperinflation and ventilator hyperinflation regarding sputum yield, including a systematic review (Anderson *et al*, 2015).

The most effective interventions to improve oxygenation in the short term appear to be manual and ventilator hyperinflation with multiple studies reporting statistically significant if not necessarily clinically significant findings in favour of this intervention (Paulus *et al*, 2012; Malekzadeh *et al*, 2016; Mohamed *et al*, 2017 Frank *et al*, 2015; El-Deen and Ahmed, 2013; Ibrahim and Mohamed, 2018; Raafat and Elbasiouny, 2011; Waqas *et al*, 2014). Improvements in respiratory mechanics such as static and dynamic lung compliance were reported with multi-modal chest physiotherapy by multiple systematic reviews (Andrews *et al*, 2013; Stiller, 2013; van der Lee *et al*, 2020). Several experimental studies reported a similar effect with airway clearance adjuncts (Chicayban *et al*, 2011; Chuang *et al*, 2017; Longhini *et al*, 2020) which seems to be an emerging area of research that warrants further attention.

Animal studies have investigated the effect of airway clearance techniques on PEF, particularly to determine whether the threshold for mucous movement can be achieved by head down positioning and manual techniques (Li Bassi *et al*, 2014; Marti *et al*, 2013). A number of clinical studies have also explored this outcome measure with adjuncts (Chicayban *et al*, 2011) and manual techniques (Amaral *et al*, 2019; Oliveira *et al*, 2019) being shown to improve PEF.

Cardiovascular and neurological stability have been investigated in a number of studies and whilst some statistically significant deteriorations have been reported during various airway clearance techniques, authors concluded that these were transient and non-clinically

significant. This suggests that ACTs are safe for the intubated patient (Blattner *et al*, 2017; Ferreira *et al*, 2013).

Experimental Research: Longer-term outcomes

Longer-term outcome measures such as mortality, ICU length of stay, ventilator-acquired pneumonia (VAP) rates and extubation outcome were reported by some studies although far less frequently than short-term outcomes. Understandably, these longer-term outcome measures have been a focus of systematic reviews (Pouzuelo-Carrascosa *et al*, 2018; Stiller, 2013; van der Lee *et al*, 2020; Wang *et al*, 2018). With the exception of mortality (Pouzuelo-Carrascosa *et al*, 2018), systematic reviews included in this scoping review report no significant effect of airway clearance techniques on any of these longer-term outcome measures (Stiller, 2013; van der Lee *et al*, 2020; Wang *et al*, 2018).

This scoping review retrieved a number of experimental studies reporting reduced VAP rates with airway clearance adjuncts (Spapen *et al*, 2015; Kuyrukluylidiz *et al*, 2016), head-down positioning (Li Bassi *et al*, 2014), manual techniques (El-Hamid *et al*, 2017) and multi-modal chest physiotherapy (Mohamed *et al*, 2017; Pattanshetty and Gaude, 2011). Similarly, an improved likelihood of extubation success has been reported in some studies with cough augmentation (Goncalves *et al*, 2012) and multi-modal chest physiotherapy (Wang *et al*, 2018). Whilst quality was not assessed as part of this scoping review, these findings suggest that the effect of airway clearance techniques on VAP rates and extubation outcome may warrant further investigation.

Limitations of this scoping review

This scoping review was intentionally limited to airway clearance techniques that are typically delivered by respiratory physiotherapists in the adult ICU. Due to this, some aspects of airway clearance such as routine suctioning, humidification, regular repositioning, bronchoscopy and pharmacological interventions were not included.

A major methodological limitation was the lack of quality assessment of the included papers. Due to the wide-ranging remit of the review and number of papers retrieved, this was beyond the scope of this review and was not deemed necessary in order to meet the aims and objectives set out in advance. Randomised clinical trials were carefully screened and if they failed to meet the criteria for randomisation as described by PEDro (de Morton, 2009), they were classified as “other experimental” studies.

Future Research

Future experimental research is still required to determine the effectiveness of airway clearance techniques in the intubated adult. Existing randomised clinical trials have focused on short-term outcomes and therefore a reasonably comprehensive body of evidence is available for common interventions such as hyperinflation and manual techniques in this regard. There are emerging interventions with an increasing evidence-base such as cough augmentation and airway clearance adjuncts. As an outcome measure, PEF appears to be influenced by ACTs, however its relevance to other, more clinical outcomes would benefit from further experimental investigation. Another under-researched area is positioning for airway clearance, despite this being a routinely used intervention for intubated adults (Matilde *et al*, 2018). A focus for future research should be the effect of airway clearance on longer-term outcomes, particularly VAP rate and extubation success as these appear to be outcomes that may be influenced by ACTs.

Future reviews should be carefully considered and rationalised. There would be merit in an updated systematic review investigating the effect of manual techniques as several recent studies have not yet been captured by such a review. Furthermore, airway clearance adjuncts for the intubated patient have not yet been investigated by systematic review and this may be an area that warrants the same. A plethora of narrative reviews on this topic are already available and therefore any future narrative reviews should have a clear and unique focus.

Conclusion

This scoping review was undertaken as an area of priority for the ACPRC Editorial Board. The extent of the available evidence regarding airway clearance for intubated adults has been summarised, thus achieving the aim of the scoping review. Included papers were diverse and wide-ranging in their findings. Overall, the evidence-base regarding the efficacy of airway clearance techniques for short-term clinical outcome measures is moderately large. Currently, there is limited evidence regarding their efficacy for longer-term outcome measures. There is a moderate body of evidence reporting that airway clearance techniques are safe for the intubated adult.

Future clinical research should focus on the effects of airway clearance techniques on longer-term outcome measures such as VAP rates and extubation outcome as well as investigating common but under-researched interventions such as positioning. As the quality of the studies was not assessed in this scoping review, future work is needed to develop clinical recommendations based on both short- and long-term efficacy of airway clearance techniques for adults who are intubated.

Acknowledgements

Association of Chartered Physiotherapists in Respiratory Care Editorial Board.

Gareth Cornell for review of the database search terms.

Funding

There was no funding provided for this scoping review. All participants gave their time voluntarily.

Conflicts of interest

There is no conflict of interest in this project.

References

- Amaral, B. L. R., de Figueiredo, A. B., Lorena, D. M., Oliveira, A. C. O., Carvalho, N. C. and Volpe, M. S. (2020) 'Effects of ventilation mode and manual chest compression on flow bias during the positive end- and zero end-expiratory pressure manoeuvre in mechanically ventilated patients: a randomised crossover trial', *Physiotherapy (United Kingdom)*, 106, pp. 145-153.
- Ambrosino, N., Janah, N. and Vaghegini, G. (2011) 'Physiotherapy in critically ill patients', *Rev Port Pneumol*, 17(6), pp. 283-288.
- Anderson, A., Alexanders, J., Sinani, C., Hayes, S. and Fogarty, M. (2015) 'Effects of ventilator vs manual hyperinflation in adults receiving mechanical ventilation: A systematic review of randomised clinical trials', *Physiotherapy (United Kingdom)*, 101(2), pp. 103-110.
- Andrews, J., Sathe, N. A., Krishnaswami, S. and McPheeters, M. L. (2013) 'Nonpharmacologic airway clearance techniques in hospitalized patients: a systematic review', *Respir Care*, 58(12), pp. 2160-86.
- Arora, S., Flower, O., Murray, N. P. and Lee, B. B. (2012) 'Respiratory care of patients with cervical spinal cord injury: a review', *Crit Care Resusc*, 14(1), pp. 64-73.
- Ashtankar, A. and Kazi, A. (2019) 'Comparative Effect of Proprioceptive Neuromuscular Facilitation (PNF) and Chest Physiotherapy with Chest Physiotherapy alone on SP02, Heart Rate, Respiratory Rate, & Lung Compliance in Mechanically Ventilated Patient', *J. Pharm. Sci. & Res*, 11(10), pp. 3514-3518.
- Bach, J. R., Siquee, D. M., Saporito, L. R. and Botticello, A. L. (2015) 'Efficacy of mechanical insufflation-exsufflation in extubating unweanable subjects with restrictive pulmonary disorders', *Respiratory Care*, 60(4), pp. 477-483.
- Belli, S., Cattaneo, D., D'Abrosca, F., Prince, I., Savio, G. and Balbi, B. (2019) 'A pilot study on the non-invasive management of tracheobronchial secretions in tracheostomised patients', *Clinical Respiratory Journal*, 13(10), pp. 637-642.
- Bennett, B. G., Thomas, P. and Ntoumenopoulos, G. (2015) 'Effect of inspiratory time and lung compliance on flow bias generated during manual hyperinflation: A bench study', *Respiratory Care*, 60(10), pp. 1449-1458.
- Berney, S., Haines, K. and Denehy, L. (2012) 'Physiotherapy in critical care in Australia', *Cardiopulm Phys Ther J*, 23(1), pp. 19-25.
- Berry, M. P. and Martí, J. D. (2014) 'Clinical management of secretion retention in critically ill patients who are intubated and mechanically ventilated', *Current Respiratory Medicine Reviews*, 10(3), pp. 163-175.

Berti, J. S. W., Tonon, E., Ronchi, C. F., Berti, H. W., de Stefano, L. M., Gut, A. L., Padovani, C. R. and Ferreira, A. L. A. (2012) 'Manual hyperinflation combined with expiratory rib cage compression for reduction of length of ICU stay in critically ill patients on mechanical ventilation', *Jornal Brasileiro de Pneumologia*, 38(4), pp. 477-486.

Beuret, P., Roux, C., Auclair, A., Nourdine, K., Kaaki, M., & Carton, M.-J. (2009). Interest of an objective evaluation of cough during weaning from mechanical ventilation. *Intensive Care Medicine*, 35(6)

Bhat, A., Chakravarthy, K. and Rao, B. K. (2014) 'Chest physiotherapy techniques in neurological intensive care units of India: A survey', *Indian Journal of Critical Care Medicine*, 18(6), pp. 363-368.

Bhoir, T., Kazi, A., Shetty, A., Gunjal, S., Lamuvell, M. and Chordiya, S. (2017) 'Acute response of manual hyperinflation versus ventilator hyperinflation in addition to chest physiotherapy on critically ill mechanically ventilated patients', *International Journal of Multidisciplinary Research and Development*, 4(6), pp. 51-56.

Blattner, C. N., Dos Santos, R. S., Dias, F. S., Dias, A. S., Mestriner, R. G. and Vieira, S. R. R. (2017) 'Acute bag-valve breathing maneuvers plus manual chest compression is safe during stable septic shock: A randomized clinical trial', *Revista Brasileira de Terapia Intensiva*, 29(1), pp. 14-22.

Borges, L. F., Saraiva, M. S., Saraiva, M. A. S., Macagnan, F. E. and Kessler, A. (2017) 'Expiratory rib cage compression in mechanically ventilated adults: systematic review with meta-analysis', *Rev Bras Ter Intensiva*, 29(1), pp. 96-104.

Bousarri, M. P., Shirvani, Y., Agha-Hassan-Kashani, S. and Nasab, N. M. (2014) 'The effect of expiratory rib cage compression before endotracheal suctioning on the vital signs in patients under mechanical ventilation', *Iran J Nurs Midwifery Res*, 19(3), pp. 285-9.

Castro, A. A. M., Calil, S. R., Freitas, S. A., Oliveira, A. B. and Porto, E. F. (2013) 'Chest physiotherapy effectiveness to reduce hospitalization and mechanical ventilation length of stay, pulmonary infection rate and mortality in ICU patients', *Respiratory Medicine*, 107(1), pp. 68-74.

Chapman, R. L., Shannon, H., Koutoumanou, E. and Main, E. (2019) 'Effect of inspiratory rise time on sputum movement during ventilator hyperinflation in a test lung model', *Physiotherapy (United Kingdom)*, 105(2), pp. 283-289.

Chen, Y. H., Yeh, M. C., Hu, H. C., Lee, C. S., Li, L. F., Chen, N. H., Huang, C. C. and Kao, K. C. (2016) 'Effects of Lung Expansion Therapy on Lung Function in Patients with Prolonged Mechanical Ventilation', *Can Respir J*, 2016, p. 5624315.

Chicayban, L. M., Hemétrio, A. C. and Azevedo, L. T. R. (2020) 'Comparison of the effects of voluntary and involuntary breath stacking techniques on respiratory mechanics and lung function patterns in tracheostomized patients: A randomized crossover clinical trial', *Jornal Brasileiro de Pneumologia*, 46(4), pp. 1-8.

Chicayban, L. M., Zin, W. A. and Guimarães, F. S. (2011) 'Can the Flutter Valve improve respiratory mechanics and sputum production in mechanically ventilated patients? A randomized crossover trial', *Heart and Lung: Journal of Acute and Critical Care*, 40(6), pp. 545-553.

Chuang, M. L., Chou, Y. L., Lee, C. Y. and Huang, S. F. (2017) 'Instantaneous responses to high-frequency chest wall oscillation in patients with acute pneumonic respiratory failure receiving mechanical ventilation: a randomized controlled study', *Medicine*, 96(9), p. e5912.

Clinkscale, D., Spihlman, K., Watts, P., Rosenbluth, D. and Kollef, M. H. (2012) 'A randomized trial of conventional chest physical therapy versus high frequency chest wall compressions in intubated and non-intubated adults', *Respiratory Care*, 57(2), pp. 221-228.

Connolly, B., Barclay, M., Blackwood, B., Bradley, J., Anand, R., Borthwick, M., Chikhani, M., Dark, P., Shyamsundar, M., Warburton, J., McAuley, D. F. and O'Neill, B. (2020) 'Airway clearance techniques and use of mucoactive agents for adult critically ill patients with acute respiratory failure: a qualitative study exploring UK physiotherapy practice', *Physiotherapy (United Kingdom)*, 108, pp. 78-87.

Cork, G., Barrett, N. and Ntoumenopoulos, G. (2014) 'Justification for chest physiotherapy during ultra-protective lung ventilation and extra-corporeal membrane oxygenation: a case study', *Physiother Res Int*, 19(2), pp. 126-8.

Coutinho, W., Vieira, P., Kutchak, F., Dias, A., Rieder, M. and Forgiarini, L., Jr. (2018) 'Comparison of mechanical insufflation-Exsufflation and endotracheal suctioning in mechanically ventilated patients: Effects on respiratory mechanics, hemodynamics, and volume of secretions', *Indian Journal of Critical Care Medicine*, 22(7), pp. 485-490.

de Camillis, M. L. F., Savi, A., Rosa, R. G., Figueiredo, M., Wickert, R., Borges, L. G. A., Galant, L. and Teixeira, C. (2018) 'Effects of mechanical insufflation-exsufflation on airway mucus clearance among mechanically ventilated icu subjects', *Respiratory Care*, 63(12), pp. 1471-1477.

de Cerqueira Neto, M. L., Moura, A. V., Fontes Cerqueira, T. C., Aquim, E. E., Reá-Neto, A., Oliveira, M. C., da Silva Júnior, W. M., Santana-Filho, V. J. and Scola, R. H. (2013) 'Acute effects of physiotherapeutic respiratory maneuvers in critically ill patients with craniocerebral trauma', *Clinics*, 68(9), pp. 1210-1214.

de Godoy, A. C., Yokota Cde, O., Araújo, II and de Freitas, M. I. (2011) 'Can manual hyperinflation maneuvers cause aspiration of oropharyngeal secretions in patients under mechanical ventilation?', *Rev Bras Anesthesiol*, 61(5), pp. 556-60, 305-7.

de Morton, N. (2009) The PEDro scale is a valid measure of the methodological quality of clinical trials: a demographic study. *Australian Journal of Physiotherapy*. 55 (2) 129-133

Della, V., Fabiana, Oliveira, Rosmari, A. R. A., Dragosavac and Desanka (2012) 'Effects of manual chest compression and decompression maneuver on lung volumes, capnography and pulse oximetry in patients receiving mechanical ventilation', *Rev Bras Fisioter*, 16(5), pp. 354-9.

Denehy, L., Granger, C. L., El-Ansary, D. and Parry, S. M. (2018) 'Advances in cardiorespiratory physiotherapy and their clinical impact', *Expert Review of Respiratory Medicine*, 12(3), pp. 203-215.

Dennis, D., Jacob, W. and Budgeon, C. (2012) 'Ventilator versus manual hyperinflation in clearing sputum in ventilated intensive care unit patients', *Anaesthesia and Intensive Care*, 40(1), pp. 142-149.

Dennis, D. M., Duncan, C. N., Pinder, M., Budgeon, C. A. and Jacob, W. J. (2016) 'Performance of manual hyperinflation: consistency and modification of the technique by intensive care unit nurses during physiotherapy', *Journal of Clinical Nursing*, 25(15-16), pp. 2295-2304.

Eden, A., Purkiss, C., Cork, G., Baddeley, A., Morris, K., Carey, L., Brown, M., McGarrigle, L. and Kennedy, S. (2017) 'In-patient physiotherapy for adults on veno-venous extracorporeal membrane oxygenation – United Kingdom ECMO Physiotherapy Network: A consensus agreement for best practice', *Journal of the Intensive Care Society*, 18(3), pp. 212-220.

El-Deen, H. and Ahmed, K. (2013) 'Acute response of manual hyperinflation technique on airway secretion clearance in postoperative Intubated patients', *Journal of medical research and practice*, 2(9), pp. 238-45.

El-Hamid, G., Shams, T., Ali, A. and Awad, S. (2017) 'Effect of Using Different Modalities of Chest Physiotherapy on Prevention of Ventilator Associated Pneumonia', *Journal of Nursing and Health Science*, 6(5), pp. 19-35.

Faculty of Intensive Care Medicine & Intensive Care Society (2019) 'Guidelines for the Provision of Intensive Care Services', Edition 2.

Fernandez-Restrepo, L., Shaffer, L., Amalakuhan, B., Restrepo, M. I., Peters, J. and Restrepo, R. (2017) 'Effects of intrapulmonary percussive ventilation on airway mucus clearance: A bench model', *World J Crit Care Med*, 6(3), pp. 164-171.

Ferreira, L. L., Valenti, V. E. and Vanderlei, L. C. M. (2013) 'Chest physiotherapy on intracranial pressure of critically ill patients admitted to the intensive care unit: A systematic review', *Revista Brasileira de Terapia Intensiva*, 25(4), pp. 327-333.

Frank, U., Frank, K. and Zimmermann, H. (2015) 'Effects of Respiratory Therapy (bagging) on Respiratory Function, Swallowing Frequency and Vigilance in Tracheotomized Patients in Early Neurorehabilitation', *Pneumologie*, 69(7), pp. 394-399.

Genc, A., Akan, M. and Gunerli, A. (2011) 'The effects of manual hyperinflation with or without rib-cage compression in mechanically ventilated patients', *Italian Journal of Physiotherapy*, 1(2), pp. 48-54.

Goni-Viguria, R., Yoldi-Arzo, E., Casajus-Sola, L., Aquerreta-Larraya, T., Fernandez-Sangil, P., Guzman-Unamuno, E. and Moyano-Berardo, B. M. (2018) 'Respiratory physiotherapy in intensive care unit: bibliographic review', *Enfermeria Intensiva*, 29(4), pp. 168-181.

Gonçalves, M. R., Honrado, T., Winck, J. C. and Paiva, J. A. (2012) 'Effects of mechanical insufflation-exsufflation in preventing respiratory failure after extubation: a randomized controlled trial', *Crit Care*, 16(2), p. R48.

Gosselink R, Bott J, Johnson M et al. Physiotherapy for adult patients with critical illness: Recommendations of the European Respiratory Society and European Society of Intensive Care Medicine task force on physiotherapy for critically ill patients. *Intensive Care Med.* 2008 Jul ;34(7):1188-99.

Gosselink, R., Clerckx, B., Robbeets, C., Vanhullebusch, T., Vanpee, G. and Segers, J. (2011) 'Physiotherapy in the intensive care unit', *Netherlands Journal of Critical Care*, 15(2), pp. 1-10.

Grammatopoulou, E., Charmpas, T. N., Strati, E. G., Nikolaos, T., Evagelodimou, A., Vlassia, B. and Skordilis, E. K. (2017) 'The scope of physiotherapy services provided in public ICUs in Greece: A pilot study', *Physiotherapy Theory and Practice*, 33(2), pp. 138-146.

Guimarães, F. S., Lopes, A. J., Constantino, S. S., Lima, J. C., Canuto, P. and de Menezes, S. L. S. (2014) 'Expiratory rib cage in mechanically ventilated subjects: A randomized crossover trial', *Respiratory Care*, 59(5), pp. 678-685.

Guner, S. I. and Korkmaz, F. D. (2015) 'Investigation of the effects of chest physiotherapy in different positions on the heart and the respiratory system after coronary artery bypass surgery', *Toxicol Ind Health*, 31(7), pp. 630-7.

Guérin, C., Bourdin, G., Leray, V., Delannoy, B., Bayle, F., Germain, M. and Richard, J. C. (2011) 'Performance of the CoughAssist insufflation-exsufflation device in the presence of an endotracheal tube or tracheostomy tube: A bench study', *Respiratory Care*, 56(8), pp. 1108-1114.

Hariedy, N., Mohammed, W., Abdel-Aziz, M. and Morsy, K. (2015) 'Chest Physiotherapy and Recruitment Maneuvers: Effects on Lung Mechanics and Pulmonary Complications among Mechanically Ventilated Patients with Acute Lung Injury', *Journal of Nursing and Health Science*, 4(4), pp. 17-32.

Hayes, K., Seller, D., Webb, M., Hodgson, C., Holland, A. (2011) 'Ventilator hyperinflation: a survey of current physiotherapy practice in Australia and New Zealand', *New Zealand Journal of Physiotherapy*, 39(3), pp. 124-130.

Hewitt, N., Bucknall, T. and Faraone, N. M. (2016) 'Lateral positioning for critically ill adult patients', *Cochrane Database Syst Rev*, 2016(5), p. Cd007205.

Hongrattana, G., Reungjui, P. and Jones, C. U. (2014) 'Acute hemodynamic responses to 30 degrees head-down postural drainage in stable, ventilated trauma patients: a randomized crossover trial', *Heart & Lung*, 43(5), pp. 399-405.

Hyun, S. E., Lee, S. M. and Shin, H. I. (2021) 'Peak expiratory flow during mechanical insufflation-exsufflation: Endotracheal tube versus face mask', *Respiratory Care*, 66(12), pp. 1815-1823.

Ibrahim, A. and Mohamed, A. (2018) 'Effect of Adding Different Physiotherapy Modalities to Standard Chest Physiotherapy for Critical Ill Patients.', *Research Journal of Pharmaceutical, Biological and Chemical Sciences*, 9(3), pp. 144-52.

Jacob, W., Dennis, D., Jacques, A., Marsh, L., Woods, P. and Hebden-Todd, T. (2021) 'Ventilator hyperinflation determined by peak airway pressure delivered: A randomized crossover trial', *Nurs Crit Care*, 26(1), pp. 14-19.

Jang, M. H., Shin, M. J. and Shin, Y. B. (2019) 'Pulmonary and Physical Rehabilitation in Critically Ill Patients', *Acute Crit Care*, 34(1), pp. 1-13.

Jiandani, M. and Patel, A. (2018) 'Cardiorespiratory Response to Chest Physiotherapy in Intensive Respiratory Care Unit', *International Journal of Health Sciences & Research*, 8(7), pp. 159-65.

Jones, C. U., Kluayhomthong, S., Chaisuksant, S. and Khrisanapant, W. (2013) 'Breathing exercise using a new breathing device increases airway secretion clearance in mechanically ventilated patients', *Heart and Lung: Journal of Acute and Critical Care*, 42(3), pp. 177-182.

Jyoti, M. (2018) 'Role of Physiotherapist in Critical Care Situations: Recent Perspective', *Journal of Cardiac Critical Care*, 2(1), pp. 14-18.

Kallet, R. H. (2013) 'Adjunct therapies during mechanical ventilation: airway clearance techniques, therapeutic aerosols, and gases', *Respiratory Care*, 58(6), pp. 1053-73.

Kluayhomthong, S., Ubolsakka-Jones, C., Domthong, P., Reechaipichitkul, W. and Jones, D. A. (2019) 'The immediate effects of breathing with oscillated inspiratory and expiratory airflows on secretion clearance in intubated patients with cervical spinal cord injury', *Spinal Cord*, 57(4), pp. 308-316.

Kohan, M., Rezaei-Adaryani, M., Najaf-Yarandi, A., Hoseini, F. and Mohammad-Taheri, N. (2014) 'Effects of expiratory ribcage compression before endotracheal suctioning on arterial blood gases in patients receiving mechanical ventilation', *Nursing in Critical Care*, 19(5), pp. 255-261.

Konrad F, Schiener R, Marx T, Georgief M. Ultrastructure and mucociliary transport of bronchial respiratory epithelium in intubated patients. *Intensive Care Med.* 1995;21:482–489

Konrad F, Schreiber T, Brecht-Kraus D, et al: Mucociliary transport in ICU patients. *Chest* 1994; 105:237–241

Kubo, T., Osuka, A., Kabata, D., Kimura, M., Tabira, K. and Ogura, H. (2021) 'Chest physical therapy reduces pneumonia following inhalation injury', *Burns*, 47(1), pp. 198-205.

Kuroiwa, R., Tateishi, Y., Oshima, T., Inagaki, T., Furukawa, S., Takemura, R., Kawasaki, Y. and Murata, A. (2021) 'Mechanical insufflation-exsufflation for the prevention of ventilator-associated pneumonia in intensive care units: A retrospective cohort study', *Indian Journal of Critical Care Medicine*, 25(1), pp. 62-66.

Kuyruklyildiz, U., Binici, O., Kupeli, I., Erturk, N., Gulhan, B., Akyol, F., Ozcicek, A., Onk, D. and Karabakan, G. (2016) 'What Is the Best Pulmonary Physiotherapy Method in ICU?', *Canadian Respiratory Journal*, 2016, pp. 1-5.

Lee, D., Park, S., Yeom, S., Han, S., Lee, S. and Ryu, J. (2011) 'High-frequency Chest Wall Oscillation Therapy: Clinical Effectiveness in the Patients with Pulmonary Contusion', *Korean Journal of Critical Care Medicine*, 26(4), pp. 256-60.

Li Bassi, G., Marti, J. D., Saucedo, L., Rigol, M., Roca, I., Cabanas, M., Muñoz, L., Ranzani, O. T., Giunta, V., Luque, N., Esperatti, M., Gabarrus, A., Fernandez, L., Rinaudo, M., Ferrer, M., Ramirez, J., Vila, J. and Torres, A. (2014) 'Gravity predominates over ventilatory pattern in the prevention of ventilator-associated pneumonia', *Critical Care Medicine*, 42(9), pp. e620-e627.

Li Bassi, G., Martí, J. D., Comaru, T., Aguilera-Xiol, E., Rigol, M., Ntoumenopoulos, G., Terraneo, S., De Rosa, F., Rinaudo, M., Fernandez-Barat, L., Battaglini, D., Meli, A., Ferrer, M., Pelosi, P., Chiumello, D. and Torres, A. (2019) 'Short-Term Appraisal of the Effects and Safety of Manual Versus Ventilator Hyperinflation in an Animal Model of Severe Pneumonia', *Respiratory care*, 64(7), pp. 760-770.

Li Bassi, G., Saucedo, L., Marti, J. D., Rigol, M., Esperatti, M., Luque, N., Ferrer, M., Gabarrus, A., Fernandez, L., Kolobow, T. and Torres, A. (2012) 'Effects of duty cycle and positive end-expiratory pressure on mucus clearance during mechanical ventilation*', *Critical Care Medicine*, 40(3), pp. 895-902.

Li Bassi, G. and Torres, A. (2011) 'Ventilator-associated pneumonia: role of positioning', *Curr Opin Crit Care*, 17(1), pp. 57-63.

Linnane, M. P., Caruana, L. R., Tronstad, O., Corley, A., Spooner, A. J., Barnett, A. G., Thomas, P. J. and Walsh, J. R. (2019) 'A comparison of the effects of manual hyperinflation and ventilator hyperinflation on restoring end-expiratory lung volume after endotracheal suctioning: A pilot physiologic study', *J Crit Care*, 49, pp. 77-83.

Longhini, F., Bruni, A., Garofalo, E., Ronco, C., Gusmano, A., Cammarota, G., Pasin, L., Frigerio, P., Chiumello, D. and Navalesi, P. (2020) 'Chest physiotherapy improves lung aeration in hypersecretive critically ill patients: a pilot randomized physiological study', *Crit Care*, 24(1), p. 479.

Lottering, M. and Van Aswegen, H. (2016) 'Physiotherapy practice in South African intensive care units', *Southern African Journal of Critical Care*, 32(1), pp. 11-16.

Mahajan, M., DiStefano, D., Satalin, J., Andrews, P., al-Khalisy, H., Baker, S., Gatto, L., Nieman, G. and Habashi, N. (2019) 'Time-controlled adaptive ventilation (TCAV) accelerates simulated mucus clearance via increased expiratory flow rate', *Intensive Care Medicine Experimental*, 7(27), pp. 1-11.

Makhabah, D. and Ambrosino, N. (2013) 'Airway Clearance in the Intensive Care Unit.', *European Medical Journal: Respiratory*, October, pp. 135-139.

Malbouisson LM, Muller JC, Constantin JM, et al. CT scan ARDS study group: computed tomography assessment of positive end-expiratory pressure-induced alveolar recruitment in patients with acute respiratory distress syndrome. *Am J Respir Crit Care Med* 2001; 163:1444-1450

Malekzadeh, J., Yazdani, M., Sedaghat, A., Mazlom, S. R., Khajebeyk A. P. (2016) 'Effect of Lung Manual Hyperinflation (MHI) on Oxygenation of Patients Following Abdominal Surgery and T-Tube Support', *Evidence Based Care Journal*, 6(3), pp. 55-66.

Marti, J. D., Ntoumenopoulos, G. and Torres, A. (2013) 'Physiotherapy in mechanically ventilated patients: Why and how', *Clinical Pulmonary Medicine*, 20(6), pp. 292-299.

Martins, L., da Silva Naue, W., Skueresky, A., Bianchi, T., Dias, A. and Forgiarini, L. (2019) 'Effects of Combined Tracheal Suctioning and Expiratory Pause: A Crossover Randomized Clinical Trial', *Indian J Crit Care Med.*, 23(10), pp. 454-7.

Martí, J. D., Li Bassi, G., Rigol, M., Saucedo, L., Ranzani, O. T., Esperatti, M., Luque, N., Ferrer, M., Vilaro, J., Kolobow, T. and Torres, A. (2013) 'Effects of manual rib cage compressions on expiratory flow and mucus clearance during mechanical ventilation', *Critical Care Medicine*, 41(3), pp. 850-856.

Martínez-Alejos, R., Martí, J. D., Li Bassi, G., Gonzalez-Anton, D., Pilar-Diaz, X., Reginault, T., Wibart, P., Ntoumenopoulos, G., Tronstad, O., Gabarrus, A., Quinart, A. and Torres, A. (2021) 'Effects of mechanical insufflation-exsufflation on sputum volume in mechanically ventilated critically ill subjects', *Respiratory Care*, 66(9), pp. 1371-1379.

Matilde, I. N. E., Eid, R. A. C., Nunes, A. F., Ambrozin, A. R. P., Moura, R. H., Carnieli-Cazati, D. and Timenetsky, K. T. (2018) 'Bronchial hygiene techniques in patients on mechanical ventilation: what are used and why?', *Einstein (Sao Paulo)*, 16(1), p. eAO3856.

Meawad, M., El Aziz, A., Obaya, H., Mohamed, S. and Mounir, K. (2018) 'Effect of Chest Physical Therapy Modalities on Oxygen Saturation and Partial Pressure of Arterial Oxygen in Mechanically Ventilated Patients', *The Egyptian Journal of Hospital Medicine*, 72(8), pp. 5005-5008.

Meyers, M. and Gonzales, J. (2019) 'Intrapulmonary percussive ventilation: An overview', *Neonatal Intensive Care*, 32(4), pp. 36-41.

Mohamed, A. and Saber, H. (2015) 'Oxygenation Effects of Manual Hyperinflation on Mechanically Ventilated Patients', *Medical Journal of Cairo University*, 81(1), pp. 633-63.

Mohamed, M., Hagag, A., Taha, M. and Ahmed, A. (2017) 'Chest Physiotherapy in Prevention of Hospital Acquired Pneumonia after Liver Transplantation', *Medical Journal of Cairo University*, 85(7), pp. 2761-8.

Moreira, F. C., Teixeira, C., Savi, A. and Xavier, R. (2015) 'Changes in respiratory mechanics during respiratory physiotherapy in mechanically ventilated patients', *Rev Bras Ter Intensiva*, 27(2), pp. 155-60.

Naue, W. D. S., Herve, B. B., Vieira, F. N., Deponti, G. N., Martins, L. F., Dias, A. S. and Vieira, S. R. R. (2019) 'Comparison of bronchial hygiene techniques in mechanically ventilated patients: a randomized clinical trial', *Rev Bras Ter Intensiva*, 31(1), pp. 39-46.

Naue, W. S., Junior, L. A. F., Dias, A. S. and Vieira, S. R. R. (2014) 'Chest compression with a higher level of pressure support ventilation: Effects on secretion removal, hemodynamics, and respiratory mechanics in patients on mechanical ventilation', *Jornal Brasileiro de Pneumologia*, 40(1), pp. 55-60.

Naue, W. S., Silva, A. C. T., Guntzel, A. M., Condessa, R. L., Olivera, R. P. and Vieira, S. R. (2011) 'Increasing pressure support does not enhance secretion clearance if applied during

manual chest wall vibration in intubated patients: a randomised trial', *Journal of Physiotherapy*, 57(1), pp. 21-26.

Newman, A. N. L., Gravesande, J., Rotella, S., Wu, S. S., Topp-Nguyen, N., Kho, M. E., Harris, J. E., Fox-Robichaud, A. and Solomon, P. (2018) 'Physiotherapy in the neurotrauma intensive care unit: A scoping review', *Journal of Critical Care*, 48, pp. 390-406.

Newstead, C. J., Seaton, J. A., Johnston, C. L. (2017) 'Australian critical care nursing professionals' attitudes towards the use of traditional "chest physiotherapy" techniques', *Hong Kong Physiotherapy Journal*, 36, pp. 33-48.

Ntoumenopoulos, G. (2014) 'Clinical impact of secretion retention', *Current Respiratory Medicine Reviews*, 10(3), pp. 158-162.

Ntoumenopoulos, G., Berry, M. and Camporota, L. (2014) 'Effects of manually-assisted cough combined with postural drainage, saline instillation and airway suctioning in critically-ill patients during high-frequency oscillatory ventilation: A prospective observational single centre trial', *Physiotherapy Theory and Practice*, 30(5), pp. 306-311.

Ntoumenopoulos, G., Berry, M. P., Camporota, L. and Lam, L. (2019) 'Indicators of Airway Secretion Weight in Mechanically Ventilated Subjects', *Respiratory care*, 64(11), pp. 1377-1386.

Ntoumenopoulos, G. and Glickman, Y. (2012) 'Computerised lung sound monitoring to assess effectiveness of chest physiotherapy and secretion removal: A feasibility study', *Physiotherapy (United Kingdom)*, 98(3), pp. 250-255.

Ntoumenopoulos, G., Hammond, N., Watts, N. R., Thompson, K., Hanlon, G., Paratz, J. D., Thomas, P. and For the George Institute for Global Health and the Australian and New Zealand Intensive Care Society Clinical Trials, G. (2018) 'Secretion clearance strategies in Australian and New Zealand Intensive Care Units', *Australian Critical Care*, 31(4), pp. 191-196.

Ntoumenopoulos, G., Shannon, H. and Main, E. (2011) 'Do commonly used ventilator settings for mechanically ventilated adults have the potential to embed secretions or promote clearance?', *Respiratory Care*, 56(12), pp. 1887-1892.

Oliveira, A. C. O., Lorena, D. M., Gomes, L. C., Amaral, B. L. R. and Volpe, M. S. (2019) 'Effects of manual chest compression on expiratory flow bias during the positive end-expiratory pressure-zero end-expiratory pressure maneuver in patients on mechanical ventilation', *J Bras Pneumol*, 45(3), p. e20180058.

Ouchi, A., Sakuramoto, H., Unoki, T., Yoshino, Y., Hosino, H., Koyama, Y., Enomoto, Y., Shimojo, N., Mizutani, T. and Inoue, Y. (2020) 'Effects of manual rib cage compressions on mucus clearance in mechanically ventilated pigs', *Respiratory Care*, 65(8), pp. 1135-1140.

Pathmanathan, N., Beaumont, N., Gratrix, A. (2015) 'Respiratory physiotherapy in the critical care unit', *Continuing Education in Anaesthesia Critical Care & Pain* 15(1), pp. 20-25.

Pattanshetty, R. B. and Gaude, G. S. (2011) 'Effect of multimodality chest physiotherapy on the rate of recovery and prevention of complications in patients with mechanical ventilation: A prospective study in medical and surgical intensive care units', *Indian Journal of Medical Sciences*, 65(5), pp. 175-185.

Paulus, F, Binnekade, J. M., Middelhoek, P., Vroom, M. B., Schultz M. J. (2014) 'Guideline implementation powered by feedback and education improves manual hyperinflation performance', *Nursing in Critical Care*, 21(1), pp. 36-43.

Paulus, F., Binnekade, J. M., Vroom, M. B. and Schultz, M. J. (2012) 'Benefits and risks of manual hyperinflation in intubated and mechanically ventilated intensive care unit patients: a systematic review', *Crit Care*, 16(4), p. R145.

Paulus, F., Veelo, D. P., de Nijs, S. B., Beenen, L. F., Bresser, P., de Mol, B. A., Binnekade, J. M. and Schultz, M. J. (2011) 'Manual hyperinflation partly prevents reductions of functional residual capacity in cardiac surgical patients-a randomized controlled trial', *Crit Care*, 15(4), p. R187.

Peters MDJ, Godfrey C, McInerney P, Munn Z, Tricco AC, Khalil, H. Chapter 11: Scoping Reviews (2020 version). In: Aromataris E, Munn Z (Editors). *JB I Manual for Evidence Synthesis*, JBI, 2020

Piccin, V. S., Calciolari, C., Yoshizaki, K., Gomes, S., Albertini-Yagi, C., Dolhnikoff, M., Macchione, M., Caldini, E. G., Saldiva, P. H. and Negri, E. M. (2011) 'Effects of different mechanical ventilation strategies on the mucociliary system', *Intensive Care Med*, 37(1), pp. 132-40.

Pozuelo-Carrascosa, D. P., Torres-Costoso, A., Alvarez-Bueno, C., Cavero-Redondo, I., López Muñoz, P. and Martínez-Vizcaíno, V. (2018) 'Multimodality respiratory physiotherapy reduces mortality but may not prevent ventilator-associated pneumonia or reduce length of stay in the intensive care unit: a systematic review', *Journal of Physiotherapy*, 64(4), pp. 222-228.

Raafat, A. and Elbasiouny, H. (2011) 'Arterial oxygenation response to manual hyperinflation as an added procedure to chest physiotherapy in critically ill mechanically ventilated patients', *Journal of American Science*, 7(12), pp. 585-90.

Ribeiro, B. S., Lopes, A. J., Menezes, S. L. S. and Guimarães, F. S. (2019) 'Selecting the best ventilator hyperinflation technique based on physiologic markers: A randomized controlled crossover study', *Heart and Lung*, 48(1), pp. 39-45.

Rose, L., McKim, D., Leasa, D., Nonoyama, M., Tandon, A., Kaminska, M., O'Connell, C., Loewen, A., Connolly, B., Murphy, P., Hart, N., Road, J. (2018) 'Monitoring Cough Effectiveness and Use of Airway Clearance Strategies: A Canadian and UK Survey', *Respiratory Care*, 63(12), pp. 1506-13.

Rose, L., Adhikari, N. K., Leasa, D., Fergusson, D. A. and McKim, D. (2017) 'Cough augmentation techniques for extubation or weaning critically ill patients from mechanical ventilation', *Cochrane Database Syst Rev*, 1(1), p. Cd011833.

Rose, L., Adhikari, N. K., Poon, J., Leasa, D. and McKim, D. A. (2016) 'Cough augmentation techniques in the critically ill: A Canadian national survey', *Respiratory Care*, 61(10), pp. 1360-1368.

Sharma, P., Prem, V. and Jain, S. (2018) 'Immediate effects of acapella® on dynamic lung compliance in mechanically ventilated patients with acute respiratory distress syndrome: A case series', *Indian Journal of Critical Care Medicine*, 22(2), pp. 100-102.

Shetty, S., Alaparathi, G. K., Shyam Krishnan, K., Upadya, M., Prabhakar, A., Bairapareddy, K. C. and Amaravadi, S. K. (2020) 'Comparison of head-down tilt versus flat position on mucus clearance and respiratory mechanics in mechanically ventilated patients: A randomized crossover trial', *Critical Reviews in Physical and Rehabilitation Medicine*, 32(1), pp. 23-38.

Shi, Z. H., Jonkman, A., de Vries, H., Jansen, D., Ottenheijm, C., Girbes, A., Spoelstra-de Man, A., Zhou, J. X., Brochard, L. and Heunks, L. (2019) 'Expiratory muscle dysfunction in critically ill patients: towards improved understanding', *Intensive Care Medicine*, 45(8), pp. 1061-1071.

Skinner, E. H., Haines, K. J., Berney, S., Warrillow, S., Harrold, M. and Denehy, L. (2015) 'Usual care physiotherapy during acute hospitalization in subjects admitted to the ICU: An observational cohort study', *Respiratory Care*, 60(10), pp. 1476-1485.

Skinner, E. H., Thomas, P., Reeve, J. C. and Patman, S. (2016) 'Minimum standards of clinical practice for physiotherapists working in critical care settings in Australia and New Zealand: A modified Delphi technique', *Physiotherapy Theory and Practice*, 32(6), pp. 468-482.

Spapen, H., Borremans, M., Diltoer, M., De Regt, J., Bruggemans, C. and Honoré, P. M. (2016) 'Intrapulmonary percussion with autogenic drainage and ventilator-associated gram-negative infection: A pilot study', *Netherlands Journal of Critical Care*, 24(2), pp. 6-10.

Spapen, H. D., Regt, J. D. and Honoré, P. M. (2017) 'Chest physiotherapy in mechanically ventilated patients without pneumonia-a narrative review', *Journal of Thoracic Disease*, 9(1), pp. E44-E49.

Stiller, K. (2013) 'Physiotherapy in intensive care. An updated systematic review', *Chest*, 144(3), pp. 825-847.

Stilma, W., van der Hoeven, S. M., Scholte Op Reimer, W. J. M., Schultz, M. J., Rose, L. and Paulus, F. (2021) 'Airway care interventions for invasively ventilated critically ill adults—A Dutch national survey', *Journal of Clinical Medicine*, 10(15).

Strickland, S. L., Rubin, B. K., Haas, C. F., Volsko, T. A., Drescher, G. S. and O'Malley, C. A. (2015) 'AARC Clinical Practice Guideline: Effectiveness of Pharmacologic Airway Clearance Therapies in Hospitalized Patients', *Respir Care*, 60(7), pp. 1071-7.

Suh, M., Heitkemper, M. and Smi, C. K. (2011) 'Chest Physiotherapy on the respiratory mechanics and elimination of sputum in paralyzed and mechanically ventilated patients with acute lung injury: A pilot study', *Asian Nursing Research*, 5(1), pp. 60-69.

Swingwood, E., Stilma, W., Tume, L., Cramp, F., Paulus, F., Schultz, M., Scholte op Reimer, W. and Rose, L. (2020) 'The use of mechanical insufflation-exsufflation in invasively ventilated critically ill adults: a scoping review protocol', *Systematic Reviews*, 9(1).

Terzi, N., Guerin, C. and Gonçalves, M. R. (2019) 'What's new in management and clearing of airway secretions in ICU patients? It is time to focus on cough augmentation', *Intensive Care Med*, 45(6), pp. 865-868.

Thomas, P., J. (2015) 'The effect of mechanical ventilator settings during ventilator hyperinflation techniques: a bench-top analysis', *Anaesth Intensive Care*, 43(1), pp. 81-7.

Tomar, G. S., Singh, G. P., Bithal, P., Upadhyay, A. D. and Chaturvedi, A. (2019) 'Comparison of Effects of Manual and Mechanical Airway Clearance Techniques on Intracranial Pressure in Patients With Severe Traumatic Brain Injury on a Ventilator: Randomized, Crossover Trial', *Phys Ther*, 99(4), pp. 388-395.

Tricco, AC, Lillie, E, Zarin, W, O'Brien, KK, Colquhoun, H, Levac, D, Moher, D, Peters, MD, Horsley, T, Weeks, L, Hempel, S et al. PRISMA extension for scoping reviews (PRISMA-ScR): checklist and explanation. *Ann Intern Med*. 2018;169(7):467-473

Twose, P., Jones, U. and Cornell, G. (2019) 'Minimum standards of clinical practice for physiotherapists working in critical care settings in the United Kingdom: A modified Delphi technique', *Journal of the Intensive Care Society*, 20(2), pp. 118-131.

van Aswegen, H., van Aswegen, A., Raan, H. D., Toit, R. D., Spruyt, M., Nel, R. and Maleka, M. (2013) 'Airflow distribution with manual hyperinflation as assessed through gamma camera imaging: A crossover randomised trial', *Physiotherapy (United Kingdom)*, 99(2), pp. 107-112.

van der Lee, L. Hill, A., M., Patman, S. (2018) 'After-hours respiratory physiotherapy for intubated and mechanically ventilated patients with community-acquired pneumonia: An Australian perspective', *Aust Crit Care*, 31(6), pp. 349-354.

van der Lee, L., Hill, A. M., Patman, S. (2020) 'Clinical validation of expert consensus statements for respiratory physiotherapy management of invasively ventilated adults with community-acquired pneumonia: A qualitative study', *Intensive and Critical Care Nursing*, 60, pp. 1-10.

van der Lee, L., Hill, A. M. and Patman, S. (2017) 'Efficacy of a respiratory physiotherapy intervention for intubated and mechanically ventilated adults with community acquired pneumonia: A systematic review protocol', *JBIC Database of Systematic Reviews and Implementation Reports*, 15(6), pp. 1508-1511.

van der Lee, L., Hill, A. M. and Patman, S. (2019) 'Expert consensus for respiratory physiotherapy management of mechanically ventilated adults with community-acquired pneumonia: A Delphi study', *J Eval Clin Pract*, 25(2), pp. 230-243.

Volpe, M. S., Guimarães, F. S., Morais, C. A. (2020) 'Airway Clearance Techniques for Mechanically Ventilated Patients: Insights for Optimization', *Respiratory Care*, 65(8), pp. 1174-88.

Volpe, M. S., Naves, J. M., Ribeiro, G. G., Ruas, G., Amato, M. B. P. (2018) 'Airway clearance with an optimized mechanical insufflation-exsufflation maneuver', *Respiratory Care*, 63(10), pp. 1214-1222.

Wang, M. Y., Pan, L. and Hu, X. J. (2019) 'Chest physiotherapy for the prevention of ventilator-associated pneumonia: A meta-analysis', *American Journal of Infection Control*, 47(7), pp. 755-760.

Wang, T. H., Wu, C. P. and Wang, L. Y. (2018) 'Chest physiotherapy with early mobilization may improve extubation outcome in critically ill patients in the intensive care units', *Clinical Respiratory Journal*, 12(11), pp. 2613-2621.

Waqas, M., Malik, A. and Javed, M. (2014) 'Effectiveness of conventional chest physiotherapy versus manual hyperinflation during postural drainage of ventilated COPD patients', *Rawal Medical Journal*, 39(1), pp. 32-34.

Yousefnia-Darzi, F., Hasavari, F., Khaleghdoost, T., Kazemnezhad-Leyli, E. and Khalili, M. (2016) 'Effects of thoracic squeezing on airway secretion removal in mechanically ventilated patients', *Iranian Journal of Nursing and Midwifery Research*, 21(3), pp. 337-342.