

What sources of bereavement support are perceived helpful by bereaved people and why? Empirical evidence for the compassionate communities approach

Samar M Aoun^{1,2} , Lauren J Breen³, Ishta White³,
Bruce Rumbold¹ and Allan Kellehear⁴

Palliative Medicine
2018, Vol. 32(8) 1378–1388
© The Author(s) 2018



Reprints and permissions:
sagepub.co.uk/journalsPermissions.nav
DOI: 10.1177/0269216318774995
journals.sagepub.com/home/pmj



Abstract

Aims: To determine who provides bereavement support in the community, what sources are perceived to be the most or least helpful and for what reason, and to identify the empirical elements for optimal support in developing any future compassionate communities approach in palliative care.

Design: A population-based cross-sectional investigation of bereavement experiences. Sources of support (informal, community and professional) were categorised according to the Public Health Model of Bereavement Support; most helpful reasons were categorised using the Social Provisions Scale, and least helpful were analysed using inductive content analysis.

Setting and participants: Bereaved people were recruited from databases of funeral providers in Australia via an anonymous postal survey (2013–2014).

Results: In total, 678 bereaved people responded to the survey. The most frequently used sources of support were in the informal category such as family, friends and funeral providers. While the professional category sources were the least used, they had the highest proportions of perceived unhelpfulness whereas the lowest proportions of unhelpfulness were in the informal category. The functional types of helpful support were Attachment, Reliable Alliance, Social Integration and Guidance. The five themes for least helpful support were: Insensitivity, Absence of Anticipated Support, Poor Advice, Lack of Empathy and Systemic Hindrance.

Conclusion: A public health approach, as exemplified by compassionate communities policies and practices, should be adopted to support the majority of bereaved people as much of this support is already provided in informal and other community settings by a range of people already involved in the everyday lives of those recently bereaved. This study has provided further support for the need to strengthen the compassionate communities approach, not only for end of life care for dying patients but also along the continuum of bereavement support.

Keywords

Public health, social support, capacity building, bereavement support, palliative care

What is already known about the topic?

- The majority of bereaved people rely on family and friends for bereavement support.
- There is little evidence as to how this support is provided and the reasons for its perceived helpfulness.
- Consequently, there is little indication about which particular support strategies are optimal for building a community's capacity to provide the type of support advocated by the compassionate communities approach now widely advocated by palliative care services.

¹Palliative Care Unit, Department of Public Health, School of Psychology and Public Health, La Trobe University, Melbourne, VIC, Australia

²Institute for Health Research, Notre Dame University, Fremantle, WA, Australia.

³School of Psychology, Faculty of Health Sciences, Curtin University, Perth, WA, Australia

⁴School of Nursing and Healthcare Leadership, Faculty of Health Studies, University of Bradford, Bradford, UK

Corresponding author:

Samar M Aoun, Palliative Care Unit, School of Psychology and Public Health, La Trobe University, Kingsbury Drive, Bundoora, 3086 Victoria, Australia.

Email: s.aoun@latrobe.edu.au

What this paper adds?

- While the professional sources were the least used, they had the highest proportions of perceived unhelpfulness.
- The types of informal social support most valued as helpful were the emotional bonds (attachment), the practical assistance (tangible alliance) and the perceived sense of belonging (social integration).
- A public health approach to bereavement care is needed to support 'everyday assets' in the community who care for the majority of the bereaved, without the overreach from professional services.
- The compassionate communities approach, so far focussed on care of the dying, is well and truly operational in the bereavement phase.

Implications for practice, theory or policy

- This study provides strong empirical indicators and directions that encourage a compassionate community approach to future bereavement support practice and policy.
- Findings provide guidance as to operationalising how to offer professionalised services to those who would most benefit, while not providing these services unnecessarily to those who are supported within their natural support networks.
- More research is needed in identifying a range of useful practice models that connect health and social care services with local communities and build upon local helping networks.

Introduction

Social support is an important factor influencing bereavement outcomes.^{1,2} Despite social support being the protective factor that is most amenable to intervention, it has received limited attention in the literature.^{3,4} Albrecht and Adelman⁵ describe social support as 'verbal and non-verbal communication between recipients and providers that reduces uncertainty about the situation, the self, the other, or the relationship, and functions to enhance a perception of personal control in one's life experiences' (p. 19). Research suggests that, following the loss of a relative or friend, social support can act as a buffer against psychological morbidity by reducing grief symptomatology and facilitating psychological adaptation.^{3,6} Importantly, it is also one of the few factors influencing the grieving process that can be modified and targeted during treatment to reduce the effects of those risk factors less modifiable, such as gender and personality features.^{2,7} Despite this, there is little empirical evidence supporting the methods employed to build the community's capacity to provide this type of support. While many studies have identified that the majority of bereaved people rely on family and friends as providers of support, there are limited studies, in number and methodological rigour, examining in detail the ways in which this support is provided and the reasons behind its perceived helpfulness.^{4,8–11}

Building the community capacity to support death, dying and loss is an important public health issue in palliative care. Central to any public health approach focussed on bereavement care in this particular context is the development of community supports and interventions that address the multiple morbidities (e.g. depression, job loss or social isolation) and mortalities (e.g. suicide or sudden death) associated with bereavement and that may be amenable to prevention, harm reduction and early intervention.

'Compassionate Communities/Cities' approaches are community development strategies, now widely discussed in recent palliative care literature, that attempt to build these kinds of supports and interventions to address the social epidemiology of grief and loss.^{12–14} These models of community support rely on identifying and developing local caring networks around the dying person and their family. Compassionate communities normalise seeking and accepting help from family, friends, neighbours, schools and workplaces for emotional and social support. In health promotion terms, support for grieving people is not only the task for health and social services but becomes everyone's responsibility.¹⁵ This is also one of six key recommendations of the recent UK palliative care framework for local action 2015–2020 – 'Ambition 6: Each community is prepared to help'.¹⁶

Qualitative studies on social support have highlighted its importance in adapting to life in the wake of loss via the death of a significant other. They indicated that connecting with bereaved others and experiencing compassion from support providers are two of the most helpful forms of social support reported by participants.^{17–19} The few quantitative studies that have examined different forms of social support and their relationship to bereavement outcomes have shown that people who would most likely benefit from support might be least likely to receive it.^{20,21}

Weis' model of social provision affords perhaps the most comprehensive assessment of perceived social support by offering a clear conceptualisation of six separate functions of social support.²² The Social Provisions Scale (SPS) was developed based on Weis' model and taps into these following six types of social support provisions: 'Attachment' refers to having an emotional and affectionate bond with another person, thus providing a sense of security; 'Social

Integration' refers to the perceived sense of belonging to a group that shares similar interests/concerns; 'Guidance' refers to the perceived availability of someone one can turn to for advice or information; 'Reliable Alliance' refers to the perceived availability of someone to rely on for tangible assistance; 'Reassurance of Worth' refers to the belief that one is valued by others; and 'Opportunity for Nurturance' refers to the belief that one is needed by other people.²³

However, no studies have examined the perceived social support in bereavement in a systematic way with the use of the SPS as a framework guiding the classification of functional types of social support.²³ Also no studies have examined the interaction between source and functional type of support in a population-based bereaved sample. Knowledge of bereavement based just on clinical encounters has resulted in many service providers taking an undifferentiated approach to bereavement care, despite research suggesting that this type of early intervention by professionals may not be effective.^{24,25} Rumbold and Aoun¹⁰ conclude that the research community, clinical practitioners and support services are well informed by the minority who seek support through health professionals; however, little is known about how bereavement is lived out in everyday life by the majority of those who do not seek these services.

In their public health model of bereavement support which advocates for partnerships between the formal and informal networks, Aoun et al.⁸ used a population-based survey to investigate the range of support received overall from professional caregivers and community sources. They classified sources of bereavement support into three: informal support, community support and professional support. Informal support included support offered by family, friends, funeral directors, financial or legal advisors, religious or spiritual advisors and the Internet or literature. Community support included support offered by General Practitioners (GPs), nursing homes, hospitals, pharmacists, community groups, palliative care providers or school-based advisors. Professional support included support offered by trained counsellors, bereavement support groups, social workers, case coordinators, psychologists and psychiatrists.

Objectives

The overall aim of this study is to determine who provides bereavement support in the community, what sources are perceived to be the most or least helpful and for what reason. Therefore, the three specific objectives were

- To identify the nature and extent of the sources of bereavement support, as outlined in the public health model of bereavement support, whether perceived helpful or unhelpful;
- To identify the nature and extent of the functional types of bereavement support, using the SPS as the guiding theoretical framework;

- To explore the empirical evidence for supporting the compassionate communities approach in bereavement support.

Methods

Ethics approval was granted by Curtin University Research Ethics Committee (HR 57/2012). We undertook the quantitative and qualitative analyses of cross-sectional questionnaire data pertaining to helpful and unhelpful sources and types of social support accessed by a population-based sample of bereaved people.

Study design

The study is a population-based cross-sectional investigation of bereavement experiences. A postal survey was used to collect information from clients of four funeral providers in Australia, specifically from metropolitan and regional areas of Western Australia and Victoria (2013–2014), 6–24 months after the death of their family member. Funeral providers were used as it was not possible to recruit through the Death Registry in Australia.

Participants and procedures

A total of 3190 study packages were delivered to the four funeral providers. These packages contained an invitation letter addressed from the funeral provider to the family, information sheet, the questionnaire, a list of support services for the family to use in case the respondent became distressed while completing the questionnaire and a reply-paid envelope. The funeral providers then selected all clients who were bereaved 6–24 months ago from their databases, attached names and address labels on the envelopes and mailed the study packages. Consent was implied by the return of the completed survey. No reminder letter was sent as it was felt to be too intrusive on the bereaved families. Clients were eligible to participate in the study if they had been bereaved by a close family member or friend in the specified time frame, were able to read, understand and write in English and were over 18 years of age.

Measures and analyses

The questionnaire was developed in consultation with a reference group comprising representatives of the funeral industry, bereavement counsellors, palliative care services, primary care and community-based services. It has eight sections with a total of 82, predominantly closed, questions. The questionnaire was designed to gather information on participant demographics, the supports they accessed, those supports they would have liked to have been able to access, their perceived support needs and the extent to which they felt these needs had been met.^{8,26} The current study focussed

predominantly on the survey section pertaining to the perceived helpfulness of bereavement supports the individual accessed. This section asked the bereaved respondent to list in order, three of the most helpful or unhelpful sources of support accessed, and to state why they were the most or least helpful. To ensure quality and rigour of the coding process, two authors independently blind-coded a random 10% of responses. These codes were compared, with differences discussed until consensus was reached and the remaining responses were coded by one author.

The question on helpful support was as follows: 'Please list up to 3 of the most helpful sources of support that you accessed and reasons why they were the most helpful'. Spaces were provided for free descriptions on the questionnaire. Responses to the first part of the question were categorised according to the three sources of support identified through the Public Health Model of Bereavement Support: informal support, community support and professional support.⁸ Descriptive statistics for variables were calculated using IBM SPSS Statistics Version 24. Responses to the second part of the question were categorised using the SPS as an initial framework for the identification of different types of functional support.²³ The SPS has undergone considerable psychometric evaluation with multiple studies demonstrating that the measure has good internal consistency, test-retest reliability and convergent and discriminant validity.^{23,27}

Similarly, the question on unhelpful support was as follows: 'Please list up to 3 of the least helpful sources of support that you accessed and reasons why they were the least helpful'. As there is no framework for categorising non- or unhelpful support, responses to the second part of the question were categorised using an inductive approach to coding whereby the data were subject to open content analysis and five themes were generated: 'Insensitivity' is related to rude or insensitive remarks, unexpected/inappropriate behaviour or behaviour that was incongruent with the situation; 'Absence of Anticipated Support' is support that was expected to be provided or available to the bereaved, but it did not materialise; 'Poor Advice' included unwanted advice or interference; 'Systemic Hindrance' is the system putting up obstacles and 'Lack of Empathy' included indifference and lack of interest.²⁸

Results

Profile of the bereaved and deceased

In total, 678 bereaved people responded to the survey (21.3% response rate). In all, 71% of the bereaved people who responded were female, mean age of 62.4 years (standard deviation (SD)=12.2); 49.6% were married and 37.0% were widowed; 36.8% were the spouse of the deceased and 45.3% were the son or daughter of the deceased; 69.5% were Australian citizens; 45.2% had finished high school,

Table 1. Characteristics of the Bereaved.

Bereaved	Total, n = 678	
	n	%
Gender		
Male	194	28.8
Female	479	71.2
Age (years)		
Mean (SD)	62.4	(12.2)
Median (range)	62.0	(20–96)
Marital status		
Single/never married	25	3.7
Married/de facto	336	49.6
Separated/divorced	59	8.7
Widowed	251	37.0
Cultural background		
Australian–non-Aboriginal	471	69.5
Australian–Aboriginal	4	0.6
Other English speaking	125	18.5
Non-English speaking	66	9.7
Highest level of education		
No formal education	3	0.4
Primary school	28	4.2
High school	304	45.2
Diploma/certificate/trade qualification	205	30.5
University degree	132	19.6
Main employment		
Paid employment	247	36.9
Retired	260	38.9
Disabled	9	1.3
Household duties	92	13.8
Unemployed	15	2.2
Other	46	6.8
Relationship to the deceased		
Spouse/partner	249	36.8
Parent	48	7.1
Sibling	25	3.7
Daughter/son	307	45.3
Other relative	32	4.7
Friend	13	1.9
Other	3	0.4
Period of bereavement: mean (SD) in months	14.3	(6.28)

SD: standard deviation.

30.5% had a diploma or trade qualification and 19.6% had a university degree; 36.9% were currently employed and 38.9% were retired. The mean period of bereavement for respondents was 14.3 months (SD=6.3; Table 1).

The mean age of the deceased was 75.4 years (SD=18.3); about half of the deceased were female; 68% of the deaths were due to life limiting illnesses (mainly cancer 30%, heart disease/organ failure 18% and dementia/old age 15%) and 54% of these care recipients used palliative care services; their death occurred in hospital




Figure 1. Perceived helpfulness of sources of bereavement support.

(39%), nursing home (32%), home (18%) and hospice (9%).

Perceived helpfulness

The most frequently used sources of support were those in the informal category such as family, friends and funeral providers (80%–90%), followed by GPs from the community category (58%) (Figure 1). While the professional category sources were the least used, they had the highest proportions of perceived unhelpfulness (Table 2): nearly half of those who used psychiatrists found them unhelpful (46%), and over a third of respondents who used the following services rated them unhelpful – bereavement support group (41%), case coordinator (39%), social worker (38%) and school-based advisor (36%). Similar high rates of unhelpfulness were reported in the community category and in particular a community group (40%), community pharmacist (39%) and palliative care service (33%). By contrast, the lowest proportions of unhelpfulness were in the informal category where only 8% found family unhelpful, followed by the funeral provider (8.7%) and friends (11.7%) (Table 2).

Types of helpful bereavement support

About 80% of respondents provided at least one reason for experiencing helpful support. Using the SPS framework, the concepts ‘reassurance of worth’ and ‘opportunity for

Table 2. Perceived helpfulness of each source of bereavement support (n = 678).

	Very/quite helpful		A little/not helpful	
	n	%	n	%
Family	585	92.00	51	8.00
Funeral provider	491	91.30	47	8.70
Friends	526	88.30	70	11.70
Psychologist	33	78.60	9	21.40
Religious or spiritual advisor	179	78.50	49	21.50
Counsellor	50	76.90	15	23.10
Nursing home	129	76.30	40	23.70
Financial or legal	182	72.50	69	27.50
GP	270	70.30	114	29.70
Hospital	162	70.10	69	29.90
Palliative care service	106	66.70	53	33.30
School-based advisor	14	63.60	8	36.40
Social worker	34	61.80	21	38.20
Community pharmacist	116	61.10	74	38.90
Case coordinator	22	61.10	14	38.90
Community group	104	60.10	69	39.90
Internet or other literature	95	59.70	64	40.30
Bereavement support group	30	58.80	21	41.20
Psychiatrist	7	53.80	6	46.20

Those who did not answer the question or who indicated ‘not applicable’ were removed from the denominator for each support source variable.

Table 3. Source and corresponding type of functional support endorsed as helpful.

Source and functional type of support	Quotes	Frequency endorsed
Informal		
Attachment	'Family, were there for me offering help and support. They listened to how I felt about losing Mum and comforted me'. 'Friends, they sensed my grief and stood alongside me'. 'Funeral provider [name], his wonderful calmness, and kindness, and sense of humour (he needed it with our family) exceptional!'	416
Social integration	'Workmates, they made me get out and about'. 'Family, we all shared the same loss and supported each other'. 'My colleagues who shared their stories of experience of suicide in their families that made me feel less alone in my grief and other mothers at the cemetery who lost loved ones in similar circumstances'.	166
Guidance	'Funeral directors, their advice was invaluable and always right'. 'Close friends, offered support and listened to me with advice'. 'Bank, helped and guided me through the process of consolidating finances and paying accounts'.	97
Reliable alliance	'Family, helped with preparation of funeral with meals'. 'Friends, helped to weather-proof the patio for funeral services'. 'Cousin of late husband, lovely, comforting, organised getting the car home as I am not a driver'.	234
Community		
Attachment	'[area] Service Club, compassionate and caring. Phone calls, notes in the mail – checking on well-being'. 'GP, would attend or phone 2x day, took on my concerns'. '[Health and community service], were caring and compassionate'.	84
Social integration	'[Community support group], because these people going exactly through the same issues and know what we are talking about'. 'Nursing home, carers attended the funeral. We weren't alone as the mourners'. 'Hospice, I get regular letters inviting me to afternoon teas and get together'.	31
Guidance	'[Health and community service] were main source of information'. 'Hospital staff, Informed us what steps to take next and what documents we would need'. 'Hospice, kind information of illness and how to handle oncoming death'.	57
Reliable alliance	'Palliative care, the nurse we contacted when my wife was critical was efficient and very quick in getting the job done'. 'Community club, helped organise the wake. Suspended normal activities for that day at club'. '[Religious community service], they gave caring support in health. They arrived, came everyday, morning and night towards the end. They had equipment and contacts and experience'.	76
Professional		
Attachment	'Psychologist, impartial third party to talk to, offload'. 'Counselling, be able to talk to a person who really listens'. 'Social worker, became close to her along the journey'.	20
Social integration	'Bereavement support group, able to connect to others that had lost their partner'. 'Trauma and grief counsellor, because they are qualified people with empathetic behavior and know mostly how to deal with these situations appropriately, also some have their own experience about loss from a loved one and they can very well relate to your pain and suffering'. '[Community support service], support, counselling, meeting parents like us'.	7
Guidance	'Victim support services; They were there to give advice and helped through each stage as it came. I did not know this service existed'. 'Bereavement support group, they listened and were experts in this field. Gave me paths to follow and offered solutions to dealing with the pain and so-called friends and family issues'. 'Psychologist, gave us tools and options with coping with the loss'.	18
Reliable alliance	NA	0

nurturance' were not included as none of the responses aligned with these concepts in this particular sample. A summary table, providing examples of the codes generated and the frequency with which each functional type of support was endorsed by the sample, is shown (Table 3).

In the informal sources of support, attachment was the form of social support most frequently endorsed as

helpful by respondents, followed by reliable alliance and social integration. Guidance came last in the informal sources but became more important in the other two sources. Within the community sources of support, attachment and reliable alliance were still prominent followed by guidance, while in the professional sources, guidance was as frequent as attachment (Table 3).

Table 4. Source and corresponding type of support endorsed as unhelpful.

Source and reasons for lack of Support	Quotes	Frequency endorsed
Informal		
Absence	(Family) 'All came over next day then disappeared – too much like “social gathering” wasn't helpful'. (Friends) 'They were not there when I needed them. I think they did not know what to do'. (Friends) 'Offers of assistance that were not followed up'.	31
Insensitivity	(Acquaintances) 'Glib and off-hand comments'. (Government Dept) 'Rude staff not interested in how we were feeling'. (Workplace) 'Told just do my job'.	25
Poor advice	(Government Dept) 'Poor advice and assistance about the correct forms to fill in and the implications'. (Pastor) 'I didn't want to be hugged all the time but he would say things about God that made me feel worse'. (Lawyer) 'Options were not explained or presented adequately. We felt the advice given was not the only option afterward'.	7
Lack of empathy	(Acting manager) 'No compassion. Threatened to sack me'. (Financial Institutions) 'Just wanting their money on outstanding loan of the deceased. No compassion during/after sorting the estate out'.	4
Systemic hindrance	(Probate office) 'Even though the will and death certificate were all in order, they threw up numerous obstacles to make the finalisation of the will difficult'. (Coronial Investigation Unit) 'By law they say that they can't tell you anything, refer you to Coroner'. (Superannuation companies) 'Red tape and bullshit I had to jump through to make claim on super/death benefit policy'.	27
Community		
Absence	(General practitioner) 'They never made contact, not once'. (Nursing home) 'Never contacted me again after my Mum passed away'. (Palliative care service) 'No contact was made to offer sympathy or help. I had messaged the two people to advise of my brother-in-law's passing and got no response'.	30
Insensitivity	(Nursing home) 'They were insensitive. It's a business for them'. (Nursing home) 'Only concerned to clear Mum's room next morning – no comment from them'. (Police) 'They treated me like a criminal'.	16
Poor advice	(General practitioner) 'Gave my tablets which I did not take (for depression)'. (Doctors) 'Didn't really help or suggest counselling etc'.	3
Lack of empathy	(General practitioner) 'Could not demonstrate any empathy or care for our loss'. (Pharmacist) 'Not interested in situation'. (Police) 'Closed off with information and empathy'.	8
Systemic hindrance	(Emergency department) 'Triage nurse didn't have a system to recognise an imminent admission'. (Police) 'Tried to stop me seeing my son where he died and when I tried to access more information, their response was inadequate'. (Not for profit organisation) 'Not able to help my nephews – only if parents and them had been in a separated relationship'.	12
Professional		
Absence	(Bereavement support) 'Not a word from them or anyone'. (Hospital caseworker) 'Never heard from them'.	5
Insensitivity	(Counsellor) 'Was always too busy to talk to me and she was always 'running out of time'.	2
Poor advice	(Psychologist) 'On returning [home] after funeral, I was told to go and "sit in the sun and pat the dog"'!	1
Lack of empathy	(Lifeline) 'Totally cold to my grief'.	1
Systemic hindrance	(Psychologist used by wife) 'Not helpful appeared to hold back recovery'	1

Types of unhelpful bereavement support

Fewer respondents provided comments on unhelpful sources compared to helpful sources (23.6% vs 78.6%). Table 4 provides examples of the frequency with which each of the five themes (Absence, Insensitivity, Poor

Advice, Lack of Empathy and Systemic Hindrance) was endorsed by the sample.

Absence (mainly by family/friends), insensitivity and systemic hindrance (mainly by financial and legal entities) were the most prominent themes in the informal sources. Similarly in the community sources, absence and

insensitivity from doctors, health services and nursing homes were highlighted (Table 4).

Discussion

Compared to existing literature, this study presents a more nuanced account of the role of both sources and functional types of social support for bereavement, in terms of the reasons why the supports accessed by a population-based sample of bereaved people were perceived as helpful or unhelpful. While about 90% of respondents listed family and friends as providers of support, we had previously lacked detail about what made this source of support helpful. This is almost inevitable in surveys where it is easier to be specific in listing professional care providers than the myriad of small contributions that come from friends, family and neighbours.¹⁰ Certainly, researchers who studied informal networks of care for people dying in the community found they had to persist to identify the contribution made by informal caregivers.²⁹

As expected, the most frequently used sources of support were those in the informal category, such as family and friends, and were deemed mostly helpful.^{8,11} What is unexpected is that funeral providers came next as most helpful and this is worthy of further investigation because of the valuable role they can play as part of the existing networks. While the professional sources were the least used, they had the highest proportions of perceived unhelpfulness (33%–46%). In particular, a third did not find palliative care services helpful and this is highlighted in the latest findings from this survey showing that just half of the bereaved had a follow-up contact from the palliative care services at 3–6 weeks, and only a quarter had a follow-up at 6 months, and that the blanket approach to bereavement support adopted by the services was deemed unhelpful.³⁰

Most health professionals have little exposure to evidence-based grief information and some draw upon outdated and ineffective models when working with bereaved clients.^{31–34} The higher levels of professionals reported as unhelpful could also reflect higher expectations of the care they should offer. But evidence suggesting that inappropriate referrals for intervention may lead to worse outcomes in bereavement should also be remembered.^{25,35} Early, particularly unrequested, referrals to a counsellor have the potential to problematise grief and compromise the support that could be offered by existing networks – grief is seen as a problem demanding professional intervention.

Using the SPS framework, the concepts exhibited by this study sample were as follows: attachment and social integration which fall in the non-assistance related functions of social ties, while reliable alliance and guidance fall in the assistance-related functions.²⁷ In the informal sources of support, the findings revealed that attachment (emotional or affectionate bond, often provided by spouse/partner or close family) was the form of social support most frequently endorsed as helpful by participants,

suggesting that the majority of bereaved individuals value this relational bond and emotional support from those closest to them. This was followed in terms of frequency by reliable alliance (tangible assistance such as providing meals and transport often offered by family members) and social integration (perceived sense of belonging often provided by friends). These findings are supported by previous research, which found that satisfaction with practical support in the months following loss predicted favourable bereavement outcomes, and that receiving practical support helps buffer the detrimental effects of stress as it reduces the situational demands of the individual.^{3,36} In addition, receiving tangible aid conveys to the individual that they are valued by their significant others which then bolsters their self-esteem and their sense of worth.

Guidance (advice and information) came last in the informal sources possibly because the bereaved prefer to obtain this from health professionals, who are equipped with the skills and knowledge though the advice and information were not necessarily about bereavement.³⁷ They also might like to receive this guidance from others who have experienced a similar loss rather than the general friends and family.³⁸ This was supported by guidance becoming more important in the community and professional sources: within the community sources of support, attachment and reliable alliance were still prominent followed by guidance, while in the professional sources, guidance became equal to attachment, but as expected there was no presence of reliable alliance.

In examining the unhelpful support, it seems that feeling unsupported was more than just the absence of support. While people did mention the absence of support, they also mentioned receiving insensitive comments and unwanted advice. However, fewer bereaved people had experienced unhelpful support (compared to helpful support) and that was mainly from government, financial and legal institutions.

Although the SPS framework is about social support rather than professional support, we took more of a social and public health approach to these care networks, both formal and informal, and considered all service providers as already existing members of communities, rather than placing them outside of these communities. This is supported by Horsfall et al. who explained that ‘In reality of course, especially in regional and rural areas, members of formal networks are also members of the community. They live, work, and play alongside people who are not necessarily providing a “service”’ (pp. 335–336).³⁹ It is worth noting that three out of four funeral service providers whose clients responded to this survey were in regional areas. This is the narrative that would operationalise the concepts of the public health approach and community capacity building through the compassionate communities approach. Horsfall et al.³⁹ called for better partnerships between such informal and formal caring networks, within the public health approach to end of life care. These partnership approaches

to community support strengthen the capacity of the whole community to respond to dying, death and loss providing synergies that enhance both formal and informal efforts for not only addressing the comorbidities of grief but also enhancing health and well-being.

In addition to improving emotional well-being and reducing isolation, research in the United Kingdom has shown that a compassionate communities approach can reduce health service usage and ultimately result in significant savings for health services.^{40,41} Therefore, communities need greater support to adopt a systematic approach to developing and mapping their own asset networks. There are numerous fruitful ways they might achieve this: through implementing suggested strategies such as the Compassionate City Charter or using Healthy End of Life Program (HELP) resources; building public policies that support dying, death, loss and grief; creating supporting environments and in particular social supports from workplaces and schools; strengthening existing community action; and developing personal skills in these areas while reorienting the existing health system towards a stronger and more enabling community and health services partnership.^{42,43}

Implications for research and practice

The implications for further research suggested by these findings indicate that we might usefully trial which elements of community development and support are universally important and which other elements are conditional upon regional and other social factors. This is not merely a call for greater evaluation of these approaches – evaluation of community development models similar to compassionate communities are commonplace in health promotion literature in general.⁴⁴ Rather it will be crucial to identify the essential professional or service elements for a programme of support that are best able to build upon local helping networks. In other words, more research is needed in identifying a range of useful practice models that connect health and social care services with local neighbourhoods and cultural life such as workplaces or schools. Evaluation studies will be only one element in a range of other essential and broader social research tasks (ethnographic studies, observational work, community policy audits, etc.) required for this programme style of development.

Furthermore in practice, we expect that members of the networks that support a dying relative or friend will continue to support each other during bereavement, unlike formal services where there is often a disjunction between the palliative care and bereavement teams.³⁰ This continuity of relationship matters and it is important to talk about the person who has died with others that have known the deceased. Compassionate communities provide this opportunity in the face of the fragmentation that characterises late modern society.⁴⁵

Further implications for practice relate to operationalising how to offer professionalised services to those who would most benefit, while not providing these services

unnecessarily to those who are supported within their natural support networks. This is a key challenge of the assets-based approach and requires further research, policy and practice investment. Implicit in this is an important question about how this balance will be achieved. Should professional services be making decisions about the assets to be involved? Or is this more properly a decision to be negotiated by the bereaved person and all involved in the support network?

In terms of what levels of bereavement care should be provided and to whom, we have attempted to offer some directions in our previous article,⁸ but more research is warranted. Our survey was constructed to provide information on the population-based experiences of bereavement, including the extent of the alignment of bereavement risk and support needs. Using the validated PG-13 (Prolonged Grief Disorder Scale) meant that we were able to determine a population rate of prolonged grief disorder of 6.4% in this sample. In addition, the low- and moderate-risk groups could be distinguished on the basis of PG-13 scores as well as the sources of support (mental health professional sources for high risk, community support for moderate risk and family and friends for low risk). In addition, the profiles of the bereaved provided a context for the scores: low scores typically arose from losses that were expected and less disruptive to daily routines (e.g. the death of an aged parent), moderate scores typically arose from deaths that were both expected and everyday-disruptive (e.g. the death of an ageing and ill partner) and high grief scores typically arose from deaths that were both unexpected or resisted and disruptive (such as a child or a younger spouse).

Limitations

The limitations of drawing on data from this anonymous survey have been highlighted in previous published findings from the survey, namely, that the study sample is not a random sample of the general bereaved population, but it nevertheless compares well with the UK mortality follow-back survey in terms of its demographic composition.^{8,30,46} The low response rate in this study is in line with others who relied on postal surveys with no reminder follow-up, acknowledging that those who did not respond may have had different experiences to those reported in this study.⁴⁷ The selection of respondents from funeral providers' database may have influenced the significant number reporting support from these providers. Although we were able to use the SPS framework to code the functions of social support, there is no comprehensive model of types of unhelpful supports, which is why we used inductive content analysis.

Other methods of qualitative data collection (e.g. interviews, focus groups) usually allow the opportunity for clarification of meaning or for probing of answers to gain depth in the participant response; this was not possible here as participants provided comments on a self-completed

questionnaire. However, the use of open response questions in surveys is increasingly being advocated as a means of enabling respondents to voice their opinion on the topic and is particularly suitable for a large sample size as in our study ($n=678$).⁴⁸

Conclusion

This is the first study to quantify the amount of support received by the bereaved from their own social networks, how this support works and why it may work. We argue, on the basis of our findings, for the importance of adopting a primary public health approach to support the majority of bereaved people, as this care is already provided in community settings.

Although this conclusion may seem rather obvious in most health policy and practice fields, it is important to note that, for the field of bereavement policy and practice, a public health approach is in its infancy. Furthermore, most research and practice development have to date been focussed on acute care models of therapeutic intervention and not, we emphasise, with a population approach to grief. Our study has revealed that most care comes from people already involved in the everyday lives of those recently bereaved. These people are assets already in place, contributing to each other's resilience. Some of them are healthcare practitioners contributing through their everyday activities, not bereavement programmes per se. In providing bereavement care, it is therefore prudent to support these 'everyday assets', ensuring that their care is recognised, appreciated and not disrupted by overreach from professional services.¹⁰ These assets will complement quality bereavement care that identifies and offers counselling and therapy to those who need it.

Much of the compassionate communities approach has been focussed on end of life care for ageing and dying, and it needs to be extended to bereavement. This study has provided further empirical evidence of the need for more community development approaches such as those exemplified by the compassionate communities movement in palliative care – developing these not solely for the direct end of life care context but also for the journey of grief and bereavement during and after those circumstances.

Acknowledgements

The authors acknowledge the financial support of the WA Cancer and Palliative Care Network, Health Department of Western Australia; the cooperation of funeral providers who assisted with the recruitment of participants, and the bereaved families who agreed to complete the survey and contributed to our understanding in this field. Special thanks to Denise Howting who assisted with the quantitative analysis.


Declaration of conflicting interests

The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

Funding

The author(s) disclosed receipt of the following financial support for the research, authorship, and/or publication of this article: This study was supported by the Health Department of Western Australia.

ORCID iD

Samar M Aoun  <https://orcid.org/0000-0002-4073-4805>

References

- Burke LA and Neimeyer RA. Complicated spiritual grief I: relation to complicated grief symptomatology following violent death bereavement. *Death Stud* 2014; 38: 259–267.
- Lobb EA, Kristjanson LJ, Aoun SM, et al. Predictors of complicated grief: a systematic review of empirical studies. *Death Stud* 2010; 34: 673–698.
- Bottomley JS, Burke LA and Neimeyer RA. Domains of social support that predict bereavement distress following homicide loss. *Omega* 2017; 75: 3–25.
- Logan EL, Thornton JA and Breen LJ. What determines supportive behaviors following bereavement? A systematic review and call to action. *Death Stud* 2017; 42: 104–114.
- Albrecht T and Adelman M. *Communicating social support*. Newbury Park, CA: SAGE, 1987.
- Stylianou SK and Vachon MLS. The role of social support in bereavement. In: Stroebe M, Stroebe W and Hansson R (eds) *Handbook of bereavement*. Cambridge: Cambridge University Press, 1993, pp. 397–410.
- Murray JA. Communicating with the community about grieving: a description and review of the foundations of a broken leg analogy of grieving. *J Loss Trauma* 2002; 7: 47–69.
- Aoun SM, Breen LJ, Howting DA, et al. Who needs bereavement support? A population based survey of bereavement risk and support need. *PLoS ONE* 2015; 10: e0121101.
- Kersting A, Braehler E, Glaesmer H, et al. Prevalence of complicated grief in a representative population-based sample. *J Affect Disord* 2011; 131: 339–343.
- Rumbold B and Aoun S. An assets-based approach to bereavement care. *Bereave Care* 2016; 34: 99–102.
- Sallnow L and Paul S. Understanding community engagement in end-of-life care: developing conceptual clarity. *Crit Public Health* 2015; 25: 231–238.
- Kellehear A. *Compassionate cities: public health and end-of-life care*. London: Routledge, 2005.
- Sallnow L, Kumar S and Kellehear A. *International perspectives on public health and palliative care*. Abingdon: Routledge, 2013.
- Wegleitner K, Heimerl K and Kellehear A. *Compassionate communities: case studies from Britain and Europe*. Abingdon: Routledge, 2016.
- Kellehear A. Compassionate communities: end-of-life care as everyone's responsibility. *QJM* 2013; 106: 1071–1075.
- National Palliative and End of Life Care Partnership. *Ambitions for palliative and end of life care: a national framework for local action 2015–2020*. London: NHS England, 2015.
- Dyregrov K. Micro-sociological analysis of social support following traumatic bereavement: unhelpful and avoidant responses from the community. *Omega* 2004; 48: 23–44.

18. Lehman DR, Ellard JH and Wortman CB. Social support for the bereaved: recipients' and providers' perspectives on what is helpful. *J Consult Clin Psych* 1986; 54: 438–446.
19. Peters K, Cunningham C, Murphy G, et al. Helpful and unhelpful responses after suicide: experiences of bereaved family members. *Int J Ment Health Nurs* 2016; 25: 418–425.
20. Bonanno GA, Wortman CB, Lehman DR, et al. Resilience to loss and chronic grief: a prospective study from preloss to 18-months postloss. *J Pers Soc Psychol* 2002; 83: 1150–1164.
21. Lichtenthal WG, Nilsson M, Kissane DW, et al. Underutilization of mental health services among bereaved caregivers with prolonged grief disorder. *Psychiatr Serv* 2011; 62: 1225–1229.
22. Weis R. The provisions of social relationships. In: Rubin Z (ed.) *Doing unto others*. Englewood Cliffs, NJ: Prentice Hall, 1974, pp. 17–26.
23. Cutrona C and Russell D. The provisions of social relationships and adaptation to stress. In: Jones W and Perlman D (eds) *Advances in personal relationships: a research annual*. Greenwich, CT: JAI Press, 1987, pp. 37–67.
24. Breen LJ, Aoun SM, O'Connor M, et al. Bridging the gaps in palliative care bereavement support: an international perspective. *Death Stud* 2014; 38: 54–61.
25. Wittouck C, Van Autreve S, De Jaegere E, et al. The prevention and treatment of complicated grief: a meta-analysis. *Clin Psychol Rev* 2011; 31: 69–78.
26. Aoun SM, Breen LJ, Rumbold B, et al. Reported experiences of bereavement support in Western Australia: a pilot study. *Aust NZ J Publ Heal* 2014; 38: 473–479.
27. Gottlieb BH and Bergen AE. Social support concepts and measures. *J Psychosom Res* 2010; 69: 511–520.
28. Hsieh HF and Shannon SE. Three approaches to qualitative content analysis. *Qual Health Res* 2005; 15: 1277–1288.
29. Horsfall D, Noonan K and Leonard R. Bringing our dying home: how caring for someone at end of life builds social capital and develops compassionate communities. *Health Sociol Rev* 2012; 21: 373–382.
30. Aoun SM, Rumbold B, Howting D, et al. Bereavement support for family caregivers: the gap between guidelines and practice in palliative care. *PLoS ONE* 2017; 12: e0184750.
31. Breen LJ, Fernandez M, O'Connor M, et al. The preparation of graduate health professionals for working with bereaved clients: an Australian perspective. *Omega* 2012; 66: 313–332.
32. Dickinson GE, Sumner ED and Frederick LM. Death education in selected health professions. *Death Stud* 1992; 16: 281–289.
33. Breen LJ. Professionals' experiences of grief counseling: implications for bridging the gap between research and practice. *Omega* 2010; 62: 285–303.
34. O'Connor M and Breen LJ. General practitioners' experiences of bereavement care and their educational support needs: a qualitative study. *BMC Med Educ* 2014; 14: 59.
35. Waller A, Turon H, Mansfield E, et al. Assisting the bereaved: a systematic review of the evidence for grief counselling. *Palliat Med* 2016; 30: 132–148.
36. Thoits P. Mechanisms linking social ties and support to physical and mental health. *J Health Soc Behav* 2011; 52: 145–161.
37. Vachon MLS and Stylianos SK. The role of social support in bereavement. *J Soc Issues* 1988; 44: 175–190.
38. Breen LJ and O'Connor M. Acts of resistance: breaking the silence of grief following traffic crash fatalities. *Death Stud* 2010; 34: 30–53.
39. Horsfall D, Leonard R, Noonan K, et al. Working together—apart: exploring the relationships between formal and informal care networks for people dying at home. *Prog Palliat Care* 2013; 21: 331–336.
40. Brown M. Compassionate Community Project. *Resurgence and Ecologist Magazine*, April 2018. <https://www.resurgence.org/magazine/article5039-compassionate-community-project.html>
41. Cronin P. Compassionate communities in Shropshire, West Midlands, England. In: Wegleitner K, Heimerl K and Kellehear A (eds) *Compassionate communities: case studies from Britain and Europe*. Abingdon: Routledge, 2016, pp. 30–45.
42. Grindrod A and Rumbold B. *Healthy End of Life Program (HELP): offering, asking for and accepting help. Creating an end of life collaborative community culture*. Melbourne, VIC, Australia: La Trobe University Palliative Care Unit, 2016.
43. Kellehear A. The compassionate city charter: inviting the cultural and social sectors into end-of-life care. In: Wegleitner K, Heimerl K and Kellehear (eds) *Compassionate communities: case studies from Britain and Europe*. Abingdon: Routledge, 2016, pp. 76–87.
44. O'Mara-Eves A, Brunton G, McDaid D, et al. Community engagement to reduce inequalities in health: a systematic review, meta-analysis and economic analysis. Southampton (UK): NIHR Journals Library; 2013. (Public Health Research, No. 1.4.) Available from: <https://www.ncbi.nlm.nih.gov/books/NBK262817/> doi: 10.3310/phr01040
45. Walter T. A new model of grief: bereavement and biography. *Mortality* 1996; 1: 7–25.
46. Calanzani N, Higginson IJ, Koffman J, et al. Factors associated with participation, active refusals and reasons for not taking part in a mortality followback survey evaluating end-of-life care. *PLoS ONE* 2016; 11: e0146134.
47. Hunt K, Shlomo N and Addington-Hall J. Participant recruitment in sensitive surveys: a comparative trial of 'opt in' versus 'opt out' approaches. *BMC Med Res Methodol* 2013; 13: 3.
48. O'Cathain A and Thomas KJ. 'Any other comments?' Open questions on questionnaires – a bane or a bonus to research? *BMC Med Res Methodol* 2004; 4: 25.