

# Northumbria Research Link

Citation: Tahmosybayat, Robin, Baker, Katherine, Godfrey, Alan, Caplan, Nick and Barry, Gill (2017) A systematic review and meta-analysis of outcome measures to assess postural control in older adults who undertake exergaming. *Maturitas*, 98. pp. 35-45. ISSN 0378-5122

Published by: Elsevier

URL: <https://doi.org/10.1016/j.maturitas.2017.02.003>  
<<https://doi.org/10.1016/j.maturitas.2017.02.003>>

This version was downloaded from Northumbria Research Link:  
<http://nrl.northumbria.ac.uk/29870/>

Northumbria University has developed Northumbria Research Link (NRL) to enable users to access the University's research output. Copyright © and moral rights for items on NRL are retained by the individual author(s) and/or other copyright owners. Single copies of full items can be reproduced, displayed or performed, and given to third parties in any format or medium for personal research or study, educational, or not-for-profit purposes without prior permission or charge, provided the authors, title and full bibliographic details are given, as well as a hyperlink and/or URL to the original metadata page. The content must not be changed in any way. Full items must not be sold commercially in any format or medium without formal permission of the copyright holder. The full policy is available online: <http://nrl.northumbria.ac.uk/policies.html>

This document may differ from the final, published version of the research and has been made available online in accordance with publisher policies. To read and/or cite from the published version of the research, please visit the publisher's website (a subscription may be required.)

[www.northumbria.ac.uk/nrl](http://www.northumbria.ac.uk/nrl)



1 **A systematic review and meta-analysis of outcome measures to assess**  
2 **postural control in older adults who undertake exergaming**

3 Robin Tahmosybayat<sup>a</sup>, BSc, Katherine Baker<sup>a</sup>, PhD, Alan Godfrey<sup>b</sup>, PhD, Nick Caplan<sup>a</sup>, PhD, Gill Barry<sup>ab</sup>, PhD.

4 <sup>a</sup> Department of Sport, Exercise and Rehabilitation, Faculty of Health and Life Science, University of Northumbria, Newcastle Upon Tyne, UK.

5 <sup>b</sup> Institute of Neuroscience, Newcastle University Institute for Ageing, Clinical Ageing Research Unit, Newcastle University, Campus for Ageing &  
6 Vitality, Newcastle upon Tyne, UK.

7  
8 **Corresponding Author**

9 Robin Tahmosybayat BSc (Hons)  
10 PhD Researcher  
11 robin.a.tahmosybayat@northumbria.ac.uk

12  
13 Department of Sport, Exercise and Rehabilitation  
14 Faculty of Health and Life Sciences  
15 Northumberland Building, Room 431  
16 Northumbria University  
17 Newcastle upon Tyne  
18 NE1 8ST

19  
20 **Abstract**

22 Exergaming has shown to be an effective tool to improve postural control (PC) in older community dwelling  
23 individuals. Outcome measures (OMs) used are varied and individually could hold limitations to the effectiveness  
24 of the intervention. This systematic review and meta-analysis aims to explore the OMs currently used to assess  
25 PC in exergaming interventions, for healthy elderly individuals > 60 years. The literature search was conducted  
26 across five databases (CINAHL, EMBASE, PubMed, ISI, SPORTdiscus and Science Direct) using a range of  
27 search terms and combinations relating to exergaming, balance, exercise, falls and elderly. Quality assessment  
28 was conducted using the PEDro Scale and a custom-made quality assessment tool. Eleven trials were included  
29 in the meta-analysis with a mean (SD) PEDro score of 5.36 (1.57). Primary and secondary OMs showed small  
30 effects in favour of alternative training modes, though insignificant for all primary OMs. Tertiary OMs could not be  
31 included in the meta-analysis due to varying output parameters from different instrumentation. Heterogeneity  
32 remained high across trials and no studies performed long term follow up of exergaming on PC. Exergaming is a  
33 potential alternative for PC training, although still in its infancy. Strong and well-designed RCTs are needed  
34 targeting specific populations > 60 years. Variability in instrumented OMs prevent generalising aspects of  
35 quantified PC. Improvements in technologies may provide data not currently available from clinical and laboratory  
36 based methods with means to measure PC more realistically and specifically to a population's ADLs, though this  
37 remains a new area of research.

38 Key words: Exergaming; Postural Control; Elderly; Outcome Measures; Meta-analysis; Community-dwelling;  
39 Balance; Falls

40  
41  
42  
43  
44  
45

46

47

48

## 49 **1.0 Introduction**

### 50 *1.1 Background*

51 Falls are associated with ageing and disease, with one third of people aged 65 years and older falling  
52 at least once per year [1, 2]. In older individuals, a strong predictor of falls is impaired postural control  
53 (PC) among other factors [3, 4]. Postural control is the ability to maintain, achieve, or restore a state  
54 of balance during any posture or activity [5]. Correct PC requires accurately timed vestibular, visual,  
55 proprioceptive and somatosensory inputs for adaptive strategies for orientation and balance [6].  
56 Participation in balance-based training is low due to the tedious and monotonous nature of the  
57 training [7]. These therapies are repetitive which reduce attention span and impair the effectiveness of  
58 the exercises, particularly the large volume of practice associated with chronic neurological and  
59 musculoskeletal conditions [7].

60 A more recent method of PC training is exergaming [7, 8]. Exergames are computer games driven by  
61 the user's gross physical movements. Due to portability, they facilitate community deployment  
62 whereby older individuals have experienced exergaming as a form of PC training [9]. The Nintendo  
63 Wii Fit™ had been the most popular exergaming instrument and results have shown beneficial effects  
64 on PC [9]. Other exergaming models include X-Box Kinect™, PlayStation Eyetoy™ and Dance Dance  
65 Revolution™. The X-Box Kinect™ is revolutionary in its development due to being the first  
66 commercial gaming system that does not require a hand held controller or external device, more so it  
67 requires the use of infra-red technology to track an individual's movements.

68 Outcome measures (OMs) used in exergaming interventions, employed for balance evaluation, have  
69 been previously categorised as functional assessment (documents balance status and change after  
70 intervention), systems assessment (determines the underlying reason for impaired balance control),  
71 static posturography (quantify postural sway while a subject remains as still as possible) and dynamic  
72 posturography (use of external balance perturbations, changing surface and visual conditions) [10].  
73 The Berg Balance Scale (BBS) [11] and the Tinetti Performance Oriented Mobility Assessment  
74 (POMA) [12] quantify functional balance in an ordinal pattern as the participant performs balance and  
75 mobility tasks that represent activities of daily living (ADLs). The Functional Reach Test (FRT) [13]  
76 uses distance to quantify limits of stability of the centre of mass. The Single Leg Stance (SLS) [14] or  
77 the Timed Up and Go (TUG) [15] use the time domain to measure the task being performed via a stop

78 watch. These measures provide information about postural control, likelihood of falling and functional  
79 capabilities. Inter-rater reliability has been previously reported excellent for BBS, TUG and FRT as  
80 has good intra-rater reliability [16]. Unobtrusive self-report questionnaires such as the Tinetti Falls  
81 Efficacy Scale (FES) [17] and the Activities-specific Balance Confidence Scale (ABC) [18] measure  
82 perception of balance confidence and fear of falling of an individual in performing ADLs.

83 Force platforms quantify the centre of pressure (COP) excursion in mediolateral (ML) and  
84 anteroposterior (AP) direction during quiet stance in varying conditions [7]. The COP has previously  
85 characterised postural control by evaluating the relative sensitivity of COP based measures to  
86 changes in postural steadiness [19] and has been correlated with poor balance and risk of falls [20].  
87 Older adults have previously demonstrated larger areas of COP excursion on a force platform with  
88 eyes open, eyes closed or with visual feedback. They displayed longer movement times, longer path  
89 lengths of the participant's centre-of-gravity (COG) to different points within their limits-of-stability, and  
90 shorter distances of functional reach when compared with younger adults [21]. Miniaturised  
91 electronic-based wearables with inertial sensors (e.g. accelerometers and gyroscopes) have  
92 objectively and reliably measured postural sway during quiet stance [22-24]. Wearables have been  
93 introduced in clinics as an alternative to evaluating PC in the hope to eliminate clinician bias, increase  
94 sensitivity to mild impairments (ceiling effects) and improve reliability of measures [25, 26]. They have  
95 been tested in clinical populations whereby a subset of sensitive, reliable and valid instrumented  
96 postural sway characteristics had been formed [27].

97 It appears necessary to systematically explore OMs used in exergaming interventions in the hope to  
98 establish if an influence on intervention effect exists and any individual limitations that OMs may hold.

## 99 1.2 Objective

100 The aim of this systematic review and meta-analysis is to explore the outcome measures currently  
101 used to assess PC in exergaming interventions for healthy elderly individuals > 60 years.

102

103

104

105

106 **2.0 Methods**

107 **2.1 Search strategy**

108 This systematic review was reported according to the PRISMA guidelines [28]. The systematic review  
 109 was beyond the stage of data collection and therefore could not be registered with PROSPERO,  
 110 however, it did receive an official statement pertaining to its satisfaction of the inclusion criteria. This  
 111 is available upon request. Electronic databases (CINAHL, EMBASE, PubMed, Web of Science,  
 112 SPORTdiscus and Science Direct) were searched for publications from January 2000 to April 2016 for  
 113 interventions performed in clinical and community based settings. The key search terms were merged  
 114 with Boolean conjunction (OR/AND) and applied on three search levels. Key Search terms used were:  
 115 (exergam\* OR exer-gam\* OR videogam\* OR video-gam\* OR video-based OR Wii OR Nintendo OR  
 116 X-box OR Kinect OR play-station OR playstation OR virtua\* realit\* OR dance dance revolution) AND  
 117 (sport\* OR train\* OR exercis\* OR intervent\* OR balanc\* OR strength OR coordina\* OR motor control  
 118 OR postur\* OR power OR physical\* OR activit\* OR health\* OR fall\* risk OR prevent\*) AND (old\* OR  
 119 elder\* OR senior\*). Three levels of screening were carried out: (1) title, (2) abstract, and (3) full-text.  
 120 The reference lists of the included articles were also searched. Inclusion/exclusion criteria were  
 121 agreed upon by the two reviewers (RT & GB).

122

123 **2.2 Selection Criteria (PICOS)**

**Table 1: Inclusion and exclusion criteria**

	Inclusion	Exclusion
Population	Older Individuals between the age of 60 and 85 years old, no neurologic or orthopaedic condition, community dwelling or independently in retirement centres, without cognitive impairment, able to ambulate independently without assistive devices were included.	Individuals who were outside the age range of 60 - 85 years old. Populations with specific neurological (i.e. stroke, Parkinson's disease, and multiple sclerosis), metabolic (i.e. diabetes), or musculoskeletal (i.e. rheumatoid arthritis) deficits that might impair PC were excluded.
Intervention	Intervention group treated with exergaming as balance training only or combined with other forms of training such as strength training were included.	Studies where the intervention group was not treated with exergaming as balance training (i.e. virtual reality treadmill training, biofeedback) was excluded.
Comparison	A comparison group treated with traditional balance training or with no intervention or both were included.	Studies not utilising any comparison groups were excluded.
Outcomes	Outcome measures designed to objectively and subjectively assess PC (functional assessment, laboratory based assessment, self-report assessment).	Balance as a tertiary measure was excluded.

Studies	Randomised controlled trials (RCT), controlled trials (CT), two group pre and post comparison studies, whereby primary outcome measures were used to assess balance or PC either/or before, during and after a bout of exergaming were included.	Studies with fewer than six participants in each intervention group were excluded. Studies in which no inferential statistics were reported were excluded. Studies that did not meet the inclusion criteria (e.g. all (non-human) animal research)
---------	--	--

124

125           2.3     *Data Extraction*

126 Quantitative data were extracted by one reviewer (RT) and checked by another (GB). Specific details  
 127 about the interventions, populations and study methods were extracted. Primary methods to assess  
 128 PC were categorised based on traditional standing and functional mobility tests categorised into rating  
 129 scales, distance based measures and timed tasks. Secondary methods were based on self-report  
 130 measures of balance and fear of falling (self-report questionnaires). Tertiary methods were  
 131 categorised as any instrumentation that quantified PC (force platforms, perturbation platforms and  
 132 accelerometers).

133           2.4     *Quality Assessment*

134 Evidence level of included studies were assessed using the Oxford 2011 Centre for Evidence-Based  
 135 Medicine Levels of Evidence [29]. Of the five levels of evidence, level 1 is deemed to be the highest  
 136 quality of evidence (supplementary file 1, A). To eliminate unintended bias while assessing the  
 137 studies, both reviewers collaborated and eliminated any conflicting opinions. Eligibility and quality of  
 138 studies was assessed using the Physiotherapy Evidence Database Scale (PEDro) and were  
 139 independently assessed by both reviewers (supplementary file 1, B). Methodological quality was also  
 140 assessed using a custom-made tool derived from a previous systematic review (supplementary file 1,  
 141 C) [30].

142

143           2.5     *Data analysis*

144 Intervention effects were assessed by grouping studies for meta-analysis by the method of assessing  
 145 PC (Primary, secondary and tertiary). The difference of the target outcome between the intervention  
 146 and the control group including the pooled standard deviations, were calculated for different  
 147 categories of outcome measure. Random effects models (Review Manager (Revman), version 5.3,  
 148 Copenhagen, Denmark) were used and between-group standardized mean differences (SMD) were  
 149 calculated based on continuous measurement scale (mean  $\pm$ SD). Hedge's g was used to quantify

150 effect sizes for SMD to account for small sample sizes ( $n < 20$ ). For trials utilising multiple intervention  
151 arms and compared an exergaming group with an alternative balance training group (group fitness,  
152 standardised balance training program, Tai Chi etc.) and a control group (no exercise), the alternative  
153 balance training control group were compared to the exergaming group. Where a secondary active  
154 control group was included in the study, the control group most representative of traditional balance  
155 training was compared to the exergaming group. If the heterogeneity test revealed a value of  $p < 0.1$   
156 or  $I^2 > 25\%$ , then heterogeneity was considered likely. Heterogeneity was deemed moderate at  $< 50\%$   
157 and considerable at  $> 50\%$  [31, 32].

### 158 **3.0 Results**

#### 159 *3.1 Search Strategy*

160 The database search yielded 809 publications (Figure 1). After removing all duplicates (346), 463  
161 publications were abstract screened whereby 435 were excluded leaving 28 publications. After  
162 searching reference lists of the 28 included publications, an additional 26 were obtained leaving 54.  
163 Of the 54 publications, 42 were excluded with reasons to give the final number of included  
164 publications for qualitative synthesis in the review ( $n=12$ ). The publications remaining for qualitative  
165 review can be found here (supplementary file 1, D). Of the 12 publications, one was excluded from  
166 the meta-analysis where insufficient data were reported. Data was acquired from one author [33] and  
167 another failed to respond [34]. Additionally, the Cochrane Central Register of Controlled Trials  
168 revealed no further publications for inclusion in this review.




Figure 1. Flow of study screening and selection

169

170

171

172

173

174

175



176           3.2     *Quality Assessment*

177     Nine of the included publications were considered level 2 evidence (RCT's) and 3 non-RCT's were  
178     considered level 3 evidence base (supplementary file 1, E). The mean (SD) methodological quality  
179     score of the 12 trials included in the review was 5.17 (1.64). This increased to 5.36 (1.57) for the  
180     eleven trials included in the meta-analysis. When excluding the level 3 evidence trials (non-RCT) from  
181     the quality assessment the score increased to 5.44 (1.74). A third of the trials reviewed were rated  
182     below the mean score which can be attributed to a lack of blinding of the participants, therapists and  
183     assessors and a lack of allocation concealment (Table 3). There was a seeming lack of explanation  
184     for randomisation across trials with only two studies adequately explaining the method for  
185     randomising participants. Six trials failed to describe location and no intervention follow up was  
186     conducted for any of the trials (supplementary file 1, F).

**Table 2.** Overview of the study design, sample characteristics, groups, intervention type and location for included studies

Author and Date	Study Design	Sample: Population; Sample Size (n); age, years (mean $\pm$ SD), M/F	Groups	Intervention & Follow up (Y/N)	Location/ Settings
Pluchino et al., 2012	RCT 3 arms (PS)	Community-dwelling older adults, n=40; 72.5 $\pm$ 8.4 years, 15/25	IG1 : Standard Balance Exercise ; (n=14), IG2 :Tai Chi (n=14), IG3 : WF (n=12)	60 minutes, 2 x per week, 8 weeks. (N)	Research laboratory/training facility, Wii group unsupervised.
Ray et al., 2012	RCT 3 arms	Community-dwelling older adults, n=87, 75 years (no SD given), 29/58	GF: (n=40), WF + weighted vest: (n=29), CG: (n=18)	GF & WF: 3 x week 45 mins duration, 15 weeks. (N)	Laboratory
Toulotte et al., 2012	RCT 4 arms	Community-dwelling older adults., n=36, 14/22. See adjacent column for mean age (SD) per group	G1: APA, (n=9, 84.2 $\pm$ 8.1 years, 3/6). G2: Wii Fit, (n=9, 72.2 $\pm$ 8.6 years 4/5). G3: APA + WF, (n=9, 76.4 $\pm$ 4.7 years, 3/6). G4: CG (n=9, 71.8 $\pm$ 8.0, 4/5).	60 minutes per week x 20 weeks. (N)	Gymnasium at retirement centre
Merriman et al., 2015	RCT 2 arms	Community-dwelling n=59 & Retired Persons n=17, subgroups: healthy n=42, fall prone n=34, 16/60. See adjacent column for mean age (SD) per group	IG: Balance Training (n=38, 17 his of falls, 74.06 (6.66) years, 21 healthy, 74.90 (8.97) years, 1/37). CG: (n=38, 17 his of falls 73.41 (7.00) years, 21 healthy 74.33 (11.09) years, 15/23)	IG: 5 weeks, 2 x 30 min BT/week CG: diary of light, med, heavy Physical Activity. (N)	Sheltered accommodation / community centre / testing laboratory
Sato et al., 2015	RCT 2 arms	Community-dwelling older adults, n=54, 69.25 $\pm$ 5.4 years, 11/43	IG: (n=29) CG: (n=28)	65.34 (9.63) days, 40 mins - 1 hour per session, 2-3 times per week, total 24 times. (N)	N/A
Whyatt et al., 2015	RCT 2 arms	Sheltered accommodation and local activity groups, n=84, 25/57. See adjacent column for mean age (SD) per group	IG: Balance Game Training, n=40, 77.18– 6.59 years, 5/35. CG: n=42, 76.62– 7.28 years 20/22. Subgroups. High Risk Falls: IG: (n=15, 77.73 – 8.01 years, 2/13). CG: (n=12, 79.00 – 7.03 years, 6/6). Low Risk Falls: IG: (n=25, 76.83– 5.64 years 3/22). CG: (n=30, 75.67 – 7.28 years, 14/16).	IG: 30 minutes per session, 10 x sessions; over 5 weeks. CG: 5 weeks of recording levels of physical activity. (N)	N/A
Lai et al., 2013	RCT 2 arms	Community-living persons n=30, 72.1 [4.8] years, 13/17	Group A: (n=15, 70.6 (3.5) years 7/8). Group B:(n=15, 74.8 (4.7) years, 6/9). Both Groups performed an intervention phase and a control phase.	12 weeks' trial. IG: 30 min, 3 times/ week x 6 weeks then 6 weeks no exercise. CG: no exercise x 6 weeks then IG 6weeks. (N)	N/A
Singh et al., 2013	RCT 2 arms	Community-dwelling older women, n=38, 36 completed intervention.	IG: balance-focused virtual-reality games 61.12 ( 3.72) years, CG: therapeutic balance exercises: 64.00 ( 5.88) years,	30 minutes, 2 x / week for 6 weeks. (N)	N/A
Chow and Mann, 2015	RCPS 2 arms	Community-dwelling, n=20, 69 (range 65 - 78), 7/13	IG: Daily Cyber Golfing n=10, 70.4 (5.4) years, 3/7 CG: regular table games n=10, 68.0 (3.0) years, 4/6.	Daily, 30-45 minutes for 2 weeks. (N)	N/A
Nicholson et al., 2015	Non-RCT	Local retirement villages and educational settings, n=41, 74.5 (5.4) years, 14/27	IG: Wii group (n = 19, 75.11 (5.85) years, 7/12, 2 fallers). CG:(n = 22, 73.91 (5.12) years, 7/15, 3 fallers)	IG: 3 x 30 min Wii Fit sessions per week for six weeks. CG: usual everyday activities and exercise routines. (N)	Unsupervised, in pairs in community hall of a retirement village
Park et al., 2015	Non-RCT	Community Dwelling Individuals, n=30	VRG: (n=15, 66.5 $\pm$ 8.1 years, 9/3) and a BEG: (n=15, 65.2 $\pm$ 7.9 years, 10/2)	30 min 3 times a week for 8 weeks. (N)	N/A
Tange et al., 2012	Non-RCT (PS)	Elderly individuals, n=39,	WSG: n=20 77 (68-82) years, WF: n=19, 84 (80-89) years	2 x / week during 6 weeks in one-hour sessions. (N)	N/A

RCT = randomised control trial; (PS) = Pilot Study; SD = Standard Deviation; M/F = Male/ Female; (n) = number; (Y/N) = Yes/No; G1 = Group 1; G2 = Group 2; G3 = Group 3; G4 = Group 4; APA = Adapted Physical Activities; WF = Wii Fit; CG = Control Group; IG = Intervention Group; GF = Group Fitness; VRG = virtual reality group; BEG = Ball Exercise Group; WSG = Wii Sports Group; N/A = Not Applicable; mins = minutes.

**Table 3.** Outcomes from PEDro scale quality assessment

Author and Date	Eligibility Criteria	Random Allocation	Concealed allocation	Baseline Comparable	Blind Subject	Blind Therapist	Blind Assessor	Adequate Follow up	Intention to treat	Between group comparison	Point Estimates and Variability	Total
<b>RCT</b>												
Pluchino et al., 2012 *	Y	Y	Y	Y	N	N	N	N	N	Y	Y	5
Ray et al., 2012 *	Y	Y	N	N	N	N	N	N	N	Y	Y	3
Toulotte et al., 2012 *	Y	Y	Y	Y	N	N	N	Y	N	N	Y	5
Merriman et al., 2015 *	N	N	N	N	N	N	N	N	Y	Y	Y	3
Sato et al., 2015 *	Y	Y	Y	Y	Y	Y	Y	Y	N	N	Y	8
Whyatt et al., 2015 *	Y	Y	N	Y	N	N	N	Y	N	Y	Y	5
Lai et al., 2013 *	Y	Y	N	Y	N	N	Y	Y	Y	Y	Y	7
Singh et al., 2013 *	Y	Y	Y	Y	N	N	Y	Y	N	Y	Y	7
Chow and Mann., 2015 *	N	Y	N	Y	N	N	N	Y	Y	Y	Y	6
<b>Non-RCT</b>												
Nicholson et al., 2015 *	N	N	N	Y	N	N	N	Y	Y	Y	Y	5
Park et al., 2015 *	Y	N	N	Y	N	N	N	N	Y	Y	Y	4
Tange et al., 2012	Y	N	N	N	N	N	N	Y	N	Y	Y	3
<b>Total</b>	<b>9</b>	<b>8</b>	<b>4</b>	<b>9</b>	<b>1</b>	<b>1</b>	<b>3</b>	<b>8</b>	<b>5</b>	<b>10</b>	<b>12</b>	

RCT = Randomised Control Trial; Non-RCT = Non Randomised Control Trial; Y = Yes; N = No; \* = Included in Meta-Analysis.

187           3.3     *Data Extraction*


188     Intervention characteristics are available in Table 2. Intervention duration ranged from 5 to 20 weeks,  
189     individual sessions ranged from 30 to 60 minutes and session frequency ranged from 1 to 3 times per  
190     week. The majority of interventions were conducted in a research facility or a dedicated testing room  
191     in a community centre. None of the interventions took place in the home environment and two trials  
192     performed exergaming unsupervised [35, 36] (Table 3). Trials were conducted in the USA [35-37], the  
193     UK [38, 39], The Netherlands [34], France [40], Malaysia [41], Hong Kong [33], Japan [42], Taiwan  
194     [43] and South Korea [44].

195           3.4     *Intervention Effect*

196                   3.4.1    *Primary and Secondary OMs*

197     Of the 11 trials included in the meta-analysis, six reported PC outcomes from rating scales [35, 38-40,  
198     42, 43], three reported stand and reach tasks, one reported a sit and reach task [33, 35-37] and seven  
199     trials included timed tasks consisting of standing balance and mobility assessment [33, 35-37, 41, 43,  
200     44]. Data for included studies can be viewed in supplementary file 1, G. Five trials used self-report  
201     methods to quantify balance confidence and fear of falling [35, 36, 38, 39, 43]. Four trials used  
202     various versions of the falls efficacy scale [35, 36, 38, 43]. Two trials administered the ABC scale [38,  
203     39], one trial administered fall risk for older individuals living in the community [35] and one trial  
204     administered a questionnaire to measure fear of falling [38].


205     Exergaming had less of an effect on PC than alternative balance training modes when measured  
206     using rating scales (SMD: -0.27, 95% CI = -0.23 to 0.78;  $I^2 = 80\%$ ) (Figure 2) and distance-based  
207     reaching tasks (SMD: -0.28, 95% CI -0.70 to 0.15,  $I^2 = 57\%$ ) (Figure 3) but no effect was seen in  
208     favour of either intervention method through timed tasks (SMD: -0.03, 95% CI -0.30 to 0.24;  $I^2 = 50\%$ )  
209     (Figure 4). Exergaming had less of an effect on balance confidence and fear of falling than active  
210     controls when measured using questionnaires (SMD: -0.23, 95% CI 0.03 to 0.44;  $I^2 = 0\%$ ).



211

212 **Figure 2.** Outcome measures using rating scales for PC assessment in Exergaming vs. active controls. BBS = Berg balance  
 213 scale; POMA = Performance Oriented Mobility Assessment; Std. = standardised; IV = inverse variance; CI = confidence interval.


214



215

216 **Figure 3.** Outcome measures using reaching tasks for Exergaming vs. active controls. FRT = Functional Reach Test; LRT – L  
 217 = Lateral Reach Test Left; LRT-R = Lateral Reach Test Right; Std. = standardised; IV = inverse variance; CI = confidence  
 218 interval.


219



220

221 **Figure 4.** Outcome measures using timed tasks for exergaming vs. active controls. TUG = Timed Up and Go; OLSEO = One  
 222 Leg Stance Eyes Open; Std. = standardised; IV = inverse variance; CI = confidence interval.

223



224

225 **Figure 5.** Self-Report Measures of balance confidence and fear of falling for exergaming vs. active controls. FES = Falls  
 226 Efficacy Scale; ABC = Activities-specific Balance Confidence Scale; FROP-COM = Falls Risk for Older People living in the  
 227 Community; I = Iconographical and M = Modified; Std. = standardised; IV = inverse variance; CI = confidence interval.

228 After excluding non-RCT's to observe for any differences in the direction of the effect, the effect made  
229 a positive transition towards exergaming for distance-based reaching tasks (SMD: 0.10, 95% CI -0.39  
230 to 0.59,  $I^2 = 26%$ ) and marginally for timed tasks (SMD: 0.01, 95% CI -0.28 to 0.30,  $I^2 = 34%$ ), though  
231 remained statistically insignificant. A noticeable reduction in heterogeneity across studies was  
232 observed for sub-categories of primary OM (supplementary file 1, H). Findings from primary and  
233 secondary OMs with insufficient data to pool into meta-analysis can be viewed in supplementary file 1,  
234 I.

### 235 *3.4.2 Tertiary OMs*

236 The instrumentation used to quantify PC had many variations of measurement output which meant  
237 inclusion in the meta-analysis was not feasible. Individual results pertaining to intervention effect can  
238 be found in supplementary file 1, J.

239

240

**Table 4. Overview of primary, secondary and tertiary outcome measures used to assess balance**

Author and Date	Systems and apparatus	Primary OMs	Secondary OMs	Tertiary OMs	Details
<b>Pluchino et al., 2012</b>	AccuSway Force Platform, Proprio 5000 Dynamic Posturography platform	One-Leg Stance (s), Functional Reach Test (cm), Timed Up & Go Test (s), Tinetti Performance Oriented Mobility Assessment	Falls Efficacy Scale (FES), Falls Risk for Older People–Community Setting (FROP-COM).	The Postural Sway Test (COP + Time to boundary), Dynamic Posturography Test (perturbation platform)	Postural Sway Test Parameters: COP characteristics in AP and ML direction
<b>Ray et al., 2012</b>	NeuroCom SOT	8ft Timed Up and Go Test (s), Chair stand x 15-25 reps weighted, 6-minute walk test, Sit and Reach Test.	N/A	Sensory Organisation Test: 6 conditions, 3 trials/ condition. 18 trials total. 20 s/ trial.	Composite Equilibrium Score of weighted value of 6 conditions: Strategy Analysis score: Scores between 0 and 100 represent a combination of the two strategies; ankle and hip.
<b>Toulotte et al., 2012</b>	Nintendo Wii Fit + WBB	Unipedal Test Eyes Open, Eyes Closed, Tinetti Balance Assessment tool.	N/A	Wii Fit Test - Position of Centre Of Gravity (COG)	The videogame console gives two percentages (right and left) for the position of the centre of gravity. We calculated the percentage difference between right and left and concluded as to the overall position of the centre of gravity.
<b>Merriman et al., 2015</b>	Wii Balance Board (embedded with safety frame surrounding)+ Custom Designed Game	Berg Balance Scale	Balance Confidence (ABC) Scale, Fear of Falling (FOF) Falls Efficacy Scale (FES)	Static and Dynamic Balance Test.	Static: No. of secs within target area (max 10) converted to a percentage. 3 trials per target zone and average score across trials was collected. Dynamic: No of time to reach targets at fixed locations in 60s.
<b>Sato et al., 2015</b>	N/A	Berg Balance Scale, Functional Reach Test (cm), Chair Stand-30s	N/A	N/A	N/A
<b>Whyatt et al., 2015</b>	Nintendo Wii Fit, Wii Balance Board, Zimmer frame for safety, The NeuroCom Balance Master	Berg Balance Scale	ABC Scale	Custom made Static Balance Test (COP Displacement), Dynamic Balance Test - Limits of stability (COP)	Static: percentage of time spent in the target area. Dynamic: No. of targets hit COP displacement. Scores represent levels of COP spatial accuracy and data for all balance tests were converted to percentage change between Session 1 and Session 2.
<b>Lai et al., 2013</b>	The Catsys 2000 system measures postural sway, Xavix Measured Step System (XMSS)	Berg Balance Scale, Timed Up and Go Test (s), Unipedal Stance Test, XMSS stepping test	Modified Falls Efficacy Scale (MFES)	Stepping Test, Sway Area (SA), postural sway (Sway Velocity (SV) of COP in bipedal stance with eyes open and closed)	Sway Area (SA) and Sway Velocity (SV) COP in a bipedal stance with eyes open and closed. Postural sway was measured for 75 s (standard test procedure: 10 s start-up period, 60 s recording period, and 5 s run-out period), while standing directly on the platform
<b>Singh et al., 2013</b>	Probalance System	Timed up and Go Test (s), Ten Step Test		Postural Sway	Anterior –posterior and medial – lateral sway scores were converted to an overall performance index (OPI) by the Probalance software program. Lower OPI scores reflect better ability to regulate postural sway.



<b>Chow and Mann, 2015</b>	N/A	Timed up and go test (s), Single leg stance test, Functional Reach test (cm).	N/A	N/A
<b>Nicholson et al., 2015</b>	N/A	Timed Up and Go Test (s) Functional reach (cm) Lateral reach left (cm) Lateral reach right (cm) Single Leg Stance left (s) Single Leg Stance right (s) 30-s chair stand, Gait speed (m/s)	N/A	N/A
<b>Park et al., 2015</b>	BioRescue	Timed Up and Go Test (s)	Static Balance	30 sec sway length (mm) & average sway speed (mm <sup>2</sup> ) EO (COP) + biofeedback
<b>Tange et al., 2012</b>	N/A	Berg Balance Scale at 0, 3, and 6 weeks	N/A	N/A

OMs = Outcome measures; N/A = Not Applicable; COP = Centre of Pressure; SOT = Sensory Organisation Test; (s) = seconds; (cm) = centimetres; (m/s) = metres per second; mm<sup>2</sup> = millimetres squared; EO = Eyes Open

## 241 **4.0 Discussion**

242 This systematic review and meta-analysis aimed to explore OMs used to assess PC in exergaming  
243 interventions in individuals aged 60 years or more. The evidence from the meta-analyses suggest  
244 that, overall, the use of primary and secondary OMs do not impact the outcome of the intervention  
245 although after dividing the meta-analyses by individual measure type, some measures favoured  
246 exergaming more so than others and heterogeneity was moderate to high for primary OMs. After  
247 removing the non-randomised studies from the meta-analyses, the overall effect swayed toward  
248 exergaming.

### 249 *4.1 Limitations with the measures*

250 The primary measures used in this systematic review consist of clinical balance assessments which  
251 were originally created to identify balance problems or the underlying cause of a problem to predict  
252 risk of falls and determine effectiveness of intervention [10]. Healthy community dwelling older adults  
253 tend to have higher functioning capabilities and the 8 points of clinically significant change [45]  
254 required in the BBS questions the validity of this assessment for already high functioning individuals  
255 and has shown ceiling effects in this regard [46]. The gait section of the Tinetti POMA is seldom used  
256 and has also shown ceiling effects [45]. The FRT, despite its purpose, has not been well correlated  
257 with centre of mass displacement due to availability of compensatory strategies to reach not  
258 accounted for in the test [47]. The TUG also suffers the inability to detect early onset of impairment  
259 and the inability to understand if it is the gait or balance component of the scale that is affected may  
260 limit this form of measure. The use of rating scales, distance-based measures and timed tasks is  
261 practical and inexpensive for PC assessment however, the ceiling effects observed in this population  
262 hinder the ability to predict any future concerns of healthy individuals, which is valuable information in  
263 order to understand changes in PC. The use of questionnaires to evaluate self-perceived balance  
264 confidence and fear of falling are useful as they are nonintrusive and support the targeted direction of  
265 an intervention [10]. The ABC scale was developed on elderly outpatients and the confidence they  
266 perceived was based on a perceived need for a walking aid and personal assistance to ambulate  
267 outdoors [18]. Balance evaluation measures have been previously rated in terms of the ability to  
268 measure different aspects of PC and only one measure assessed all 6 aspects of postural control [48].  
269 Adapted measures could discriminate higher functional balance ability in this specific population,

270 which could result in a greater understanding of the effect of the intervention on PC. The needs of  
271 higher functioning older adults are less dependent and more focused on higher levels of activities of  
272 daily living [18].

273 The range of equipment and output parameters relating to the COP characteristics of PC requires  
274 consistency in order for instrumented outcome measures to be generalizable in the future. For  
275 example, comparing COP parameters using a force platform in Pluchino et al.'s [35] trial with the  
276 percentage change of the COG measured on a Wii Balance Board in a trial by Toulotte et al. [40].  
277 Several studies did report that participants tended to enjoy exergaming and increased motivation was  
278 observed but not measured in several trials. This concurs with several previous systematic reviews  
279 [49-51]. A limitation to force plate PC assessment is the inability to measure stepping action of  
280 dynamic balance, or indeed the dynamic balance accounted for during gait [52]. Individuals perform  
281 reactive and proactive PC adjustments on a force platform [20], but with the individual rooted to the  
282 platform, whether it is embedded or raised, not all components of the PC system are challenged as  
283 the base of support remains in a static state. Recent research has shown the importance of stepping  
284 action for prevention of falls and improving PC [53]. Postural control demands may be influenced by  
285 the complexity of the task and the environment in which the task is performed [54]. The use of a body  
286 worn accelerometer (BWA) to track PC and gait in any environment has previously been  
287 demonstrated as part of the development of an instrumented physical capability assessment (ICAP)  
288 [26], yet was not used to quantify PC in any of the trials in this review. The ability of BWA to track PC  
289 over a period of time with standardised protocols [25] could enable accurate assessment of PC in  
290 community environments for both healthy and fall prone individuals, with varying complexity of task  
291 and environmental demands. The potential for BWAs to be able to track higher functioning older  
292 individuals may eliminate the psychometric limitations seen in more traditional methods.

#### 293 4.2 Overall effect

294 The meta-analyses did show that exergaming interventions are less effective when compared to  
295 alternative balance training modes. After adjusting the meta-analyses to include only RCT's there  
296 was a shift in effect which could be attributed to the removal of non-RCTs. This is an assumption and  
297 must be considered lightly. None of the trials included in the current review performed follow-up  
298 measurements leaving a gap in the knowledge of long-term effects of exergaming on PC. Previous

299 systematic reviews have also reported similar findings [9, 55] although reported on  $p$  values alone.  
300 The use of meta-analyses to report effect sizes are arguably more appropriate for intervention  
301 evaluation [50].

#### 302 4.3 *Strengths and Limitations*

303 This systematic review was conducted in line with the PRISMA statement. The effects of the current  
304 meta-analysis must be taken with caution due to the small number of trials included in the review. The  
305 high heterogeneity and a lack of intention-to-treat analysis may not give a comprehensive picture of  
306 the effects of exergaming on PC. Furthermore, this review reported on healthy community dwelling  
307 individuals only and not those with pathological conditions and at higher risks of falls. The non-RCT's  
308 used in the meta-analyses sway potential biases and although we attempted to account for the  
309 differences, results should be interpreted carefully, particularly concerning selection bias and  
310 reporting bias.

311

#### 312 **5.0 Conclusion**

313 Exergaming is still in its infancy and heterogeneity in intervention design may affect the overall  
314 intervention effect. High quality RCTs with long periods of follow up are needed in order to inform  
315 recommendations for exergaming interventions focusing on improving PC. OMs used to assess PC in  
316 this population hold psychometric limitations and balance measures do not assess all aspect of PC.  
317 OMs that can differentiate balance problems within this population may help direct exergaming  
318 interventions. Improvements in technologies may provide further insight with means to measure PC  
319 more specifically to a population's ADLs.

320

#### 321 **Conflict of Interest statement**

322 The authors declare that there are no conflicts of interests.

323

#### 324 **Additional File 1. Supplementary data and Figures**

325 Supplementary data to this article can be found in additional file 1.

326

327 **Acknowledgements**

328 This research is funded as part of a PhD programme within the Faculty of Health and Life Sciences at  
329 Northumbria University, Newcastle Upon Tyne, UK.

330

331 **References**

- 332 1. Spaniolas, K., et al., *Ground level falls are associated with significant mortality in elderly*  
333 *patients*. Journal of Trauma and Acute Care Surgery, 2010. **69**(4): p. 821-825.
- 334 2. Gill, T.M., et al., *Association of injurious falls with disability outcomes and nursing home*  
335 *admissions in community-living older persons*. American Journal of Epidemiology, 2013.  
336 **3**(178): p. 418-425.
- 337 3. Lajoie, Y., *Effect of computerized feedback postural training on posture and attentional*  
338 *demands in older adults*. Aging Clinical and Experimental Research, 2004. **16**(5): p. 363-368.
- 339 4. Delbaere, K., et al., *A multifactorial approach to understanding fall risk in older people*.  
340 Journal of the American Geriatrics Society, 2010. **58**(9): p. 1679-1685.
- 341 5. Pollock, A.S., et al., *What is balance?* Clinical Rehabilitation, 2000. **14**(4): p. 402-406.
- 342 6. Laughton, C.A., et al., *Aging, muscle activity, and balance control: physiologic changes*  
343 *associated with balance impairment*. Gait & Posture, 2003. **18**(2): p. 101-108.
- 344 7. van Diest, M., et al., *Exergaming for balance training of elderly: state of the art and future*  
345 *developments*. Journal of Neuroengineering and Rehabilitation, 2013. **10**: p. 101.
- 346 8. Bateni, H., *Changes in balance in older adults based on use of physical therapy vs the Wii Fit*  
347 *gaming system: a preliminary study*. Physiotherapy, 2012. **98**(3): p. 211-216.
- 348 9. Laufer, Y., G. Dar, and E. Kodesh, *Does a Wii-based exercise program enhance balance*  
349 *control of independently functioning older adults? A systematic review*. Clinical Interventions  
350 In Aging, 2014. **9**: p. 1803-13.
- 351 10. Mancini, M. and F.B. Horak, *The relevance of clinical balance assessment tools to*  
352 *differentiate balance deficits*. European journal of physical and rehabilitation medicine, 2010.  
353 **46**(2): p. 239.
- 354 11. Berg, K., *Measuring balance in the elderly: development and validation of an instrument*.  
355 1992.
- 356 12. Tinetti, M.E., *Performance - oriented assessment of mobility problems in elderly patients*.  
357 Journal of the American Geriatrics Society, 1986. **34**(2): p. 119-126.
- 358 13. Duncan, P.W., et al., *Functional reach: a new clinical measure of balance*. Journal of  
359 gerontology, 1990. **45**(6): p. M192-M197.
- 360 14. Michikawa, T., et al., *One-leg standing test for elderly populations*. Journal of Orthopaedic  
361 Science, 2009. **14**(5): p. 675-685.
- 362 15. Podsiadlo, D. and S. Richardson, *The timed "Up & Go": a test of basic functional mobility for*  
363 *frail elderly persons*. Journal of the American geriatrics Society, 1991. **39**(2): p. 142-148.
- 364 16. Langley, F.A. and S.F. Mackintosh, *Functional balance assessment of older community*  
365 *dwelling adults: a systematic review of the literature*. Internet Journal of Allied Health  
366 Sciences and Practice, 2007. **5**(4): p. 13.
- 367 17. Tinetti, M.E., D. Richman, and L. Powell, *Falls efficacy as a measure of fear of falling*. Journal  
368 of Gerontology, 1990. **45**(6): p. P239-P243.

- 369 18. Powell, L.E. and A.M. Myers, *The activities-specific balance confidence (ABC) scale*. The  
370 Journals of Gerontology Series A: Biological Sciences and Medical Sciences, 1995. **50**(1): p.  
371 M28-M34.
- 372 19. Prieto, T.E., et al., *Measures of postural steadiness: differences between healthy young and*  
373 *elderly adults*. IEEE Transactions on Biomedical Engineering, 1996. **43**(9): p. 956-966.
- 374 20. Piirtola, M. and P. Era, *Force platform measurements as predictors of falls among older*  
375 *people—a review*. Gerontology, 2006. **52**(1): p. 1-16.
- 376 21. Hageman, P.A., J.M. Leibowitz, and D. Blanke, *Age and gender effects on postural control*  
377 *measures*. Archives of Physical Medicine and Rehabilitation, 1995. **76**(10): p. 961-965.
- 378 22. Moe-Nilssen, R. and J.L. Helbostad, *Trunk accelerometry as a measure of balance control*  
379 *during quiet standing*. Gait & Posture, 2002. **16**(1): p. 60-68.
- 380 23. Whitney, S., et al., *A comparison of accelerometry and center of pressure measures during*  
381 *computerized dynamic posturography: a measure of balance*. Gait & Posture, 2011. **33**(4): p.  
382 594-599.
- 383 24. Rine, R.M., et al., *Vestibular function assessment using the NIH Toolbox*. Neurology, 2013.  
384 **80**(11 ): p. S25-S31.
- 385 25. Lara, J., et al., *Towards measurement of the Healthy Ageing Phenotype in lifestyle-based*  
386 *intervention studies*. Maturitas, 2013. **76**(2): p. 189-199.
- 387 26. Godfrey, A., et al., *iCap: Instrumented assessment of physical capability*. Maturitas, 2015.  
388 **82**(1): p. 116-122.
- 389 27. Mancini, M., et al., *ISway: a sensitive, valid and reliable measure of postural control*. Journal  
390 of Neuroengineering and Rehabilitation, 2012. **9**(1): p. 1.
- 391 28. Moher, D., et al., *Preferred reporting items for systematic reviews and meta-analyses: the*  
392 *PRISMA statement*. Annals of Internal Medicine, 2009. **151**(4): p. 264-269.
- 393 29. Oxford, U.o. *The Oxford 2011 Levels of Evidence*. 2016 [cited 2016 29th June]; Available  
394 from: <http://www.cebm.net/index.aspx?o=5653>.
- 395 30. Barry, G., B. Galna, and L. Rochester, *The role of exergaming in Parkinson's disease*  
396 *rehabilitation: a systematic review of the evidence*. Journal of Neuroengineering and  
397 Rehabilitation, 2014. **11**.
- 398 31. Higgins, J.P., et al., *Measuring inconsistency in meta-analyses*. British Medical Journal, 2003.  
399 **327**(7414): p. 557-560.
- 400 32. Deeks, J.J., J. Higgins, and D.G. Altman, *Analysing data and undertaking meta - analyses*.  
401 Cochrane handbook for systematic reviews of interventions: Cochrane book series, 2008: p.  
402 243-296.
- 403 33. Chow, D.H.K. and S.K.F. Mann, *Effect of Cyber-Golfing on Balance Amongst the Elderly in*  
404 *Hong Kong: A Pilot Randomised Trial*. Hong Kong Journal of Occupational Therapy, 2015. **26**:  
405 p. 9-13.
- 406 34. Tange, H., et al. *A pilot with Exergames in Elderly Homes*. in *23rd International Conference of*  
407 *the European Federation for Medical Informatics: User Centred Networked Health Care*. 2012.
- 408 35. Pluchino, A., et al., *Pilot Study Comparing Changes in Postural Control After Training Using a*  
409 *Video Game Balance Board Program and 2 Standard Activity-Based Balance Intervention*  
410 *Programs*. Archives of Physical Medicine and Rehabilitation, 2012. **93**(7): p. 1138-1146.
- 411 36. Nicholson, V.P., et al., *Six weeks of unsupervised Nintendo Wii Fit gaming is effective at*  
412 *improving balance in independent older adults*. Journl of Aging and Physical Activity, 2015.  
413 **23**(1): p. 153-158.
- 414 37. Ray, C., et al., *The Effects of a 15-Week Exercise Intervention on Fitness and Postural Control*  
415 *in Older Adults*. Activities, Adaptation & Aging, 2012. **36**(3): p. 227-241 15p.
- 416 38. Merriman, N.A., et al., *Successful balance training is associated with improved multisensory*  
417 *function in fall-prone older adults*. Computers in Human Behavior, 2015. **45**: p. 192-203.
- 418 39. Whyatt, C., et al., *A Wii Bit of Fun: A Novel Platform to Deliver Effective Balance Training to*  
419 *Older Adults*. Games for Health Journal, 2015. **4**(6): p. 423-433.

- 420 40. Toulotte, C., C. Toursel, and N. Olivier, *Wii Fit® training vs. Adapted Physical Activities: which*  
421 *one is the most appropriate to improve the balance of independent senior subjects? A*  
422 *randomized controlled study*. *Clinical Rehabilitation*, 2012. **26**(9): p. 827-835 9p.
- 423 41. Singh, D.K.A., et al., *Effects of balance-focused interactive games compared to therapeutic*  
424 *balance classes for older women*. *Climacteric*, 2013. **16**(1): p. 141-146.
- 425 42. Sato, K., et al., *Improving Walking, Muscle Strength, and Balance in the Elderly with an*  
426 *Exergame Using Kinect: A Randomized Controlled Trial*. *Games for Health Journal*, 2015. **4**(3):  
427 p. 161-167.
- 428 43. Lai, C.-H., et al., *Effects of interactive video-game based system exercise on the balance of*  
429 *the elderly*. *Gait & Posture*, 2013. **37**(4): p. 511-515.
- 430 44. Park, E.-C., S.-G. Kim, and C.-W. Lee, *The effects of virtual reality game exercise on balance*  
431 *and gait of the elderly*. *Journal of Physical Therapy Science*, 2015. **27**(4): p. 1157-1159.
- 432 45. Yelnik, A. and I. Bonan, *Clinical tools for assessing balance disorders*. *Neurophysiologie*  
433 *Clinique/Clinical Neurophysiology*, 2008. **38**(6): p. 439-445.
- 434 46. Pardasaney, P.K., et al., *Sensitivity to change and responsiveness of four balance measures*  
435 *for community-dwelling older adults*. *Physical therapy*, 2012. **92**(3): p. 388-397.
- 436 47. Jonsson, E., M. Henriksson, and H. Hirschfeld, *Does the functional reach test reflect stability*  
437 *limits in elderly people?* *Journal of rehabilitation medicine*, 2003. **35**(1): p. 26-30.
- 438 48. Sibley, K.M., et al., *Using the systems framework for postural control to analyze the*  
439 *components of balance evaluated in standardized balance measures: a scoping review*.  
440 *Archives of physical medicine and rehabilitation*, 2015. **96**(1): p. 122-132. e29.
- 441 49. Bleakley, C.M., et al., *Gaming for Health: A Systematic Review of the Physical and Cognitive*  
442 *Effects of Interactive Computer Games in Older Adults*. *Journal of Applied Gerontology*, 2015.  
443 **34**(3): p. 166-189.
- 444 50. Donath, L., R. Rössler, and O. Faude, *Effects of Virtual Reality Training (Exergaming)*  
445 *Compared to Alternative Exercise Training and Passive Control on Standing Balance and*  
446 *Functional Mobility in Healthy Community-Dwelling Seniors: A Meta-Analytical Review*.  
447 *Sports Medicine (Auckland, N.Z.)*, 2016. **46**(9): p. 1293–1309.
- 448 51. Kümmel, J., et al., *Specificity of Balance Training in Healthy Individuals: A Systematic Review*  
449 *and Meta-Analysis*. *Sports Medicine*, 2016: p. 1-11.
- 450 52. Hwa-ann, C. and D.E. Krebs, *Dynamic balance control in elders: gait initiation assessment as*  
451 *a screening tool*. *Archives of Physical Medicine and Rehabilitation*, 1999. **80**(5): p. 490-494.
- 452 53. Skjaeret, N., et al., *Designing for Movement Quality in Exergames: Lessons Learned from*  
453 *Observing Senior Citizens Playing Stepping Games*. *Gerontology*, 2015. **61**(2): p. 186-194.
- 454 54. Pardasaney, P.K., et al., *Conceptual limitations of balance measures for community-dwelling*  
455 *older adults*. *Physical Therapy*, 2013. **93**(10): p. 1351-1368.
- 456 55. Larsen, L.H., et al., *The physical effect of exergames in healthy elderly—a systematic review*.  
457 *Games For Health: Research, Development, and Clinical Applications*, 2013. **2**(4): p. 205-212.

458