

Northumbria Research Link

Citation: Scanlan, Aaron, Berkelmans, Daniel, Vickery, Will and Kean, Crystal (2016) A Review of the Internal and External Physiological Demands Associated With Batting in Cricket. *International Journal of Sports Physiology and Performance*, 11 (8). pp. 987-997. ISSN 1555-0265

Published by: Human Kinetics

URL: <http://dx.doi.org/10.1123/ijsp.2016-0169> <<http://dx.doi.org/10.1123/ijsp.2016-0169>>

This version was downloaded from Northumbria Research Link:
<http://nrl.northumbria.ac.uk/29157/>

Northumbria University has developed Northumbria Research Link (NRL) to enable users to access the University's research output. Copyright © and moral rights for items on NRL are retained by the individual author(s) and/or other copyright owners. Single copies of full items can be reproduced, displayed or performed, and given to third parties in any format or medium for personal research or study, educational, or not-for-profit purposes without prior permission or charge, provided the authors, title and full bibliographic details are given, as well as a hyperlink and/or URL to the original metadata page. The content must not be changed in any way. Full items must not be sold commercially in any format or medium without formal permission of the copyright holder. The full policy is available online: <http://nrl.northumbria.ac.uk/policies.html>

This document may differ from the final, published version of the research and has been made available online in accordance with publisher policies. To read and/or cite from the published version of the research, please visit the publisher's website (a subscription may be required.)

www.northumbria.ac.uk/nrl



1 **A review of the internal and external physiological**
2 **demands associated with batting in cricket**

3
4 **Submission Type:** Brief Review

5
6 Aaron T. Scanlan,^{1,2} Daniel M. Berkelmans,¹ William M.
7 Vickery,³ and Crystal O. Kean¹

8
9 ¹School of Medical and Applied Sciences, Building 81, Central
10 Queensland University, Bruce Highway, Rockhampton,
11 Australia.

12 ²Human Exercise and Training Laboratory, Building 81,
13 Central Queensland University, Bruce Highway, Rockhampton,
14 Australia.

15 ³Department of Sport, Exercise and Rehabilitation,
16 Northumberland Building, Northumbria University, Newcastle
17 Upon Tyne, United Kingdom.

18
19 ✉ Dr Aaron T Scanlan
20 Human Exercise and Training Laboratory Director
21 Central Queensland University
22 Bruce Highway
23 Rockhampton, Queensland, 4702
24 Australia
25 Phone (international): +61 7 4923 2538
26 Fax (international): +61 7 4930 6781
27 Email: a.scanlan@cqu.edu.au

28
29 **Preferred Running Head:** Batting demands in cricket

30 **Manuscript word count:** 4607

31 **Abstract word count:** 226

32 **Number of References:** 46

33 **Number of Figures and Tables:** 3 tables

34 **ABSTRACT**

35 Cricket is a popular, international team sport with various game
36 formats ranging from long duration multi-day tests to short
37 duration Twenty20 game-play. The role of batsmen is critical to
38 all game formats with differing physiological demands imposed
39 during each format. Investigation of the physiological demands
40 imposed during cricket batting has historically been neglected
41 with much of the research focusing on bowling responses and
42 batting technique. A greater understanding of the physiological
43 demands of the batting role in cricket is required to assist
44 strength and conditioning professionals and coaches with the
45 design of training plans, recovery protocols, and player
46 management strategies. This brief review provides an updated
47 synthesis of the literature examining the internal (e.g. metabolic
48 demands, heart rate) and external (e.g. activity work rates)
49 physiological responses to batting in the various game formats
50 as well as simulated play and small-sided games training.
51 While few studies in this area exist, the summary of data
52 provides important insight regarding physiological responses to
53 batting, and highlights that more research on this topic is
54 required. Future research is recommended to combine internal
55 and external measures during actual game-play as well as
56 comparing different game formats and playing levels. In
57 addition, understanding the relationship between batting
58 technique and physiological responses is warranted to gain a
59 more holistic understanding of batting in cricket as well as
60 develop appropriate coaching and training strategies.

61

62 **Key Words:** batsmen; heart rate; lactate; RPE; GPS; time-
63 motion analysis.

64 **INTRODUCTION**

65 Cricket is an international team sport with 105 member
66 countries recognized by the International Cricket Council,
67 spanning Africa, the Americas, Asia, East-Asia-Pacific, and
68 Europe.¹ Modern cricket involves two teams of 11 players and
69 is played on a field containing a pitch with a set of three
70 wooden stumps at each end.² The focal point of cricket is the
71 contest between bat and ball, with three primary functional
72 roles being identified: batting, bowling, and fielding. The
73 objective when bowling and fielding is to dismiss 10 batsmen
74 (10 wickets) while minimizing the amount of runs scored.
75 Conversely, the batting team aims to strike the ball through or
76 over the field, scoring runs if the ball reaches the boundary (4
77 or 6 runs) or if the batsmen run the length of the pitch
78 (individual runs given per length completed). Ultimately, the
79 batting team attempts to accumulate more runs than the
80 opposing team. Batsmen can be dismissed by various means
81 including being bowled, caught, stumped, run out, leg before
82 wicket (stumps), and hitting the stumps.

83 In recent times there has been an increase in the volume
84 of cricket played across the annual season, as well as enhanced
85 commercialization of the sport. This evolution has promoted
86 a more professional, structured approach to travelling, training,
87 game preparation, and recovery using scientific concepts.
88 Consequently, greater research attention is being given to
89 various aspects of cricket to better understand the demands
90 placed on players during games, simulated play, and training.³⁻
91 ¹² Researchers have primarily focused on examination of
92 bowling and fielding in cricket, resulting in numerous focused
93 reviews in this area.¹³⁻¹⁷ Although less inquiry is available
94 regarding batting in cricket, a greater understanding of the
95 physiological responses and technical attributes associated with
96 this role has emerged in the literature across several
97 examinations in the past decade. Consequently, there is a need
98 for a contemporary synthesis of the literature regarding batting
99 responses in cricket, with the only available review examining
100 physiological responses conducted in 2000.¹⁸ In turn, a more
101 recent review focusing on batting technique and biomechanics
102 was conducted in 2012.¹⁹ As very little biomechanical research
103 related to cricket batting has been conducted since the review
104 by Penn and Spratford,¹⁹ the aim of this review is to focus on
105 synthesizing the literature related to internal and external
106 physiological responses to batting in cricket.

107
108 **SEARCH STRATEGY**

109 A comprehensive search of the online library databases
110 provided by Central Queensland University, as well as Google
111 Scholar and PubMed was conducted to locate potential sources
112 for this review. No time restrictions were set and the following
113 combination of terms were entered: ‘cricket’, ‘batting’,

114 'batsmen', 'responses', 'physiology', 'heart rate', 'metabolic',
115 'perceptual', and 'activity'. Reference lists of identified
116 publications were searched to locate additional sources. The
117 quality of retrieved publications were assessed using various
118 items from the critical review form for quantitative studies
119 developed by Law et al.²⁰ Given the observational nature of the
120 included studies, aspects related to study design, intervention,
121 and drop-outs contained in the original critical review form
122 were excluded in our evaluation. Evaluations for each
123 publication are presented in Table 1, with study limitations
124 provided in further detail. Almost all of the retrieved
125 publications (89%) scored ≥ 7 out of 10 in the evaluation.

126

127

INSERT TABLE 1 AROUND HERE

128

129 **GAME FORMAT**

130 Various factors can alter the physiological responses to batting
131 in cricket, the most prominent being game format. The
132 traditional game format in cricket involves multi-day
133 competition with up to 5 days (90 overs per day) for each team
134 to bat and bowl across two separate innings. A winning
135 outcome is achieved if a team dismisses 20 batsmen (2 innings
136 x 10 wickets) for a lower aggregate run total. A shift toward
137 shorter games emerged in the 1970s with the game outcome
138 produced in a single day.²¹ The One-Day format adopts a
139 similar approach to multi-day competition, but each team only
140 bats and bowls across a single innings (1 innings x 10 wickets)
141 and games are limited to 50 overs per team. The trend of
142 producing shorter game formats continued with Twenty20
143 cricket being developed in 2003. Twenty20 cricket is played
144 across 20 overs per team, with each game lasting approximately
145 3 hours.

146 Player requirements in cricket can be altered across the
147 different game formats.^{8,9,22} For instance, batting performance
148 during a Twenty20 game necessitates a higher rate of scoring
149 strokes compared to a multi-day game where less time
150 restrictions are encountered.²³ In turn, these temporal
151 constraints promote a higher urgency for attacking play and
152 running between wickets to score runs during shorter game
153 formats. Thus when gathering evidence regarding batting
154 responses from the available literature, readers should be aware
155 of the game format being investigated as each format is likely
156 to impose unique requirements upon players. We have included
157 all game formats in this review.

158

159 **PHYSIOLOGICAL RESPONSES TO BATTING IN** 160 **CRICKET**

161 Understanding the physiological requirements of team sports
162 forms the basis of designing conditioning programs which
163 promote adaptation in players to optimize physical

164 preparedness for competition.⁸ With the growing application of
165 sport science and evolution of advanced measurement
166 techniques, increased research has been conducted examining
167 the physiological responses to batting in cricket. The primary
168 measures examined in the literature can be broadly categorized
169 as internal and external responses.

170

171 **Internal Responses**

172 Measurement of internal responses to game-play and training
173 provide important insight into the physiological stress imposed
174 upon athletes across various body systems.²⁴ To date, a range of
175 internal responses to batting in cricket have been reported in the
176 literature. Specifically, internal measures primarily examined
177 include metabolic responses, heart rate (HR), blood lactate
178 concentration ($[BLa^-]$), and rating of perceived exertion (RPE).
179 A summary of the internal physiological responses observed
180 during batting in cricket studies is presented in Table 2.

181

182 ***INSERT TABLE 2 AROUND HERE***

183

184 *Metabolic Responses*

185 Early research reporting on the physiological responses to
186 batting utilized portable calorimetry during batting tasks during
187 practice to estimate the energy expenditure during actual game-
188 play.²⁵ Fletcher²⁵ estimated energy expenditure to be $648 \text{ kJ}\cdot\text{h}^{-1}$
189 for batsmen based upon running between wickets for 26.6 runs
190 per hour during international multi-day cricket. The primary
191 limitations of this research were the inclusion of drink breaks,
192 lunch breaks, and time spent waiting to bat in calculations.
193 These limitations, likely underestimating the requirements of
194 competition and this notion is substantiated by more a recent
195 investigation which shows a higher energy expenditure during
196 batting in cricket.³

197 To analyze the physiological responses to batting,
198 Christie et al.³ utilized a single 7-over bout, with 30-s and 1-
199 min rest periods between deliveries and overs respectively. A
200 live bowler was used and batsmen completed 2 runs every 3
201 balls (28 runs across the protocol).³ This configuration was
202 based on observations made during One-Day international
203 games, and allowed a more definitive metabolic assessment of
204 batting to be conducted with direct use of a portable metabolic
205 analyzer during batting. Accordingly, an energy expenditure of
206 $2,536 \text{ kJ}\cdot\text{h}^{-1}$ was recorded in first team university batsmen
207 during the protocol.³ Furthermore, a mean oxygen uptake
208 (VO_2) of $26.7 \pm 1.4 \text{ mL}\cdot\text{kg}^{-1}\cdot\text{min}^{-1}$ and respiratory exchange
209 ratio (RER) of 1.05 ± 0.05 were also observed.³ While these
210 data suggest a predominant recruitment of aerobic metabolic
211 pathways (mean $\text{VO}_2 \approx 50\% \text{ VO}_{2\text{max}}$) during batting, the
212 increase in RER to 1.09 ± 0.05 following the four sprints in the
213 first over (out of 7 overs) indicates anaerobic energy systems

214 are also important during the short high-intensity running bouts
215 and likely contribute to sustained elevated physiological
216 responses thereafter. Furthermore, given RER was consistently
217 >1 following the first over, Christie et al.³ concluded that
218 carbohydrates were the preferred source of energy substrate
219 during batting tasks.

220 The discrepancies in metabolic responses observed in
221 early research compared with more recent data are likely due to
222 temporal changes in game demands and/or greater
223 physiological stress being imposed upon batsmen during
224 shorter game formats.⁸ Nevertheless, the collective research
225 emphasizes the lack of metabolic data representative of batting
226 tasks in cricket, particularly during game-play. This lack of
227 inquiry is likely due to the logistical limitations associated with
228 administering metabolic measurement techniques using bulky
229 and costly equipment during actual batting tasks. Consequently,
230 many researchers have opted to use telemetric heart rate (HR)
231 devices to estimate the metabolic demands experienced by
232 batsmen during games and training given the greater practical
233 utility and indirect indication of aerobic energy system
234 recruitment accompanying HR monitoring.^{26,27}

235

236 *Heart Rate*

237 HR measurement has emerged as the most popular approach to
238 monitor the internal responses of batsmen in cricket.
239 Researchers have provided HR measurements for batsmen
240 across actual games, simulated play, and training scenarios.
241 Specifically, mean absolute HR of 149 ± 17 beats·min⁻¹ have
242 been observed during Twenty20 games in second-tier
243 international batsmen,⁸ while responses of 144 ± 13 beats·min⁻¹
244 and 159 ± 12 beats·min⁻¹ were reported in second-tier
245 international⁸ and first team club batsmen²⁸ during One-Day
246 games, respectively. The reported HR data also demonstrate
247 some important time-course responses with spikes in HR
248 showing players reach 97% of HR_{max} and spend considerable
249 proportions of batting time (63%) working above an intensity
250 of 75% HR_{max}.²⁸ Thus, while the overall mean HR in batsmen
251 might be considered *hard* (73-78% of age-predicted HR_{max}),²⁹
252 periodic bursts of high-intensity efforts are required which
253 suggests phosphocreatine (PCr) stores and the glycolytic
254 energy system are relied upon for energy provision, placing the
255 body into an oxygen deficient state and producing metabolic
256 by-products.³⁰ Consequently, the oxidative energy system
257 likely plays major roles not only in adenosine triphosphate re-
258 synthesis, but also in PCr restoration and lactate oxidation
259 during lower-intensity activity.³¹ Moreover, comparisons across
260 studies indicate that game format, playing level, and
261 competition locality are likely to influence the energetic
262 demands of batting as evidenced by varied HR responses.

263 While limited HR data exist representative of actual
264 game-play, a wider scope of studies have used simulated play
265 and training scenarios to measure the internal responses of
266 batsmen.^{3,5,6,28,32,33} Novel approaches such as BATEX^{5,6,32} and
267 The Battlezone^{28,33} have been developed to replicate batting
268 requirements during game-play. BATEX is a simulated batting
269 innings consisting of 6 x 21-min stages with embedded rest
270 periods, producing a performance duration indicative of scoring
271 100 runs in a One-Day international game.⁵ Each stage consists
272 of 5 overs in a net-based practice setting.⁵ Timings of ball
273 deliveries are based on archived game data, with each stage
274 containing different running requirements to represent typical
275 distributions at different game stages.⁵ In contrast, The
276 Battlezone is a small-sided game (SSG) whereby batsmen
277 complete pre-determined frequencies of 6 bouts of 8 overs
278 separated by 5 min of rest.³³ The Battlezone is representative of
279 play in close proximity to the pitch within the inner circle
280 (27.4-m diameter).³³ A net encloses the inner circle which
281 contains two batsmen, three fielders, two bowlers (who
282 alternate between overs), and a wicket-keeper.³³ Generally, the
283 aim for batsmen during The Battlezone is to score as many runs
284 as possible with encouragement to hit the ball along the ground
285 and not over the net³³; however coaches have the option to alter
286 this approach to suit session objectives.

287 Reported HR across each of these approaches, as well
288 as the simulated 7-over bout developed by Christie et al.³ have
289 been shown to vary across studies (BATEX: 130-144
290 beats·min⁻¹; The Battlezone: 164 ± 12 beats·min⁻¹; 7-over bout:
291 145 ± 11 beats·min⁻¹).^{3,5,6,28,32,33} Furthermore, HR during
292 BATEX and 7-over bout increased with protocol progression,
293 reaching 147-159 beats·min⁻¹ by the final stage/over.^{3,5,6,32}
294 Subsequently, these simulated protocols appear to invoke
295 comparable or greater HR than those evident during Twenty20
296 and One-Day game-play.^{8,28} Thus, the established simulation
297 and SSG approaches in the literature may be particularly useful
298 as training stimuli for batsmen to optimize cardiovascular
299 adaptation in preparation for game-play. However, these
300 observations should be treated with caution given the existing
301 data during simulated play and SSG training are indicative of
302 club players (first to fourth teams),^{3,5,6,28,32,33} while findings
303 during actual game-play are representative of club (first and
304 second teams) and international players.^{8,28} The higher level
305 players likely possessed superior levels of training experience
306 and fitness which might have influenced the HR results
307 observed.²⁷ Indeed, Houghton et al.³² showed lower-level club
308 players (third and fourth teams) to produce greater HR
309 responses (4-8 beats·min⁻¹) across all stages of BATEX than
310 higher-level club players (first and second teams).

311

312 *Blood Lactate Concentration*

313 While metabolic and HR data permit assumptions to be made
314 regarding the reliance on anaerobic metabolic pathways during
315 batting in cricket, [BLa⁻] is a more direct indicator of energy
316 production from anaerobic glycolysis.³⁴ Subsequently, research
317 reporting on [BLa⁻] highlights the variable recruitment of
318 anaerobic metabolism during batting across simulated play and
319 SSG training. More precisely, Houghton et al.³² observed
320 comparable [BLa⁻] in first and second team club batsmen ($3.2 \pm$
321 1.6 to 4.5 ± 1.6 mmol·L⁻¹) and third and fourth team club
322 batsmen (3.0 ± 0.9 to 4.1 ± 1.2 mmol·L⁻¹) during BATEX. The
323 range of responses indicate that during BATEX, batsmen work
324 at intensities above those associated with anaerobic threshold
325 using fixed [BLa⁻] (3-4 mmol·L⁻¹).³⁵ These responses were also
326 higher than those evident during SSG training (The Battlezone)
327 in first and second team club batsmen (1.8 ± 0.7 to 3.2 ± 1.4
328 mmol·L⁻¹).^{28,33} Collectively, these data contradict the higher
329 cardiovascular intensities (HR responses) during The
330 Battlezone, as the [BLa⁻] results indicate that batsmen do not
331 reach intensities concomitant with anaerobic threshold as
332 readily during The Battlezone compared to BATEX. These
333 variations might be due to the protocols adopted across studies.
334 The Battlezone bouts lasted between 14-18 min and batting
335 performance was self-determined, with running between
336 wickets conducted in an ad-hoc manner during live game
337 scenarios.^{36,37} In contrast, BATEX bouts were performed across
338 6 x 21-min stages with included recovery periods (total = 2 h
339 20 min) and increased running demands periodically elicited at
340 maximum exertion with protocol progression.⁵ Alternatively,
341 the lower playing levels of participants examined during
342 BATEX (club-level from first to fourth team) might have
343 possessed inferior aerobic fitness compared with those
344 completing The Battlezone (club-level first and second teams).
345 In turn, this would promote recruitment of anaerobic metabolic
346 pathways for energy provision at lower relative HR intensities
347 in participants completing BATEX, leading to greater lactate
348 accumulation. However, this postulation remains speculative as
349 aerobic fitness measures were not provided in these
350 studies.^{28,32,33}

351

352 *Rating of Perceived Exertion*

353 The reported HR and [BLa⁻] data provide useful insight
354 regarding aerobic and anaerobic metabolic recruitment
355 respectively. In turn, RPE has been suggested to be a global
356 indicator of exercise demands encompassing both of these
357 measures during intermittent activity.³⁸ RPE responses have
358 been recorded in batsmen using 1-10 and 6-20 Borg Scales.
359 Reported RPE scores of 4-5 (1-10 scale) and 10-17 (6-20 scale)
360 have been observed across The Battlezone^{28,33} and BATEX
361 protocols,^{5,6,32} respectively. Furthermore, RPE of 5 ± 2 (1-10
362 scale) has been reported during One-Day game-play in first

363 team club batsmen.²⁸ These observations reflect *fairly light* to
364 *very hard* intensities, which tend to overlap and/or exceed the
365 descriptive intensity zones observed for HR responses,²⁹ adding
366 credence to the combined anaerobic and aerobic contribution to
367 perceived exertion in cricket.³⁸ Time-course comparisons
368 across BATEX in first and second team club batsmen
369 suggested that RPE increased as a function of duration rather
370 than intensity, given greater increases in perceptions of effort
371 were observed than other internal and external markers of
372 intensity.⁶ Moreover, non-significant differences in RPE
373 between One-Day game-play and SSG training (The
374 Battlezone) were reported by Vickery et al.²⁸ suggesting batting
375 stimuli presented during each of these formats exert similar
376 perceptual demands. Given the practical benefits in gathering
377 and interpreting RPE scores,³⁹ it is apparent more research is
378 needed to establish the utility of this approach in representing
379 the internal demands associated with batting in cricket,
380 particularly during games across various formats.

381

382 **External responses**

383 The added measurement of activity demands gives detail about
384 the external responses to batting, from which further
385 physiological inferences can be made. Technological
386 advancements in video-based approaches and the development
387 of micro-technologies, such as global positioning system (GPS)
388 units, permit reliable and valid measurement of activity
389 responses in cricket.^{22,40,41} These approaches are becoming
390 more routinely used to monitor the external physiological
391 responses of players and thus an increasing number of
392 researchers have reported these data for batsmen during cricket
393 games, simulated play, and training scenarios. Activities
394 performed are typically categorized according to intensity,
395 whereby most studies adopt the following criteria:
396 standing/walking = $\leq 2 \text{ m}\cdot\text{s}^{-1}$; jogging = $2.01\text{-}3.5 \text{ m}\cdot\text{s}^{-1}$; running
397 = $3.51\text{-}4 \text{ m}\cdot\text{s}^{-1}$; striding = $4.01\text{-}5 \text{ m}\cdot\text{s}^{-1}$; and sprinting = $>5 \text{ m}\cdot\text{s}^{-1}$
398 ^{1, 5,8,28,33,42} In turn, insight regarding activity frequencies,
399 durations, and distances within these categories has been
400 provided for batsmen. A summary of the external physiological
401 responses observed during batting in cricket studies is
402 presented in Table 3.

403

404 ***INSERT TABLE 3 AROUND HERE***

405

406 *Activity Frequencies*

407 Using video-based time-motion analyses, Duffield and
408 Drinkwater²² examined the external responses of batsmen
409 across 50-, 80- and 100-run innings during One-Day and multi-
410 day international games. The comparative analyses showed
411 consistent jogging, striding, and sprinting frequencies across
412 game formats for each innings category. Furthermore,

413 significantly greater standing, and walking frequencies were
414 observed for multi-day compared to One-Day game-play for
415 each innings category. Similarly, Petersen et al.⁸ showed
416 greater frequency of high intensity (running, striding and
417 sprinting) during Twenty20 compared to One-Day and multi-
418 day game-play as well as One-Day compared to multi-day
419 game-play but did not report frequency of low-intensity
420 activities (standing, walking, or jogging). With regards to the
421 simulated game play and SSG (The Battlezone), Vickery et
422 al.^{28,33} reported The Battlezone has higher frequency of
423 sprinting and high-intensity activities compared to One-Day
424 game-play, with the frequency of sprinting comparable to that
425 noted by Petersen et al.^{8,42} during Twenty20 (Table 3).
426 However, the frequency of high-intensity activities during The
427 Battlezone is greater than that presented in actual game-play
428 studies (Table 3).

429 Duffield and Drinkwater²² also showed an increase in
430 shot frequency during the multi-day versus One-Day game-play
431 for the 80- and 100-run innings (80 runs: 95 ± 17 vs. 122 ± 23 ;
432 100 runs: 105 ± 18 vs. 151 ± 22), indicating that a greater
433 number of shots are required to reach pre-determined scores
434 during longer duration multi-day than One-Day games. Limited
435 information regarding the frequency of attacking and defensive
436 shots exists with only Vickery et al.²⁸ noting that during One-
437 Day game-play the frequency of attacking shots was $21 \pm 4 \cdot \text{h}^{-1}$
438 compared to $12 \pm 5 \cdot \text{h}^{-1}$ defensive shots. To date, no studies
439 have compared shot type frequency between different game
440 formats.

441 Based on the present research it appears batting during
442 multi-day games has a greater contribution from low-intensity
443 activity than shorter games (One-Day or Twenty20). The
444 current research also shows a greater frequency of recovery
445 activities around high-intensity bouts during multi-day cricket,
446 reflecting the more attacking style of play through an
447 augmented shot frequency in shorter game formats. However,
448 data regarding activity frequencies for batting during Twenty20
449 cricket is limited with only Petersen et al.^{8,42} reporting the
450 frequency of sprinting and high intensity efforts (Table 3).
451 Further research comparing the influence of game formats and
452 shot types on activity frequencies during batting is warranted to
453 improve the understanding in this area.

454 455 *Activity Durations*

456 Duration data provide useful information regarding the
457 proportions of game-play or training spent working at different
458 intensities. The durations spent performing different activities
459 during batting have been provided across all game formats and
460 concur with the findings for activity frequencies. Petersen et
461 al.⁴² utilized GPS technology to measure the activity demands
462 imposed upon state-level batsmen scaled to 30 min of activity

463 during Twenty20 games. Activity categories were grouped as
464 low- and high-intensity, with players spending more time
465 engaged in walking and jogging (low-intensity activity) than
466 running, striding, and sprinting (high-intensity activity). Across
467 longer game formats, Duffield and Drinkwater²² reported
468 activity durations for 50-, 80- and 100-run innings during
469 international One-Day and multi-day games. Consistent
470 jogging, striding, and sprinting durations were apparent across
471 game formats for each run innings category. For 50-run innings
472 there was also consistency in walking and shot (0.9 ± 0.2 vs 1.1
473 ± 0.4 min) durations between game formats. In contrast, there
474 was significantly greater walking durations between game
475 formats for the 80- and 100- run innings categories and shot
476 durations (80 runs: 95 ± 17 vs. 122 ± 23 min; 100 runs: $105 \pm$
477 18 vs. 151 ± 22 min). In addition, significantly greater standing
478 durations were evident between game formats for each run
479 innings category (Table 3). Overall, innings durations during
480 One-Day games were significantly shorter than multi-day
481 games for 50- and 100-run innings (50 runs: 84.5 ± 17.7 vs
482 108.9 ± 26.6 min; 100 runs: 135.5 ± 21.4 vs 213.4 ± 31.9
483 min).²²

484 The literature indicates that the majority of batting time
485 is spent engaged in low-intensity activity. When grouped
486 according to low- and high-intensity, 95.5%, 97.7%, and 98.6%
487 of batting time were spent engaged in low-intensity activity
488 during Twenty20, One-Day, and multi-day cricket
489 respectively.^{22,42} While the volume of high-intensity activity
490 during batting is comparable across One-Day and multi-day
491 game formats (2.2 vs. 2.1 min in 50-run innings), greater
492 standing and walking activity were apparent during multi-day
493 games predisposing to larger work:rest ratios in batsmen during
494 this format (One-Day: 1:47 s; multi-day: 1:67 s).²²
495 Furthermore, batting during Twenty20 cricket invoked an even
496 lower work:rest ratio (1:24 s) than both One-Day and multi-day
497 formats.⁴² Separately, Petersen et al.⁸ observed comparable
498 work:rest ratios across One-Day (1:50 s) and multi-day (1:61 s)
499 games to those reported in international batsmen by Duffield
500 and Drinkwater²² with a considerably higher work:rest ratio
501 evident during Twenty20 game-play (1:38 s). Likewise,
502 Vickery et al.²⁸ observed a higher work:rest ratio in One-Day
503 games (1:66 s) than those reported in other studies.
504 Discrepancies across studies might be related to playing level
505 of the batsmen investigated as second-tier international,⁸ state-
506 level,⁴² and club-level²⁸ players have been examined.

507 Nevertheless, the available evidence suggests as the
508 duration of the game decreases, external physiological intensity
509 increases primarily through a reduction in recovery time
510 between high-intensity efforts. Given that PCr depletion has
511 been proposed as a prominent fatigue mechanism during
512 intermittent exercise, longer recovery periods would promote

513 greater PCr restoration between high-intensity bouts to
514 optimize performance maintenance.⁴³ Consequently, other
515 fatigue mediators such as glycogen depletion, dehydration, or
516 neural mechanisms might be more influential during longer
517 game formats.²²

518

519 *Activity Distances*

520 Distance data have also been provided in all game formats
521 across varied playing levels for batsmen in cricket. During
522 state-level Twenty20 games, Petersen et al.⁴² recorded various
523 distances for different activities when scaled to 30 min of
524 batting with a greater distance covered during walking
525 activities than jogging, running, striding, or sprinting.
526 Following this study, Petersen et al.⁸ compared the activity
527 demands imposed upon second-tier international batsmen
528 during Twenty20, One-Day, and multi-day game formats.
529 Activity distances were scaled and presented as $\text{m}\cdot\text{h}^{-1}$ to
530 provide comparable data across game formats. Batsmen
531 completed greater (*moderate* to *large*) relative distances
532 jogging, running, striding, and sprinting during Twenty20 than
533 multi-day games. Furthermore, batsmen covered greater
534 (*moderate*) relative distances jogging, striding, and sprinting
535 during One-Day than multi-day games. Analogous overall
536 relative distances were also observed between Twenty20 and
537 One-Day games, with lower measures recorded during multi-
538 day games (Table 3).⁸ Following a similar approach using GPS
539 technology, Vickery et al.²⁸ reported lower overall relative
540 distances for club-level batsmen during One-Day games than
541 across all formats observed by Petersen et al..^{8,42} Overall
542 distance was further analyzed according to low- and high-
543 intensity activity and again showed the majority of distance
544 covered was covered while engaged in low-intensity activity²⁸.

545 Together, the distance data reported during actual game-
546 play shows that shorter game formats (Twenty20 and One-Day)
547 carry higher work rates ($\text{m}\cdot\text{h}^{-1}$) than longer formats (multi-
548 day).^{8,28,42} Interestingly, examinations of Twenty20 cricket
549 showed wide variation in activity distances, with 1.6-2.4 times
550 the work rate evident across activities during state-level⁴²
551 compared to second-tier international⁸ game-play. These
552 differences highlight that the external physiological demands
553 imposed upon batsmen during Twenty20 cricket might be
554 highly variable. Furthermore, across separate studies, work rate
555 decreased with the playing level investigated. Specifically,
556 second-tier international players demonstrated the highest work
557 rates ($\approx 2.6 \text{ km}\cdot\text{h}^{-1}$) during Twenty20 and One-Day games,⁸
558 followed by state-level batsmen ($2.4 \text{ km}\cdot\text{h}^{-1}$) during
559 Twenty20,⁴² and club-level batsmen ($1.9 \text{ km}\cdot\text{h}^{-1}$) during One-
560 Day games.²⁸

561 In addition to observations made during game-play,
562 distance data have also been provided using GPS units for

563 simulated play and SSG training in batsmen.^{5,28,33} Comparable
564 total work rates were evident across BATEX in club-level
565 players ($2.2 \pm 0.2 \text{ km}\cdot\text{h}^{-1}$) compared with One-Day ($1.9\text{-}2.5$
566 $\text{km}\cdot\text{h}^{-1}$) and multi-day ($2.1 \pm 0.6 \text{ km}\cdot\text{h}^{-1}$) game-play.^{5,8,28}
567 However, the work rates during BATEX were lower than those
568 reported for Twenty20 ($2.4\text{-}4.9 \text{ km}\cdot\text{h}^{-1}$).^{8,42} Analyzed further,
569 BATEX imposed lower relative distances during low-intensity
570 activities (standing, walking, and jogging) and consistent or
571 greater high-intensity demands (running, striding, and
572 sprinting) than all game formats (Table 3).^{5,8,28,42} Thus, BATEX
573 appears to match the overall work rates and exceed the high-
574 intensity demands reported for batsmen during One-Day and
575 multi-day game-play, while also matching the high-intensity
576 work rates seen during Twenty20 cricket. Consequently,
577 BATEX might hold useful utility across all game formats as an
578 assessment tool to gauge the preparedness of batsmen for
579 different competitions as well as a training stimulus to
580 adequately prepare batsmen for game demands. Conversely,
581 batting during SSG training has been shown to elicit
582 considerably higher work rates across all activity categories
583 than BATEX in club-level batsmen.^{28,33}

584 Vickery et al.^{28,33} reported batsmen to cover between
585 $3.8\text{-}3.9 \text{ km}\cdot\text{h}^{-1}$ across 14-18-min bouts of The Battlezone,
586 including $3.3\text{-}3.4 \text{ km}\cdot\text{h}^{-1}$ and $0.6\text{-}0.7 \text{ km}\cdot\text{h}^{-1}$ performing low-
587 and high-intensity activity, respectively. These work rates are
588 greater than those reported during BATEX⁵ and game-play^{8,28}
589 (Table 3). It has been proposed that the fewer number of
590 fielders in the game-play scenarios encountered during The
591 Battlezone might have permitted batsmen to score more freely
592 and the protocol objectives might promote an attacking mind-
593 set to secure as many runs as possible across the short playing
594 durations by executing frequent high-intensity sprints.³³
595 Comparisons with game-play across studies confirm the
596 practical usefulness of SSG training to elicit elevated work
597 rates and provide a beneficial training stimulus for batsmen in
598 preparation for all game formats.

599

600 CONCLUSIONS

601 Findings pertaining to the internal and external physiological
602 responses during batting in cricket vary between game format,
603 as well as simulated play and SSG training. The collective
604 works in this area provide important insight regarding player
605 responses to batting, and highlight the need for more research
606 on this topic, particularly combining internal and external
607 measures during actual game-play, comparing different game
608 formats and playing levels. Investigation of fatigue-mediating
609 mechanisms during batting across games are also encouraged
610 as well as studies examining responses to different shot types
611 (attacking vs defensive). The physiological demands of batting
612 should be considered in combination with other responses, as

613 the importance of technique to batting performance has been
614 reiterated across various sources^{19,23,28,44-46} and was not covered
615 in the present review. Future studies should examine the
616 physiological responses to batting and biomechanical attributes
617 of batting technique to provide greater insight into the
618 relationship of these variables and overall performance.
619 Further, given much of the available data (Table 1) has been
620 provided during simulation and games training scenarios, more
621 research examining player responses during actual game-play is
622 required.

623

624 PRACTICAL APPLICATIONS

625 The data synthesized in this review provide a useful reference
626 for internal and external physiological stimuli relative to game
627 format as well as simulation/training protocol for strength and
628 conditioning professionals and coaches to use when developing
629 training plans, recovery protocols, and player management
630 strategies to best prepare players for competition. Specifically,
631 work:rest ratio data highlight physiological intensity is
632 heightened across shorter game durations through a reduction
633 in recovery time between high-intensity efforts. Thus player
634 conditioning plans should account for these metabolic
635 variations and be adjusted to best prepare players for specific
636 game formats across the season. In addition, variations in work
637 rates across playing levels suggest that training and assessment
638 approaches relative to playing level are warranted.
639 Conditioning drills might incorporate batting exercise
640 simulation (BATEX) and SSG training (The Battlezone), which
641 appear to provide adequate physiological overload to prepare
642 for the batting demands associated with all game formats.

643

644 REFERENCES

- 645 1. International Cricket Council. *Members overview*.
646 Available at [http://www.icc-cricket.com/about/96/icc-](http://www.icc-cricket.com/about/96/icc-members/overview)
647 [members/overview](http://www.icc-cricket.com/about/96/icc-members/overview), Accessed November 1 2015.
- 648 2. Scalmer S. Cricket, imperialism and class domination.
649 *WorkingUSA*. 2007;10(4):431-442.
- 650 3. Christie C, Todd A, King G. Selected physiological
651 responses during batting in a simulated cricket work bout:
652 A pilot study. *J Sci Med Sport*. 2008;11(6):581-584.
- 653 4. Elliott B, Baker J, Foster D. The kinematics and kinetics of
654 the off-drive and on-drive in cricket. *Aus J Sci Med Sport*.
655 1993;25(2):48-54.
- 656 5. Houghton L, Dawson B, Rubenson J, Tobin M. Movement
657 patterns and physical strain during a novel, simulated
658 cricket batting innings (BATEX). *J Sports Sci*.
659 2011;29(8):801-809.
- 660 6. Pote L, Christie C. Selected physiological and perceptual
661 responses during a simulated limited overs century in non-
662 elite batsmen. *Eur J Sport Sci*. In press.

- 663 7. Petersen C, Pyne D, Dawson B, Kellett A, Portus M.
664 Comparison of training and game demands of national
665 level cricketers. *J Strength Cond Res.* 2011;25(5):1306-
666 1311.
- 667 8. Petersen C, Pyne D, Dawson B, Portus M, Kellett A.
668 Movement patterns in cricket vary by both position and
669 game format. *J Sports Sci.* 2010;28(1):45-52.
- 670 9. Petersen C, Pyne D, Portus M, Dawson B. Comparison of
671 player movement patterns between 1-day and test cricket. *J*
672 *Strength Cond Res.* 2011;25(5):1368-1373.
- 673 10. McNamara D, Gabbett T, Chapman P, Naughton G,
674 Farhart P. Variability of PlayerLoad, bowling velocity, and
675 performance execution in fast bowlers across repeated
676 bowling spells. *Int J Sports Physiol Perform.*
677 2015;10(8):1009-1014.
- 678 11. Orchard J, Blanch P, Paoloni J, Kountouris A, Sims K,
679 Orchard J, Brukner P. Fast bowling match workloads over
680 5–26 days and risk of injury in the following month. *J Sci*
681 *Med Sport.* 2015;18(1):26-30.
- 682 12. Peterson C, Pyne D, Portus M, Karppinen M, Dawson B.
683 Variability in movement patterns during one day
684 internationals by a cricket fast bowler. *Int J Sports Physiol*
685 *Perform.* 2009;4(2):278-281.
- 686 13. MacDonald D, Cronin J, Mills J, McGuigan M, Stretch R.
687 A review of cricket fielding requirements. *S Afr J Sports*
688 *Med.* 2013;25(3):87-92.
- 689 14. Bartlett R, Stockill N, Elliott B, Burnett A. The
690 biomechanics of fast bowling in men's cricket: A review. *J*
691 *Sports Sci.* 1996;14(5):403-424.
- 692 15. Johnstone J, Mitchell A, Hughes G, Watson T, Ford P,
693 Garrett A. The athletic profile of fast bowling in cricket: A
694 review. *J Strength Cond Res.* 2014;28(5):1465-1473.
- 695 16. Stronach B, Cronin J, Portus M. Part 1: Biomechanics,
696 injury surveillance, and predictors of injury for cricket fast
697 bowlers. *Strength Cond J.* 2014;36(4):65-72.
- 698 17. Stronach B, Cronin J, Portus M. Part 2: Mechanical and
699 anthropometric factors of fast bowling for cricket, and
700 implications for strength and conditioning. *Strength Cond*
701 *J.* 2014;36(5):53-60.
- 702 18. Stretch R, Bartlett R, Davids K. A review of batting in
703 men's cricket. *J Sports Sci.* 2000;18(12):931-949.
- 704 19. Penn M, Spratford W. Are current coaching
705 recommendations for cricket batting technique supported
706 by biomechanical research? *Sports Biomech.*
707 2012;11(3):311-323.
- 708 20. Law M, Stewart D, Pollock N, Letts L, Bosch J,
709 Westmorland M. *Critical review form - quantitative*
710 *studies.* Hamilton: MacMaster University; 1998.
- 711 21. Cashman R. Crisis in contemporary cricket. In: Cashman
712 R, Mckernan M, eds. *Sport: Money, Morality and the*

- 713 *Media*. Kensington, NSW: New South Wales University
714 Press Limited; 1981:305-312.
- 715 22. Duffield R, Drinkwater. Time-motion analysis of Test and
716 One-Day international cricket centuries. *J Sports Sci*.
717 2008;26(5):457-464.
- 718 23. Portus M, Farrow D. Enhancing cricket batting skill:
719 Implications for biomechanics and skill acquisition
720 research and practice. *Sports Biomech*. 2011;10(4):294-
721 305.
- 722 24. Scanlan A, Wen N, Tucker P, Dalbo V. The relationships
723 between internal and external training load models during
724 basketball training. *J Strength Cond Res*. 2014;28(9):2397-
725 2405.
- 726 25. Fletcher A. Calories and cricket. *Lancet*.
727 1955;268(6875):1165-1166.
- 728 26. Bot S, Hollander A. The relationship between heart rate
729 and oxygen uptake during non-steady state exercise.
730 *Ergonomics*. 2000;43(10):1578-1592.
- 731 27. Alexandre D, da Silva C, Hill-Haas S, del Wong P, Natali
732 A, De Lima J, Bara Filho M, Marins J, Garcia E, Karim C.
733 Heart rate monitoring in soccer: Interest and limits during
734 competitive match play and training, practical application.
735 *J Strength Cond Res*. 2012;26(10):2890-2906.
- 736 28. Vickery W, Dascombe B, Duffield R. Physiological,
737 movement and technical demands of centre-wicket
738 Battlezone, traditional net-based training and one-day
739 cricket matches: A comparative study of sub-elite cricket
740 players. *J Sports Sci*. 2014;32(8):722-737.
- 741 29. Balady G, Chaitman B, Driscoll D, Foster C, Froelicher E,
742 Gordon N, Pate R, Rippe J, Bazzarre T. AHA/ACSM joint
743 position statement: Recommendations for cardiovascular
744 screening, staffing, and emergency policies at
745 health/fitness facilities. *Med Sci Sports Exerc*.
746 1998;30(6):1009-1018.
- 747 30. Noakes T, Durandt J. Physiological requirements of
748 cricket. *J Sports Sci*. 2000;18(12):919-929.
- 749 31. Dupont G, Blondel N, Berthoin S. Performance for short
750 intermittent runs: active recovery vs. passive recovery. *Eur*
751 *J Appl Physiol*. 2003;89(6):548-554.
- 752 32. Houghton L, Dawson B, Rubenson J. Performance in a
753 simulated cricket batting innings (BATEX): Reliability and
754 discrimination between playing standards. *J Sports Sci*.
755 2011;29(10):1097-1103.
- 756 33. Vickery W, Dascombe B, Duffield R, Kellett A, Portus M.
757 Battlezone: An examination of the physiological responses,
758 movement demands and reproducibility of small-sided
759 cricket games. *J Sports Sci*. 2012;31(1):77-86.
- 760 34. Matthew D, Delextrat A. Heart rate, blood lactate
761 concentration, and time-motion analysis of female

- 762 basketball players during competition. *J Sports Sci.*
763 2009;27(8):813-821.
- 764 35. Garcia-Tabar I, Llodio I, Sánchez-Medina L, Ruesta M,
765 Ibañez J, Gorostiaga E. Heart rate-based prediction of
766 fixed blood lactate thresholds in professional team-sport
767 players. *J Strength Cond Res.* 2015;29(10):2794-2801.
- 768 36. Austin D, Kelly S. Positional differences in professional
769 rugby league match play through the use of global
770 positioning systems. *J Strength Cond Res.* 2013;27(1):14-
771 19.
- 772 37. Scanlan A, Dascombe B, Reaburn P, Dalbo V. The
773 physiological and activity demands experienced by
774 Australian female basketball players during competition. *J*
775 *Sci Med Sport.* 2012;15(4):341-347.
- 776 38. Coutts A, Rampinini E, Marcora S, Castagna C,
777 Impellizzeri F. Heart rate and blood lactate correlates of
778 perceived exertion during small-sided soccer games. *J Sci*
779 *Med Sport.* 2009;12(1):79-84.
- 780 39. Scanlan A, Wen N, Tucker P, Borges N, Dalbo V. Training
781 mode's influence on the relationships between training-
782 load models during basketball conditioning. *Int J Sports*
783 *Physiol Perform.* 2014;9(5):853-859.
- 784 40. Petersen C, Pyne D, Portus M, Dawson B. Validity and
785 reliability of GPS units to monitor cricket-specific
786 movement patterns. *Int J Sports Physiol Perform.*
787 2009;4(3):381-393.
- 788 41. Vickery W, Dascombe B, Baker J, Higham D, Spratford
789 W, Duffield R. Accuracy and reliability of GPS devices for
790 measurement of sports-specific movement patterns related
791 to cricket, tennis, and field-based team sports. *J Strength*
792 *Cond Res.* 2014;28(6):1697-1705.
- 793 42. Petersen C, Portus D, Dawson M. Quantifying positional
794 movement patterns in Twenty20 cricket. *Int J Perform*
795 *Anal Sport.* 2009;9(2):165-170.
- 796 43. Padulo J, Tabben M, Ardigo L, Ionel M, Popa C, Gevat C,
797 Zagatto A, Dello Iacono A. Repeated sprint ability related
798 to recovery time in young soccer players. *Res Sports Med.*
799 2015;23(4):412-423.
- 800 44. Stretch R, Buys F, Toit E, Viljoen G. Kinematics and
801 kinetics of the drive off the front foot in cricket batting. *J*
802 *Sports Sci.* 1998;16(8):711-720.
- 803 45. Stuelcken M, Portus M, Mason B. Off-side front foot
804 drives in men's high performance cricket. *Sports Biomech.*
805 2005;4(1):17-35.
- 806 46. Taliep M, Galal U, Vaughan C. The position of the head
807 and centre of mass during the front foot off-drive in skilled
808 and less-skilled cricket batsmen. *Sports Biomech.*
809 2007;6(3):345-360.

810 **TABLE LEGEND**

811

812 **Table 1.** Quality evaluation for each retrieved publication.

813

814 **Table 2.** Summary of studies reporting on the internal
815 physiological responses to batting in cricket during
816 games and training.

817

818 **Table 3.** Summary of studies reporting on the external
819 physiological responses to batting in cricket during
820 games and training.

Table 1. Quality evaluation for each retrieved publication.

Study	Purpose	Literature	Participants	Reliability	Validity	Results	Analysis	Clinical	Conclusions	Practical	Study limitations	Score
Christie et al. ³	Yes	No	Yes	No	No	Yes	Yes	No	Yes	No	Simulated batting bout used. Single live bowler used.	5
Duffield and Drinkwater ²²	Yes	Yes	No	Yes	No	Yes	Yes	Yes	Yes	Yes	Coding approach likely lowered mean durations of high-intensity bouts.	7
Houghton et al. ³²	Yes	No	Yes	Yes	No	Yes	Yes	Yes	Yes	No	Simulated batting bout used. Bowling machine used.	7
Houghton et al. ⁵	Yes	Yes	Yes	No	No	Yes	Yes	Yes	Yes	Yes	Simulated batting bout used. Single live bowler used.	8
Petersen et al. ⁴²	Yes	Yes	No	Yes	Yes	No	Yes	Yes	Yes	Yes	Data were scaled to a 30-min innings. Positional analyses conducted limiting focus on batsmen.	8
Petersen et al. ⁸	Yes	Yes	Yes	Yes	Yes	No	Yes	Yes	Yes	Yes	Only one 3-day game analyzed for multi-day data. Positional analyses conducted limiting focus on batsmen.	9
Pote and Christie ⁶	Yes	Yes	Yes	No	No	Yes	Yes	Yes	Yes	No	Simulated batting bout used. Bowling machine used.	7
Vickery et al. ²⁸	Yes	Yes	Yes	No	No	No	Yes	Yes	Yes	Yes	Small-sided games training bout used. Positional analyses conducted limiting focus on batsmen.	7
Vickery et al. ³³	Yes	Yes	Yes	No	No	Yes	Yes	No	Yes	Yes	Small-sided games training bout used. Positional analyses conducted limiting focus on batsmen.	7

Note: Quality evaluation adapted from Law et al.²⁰ Purpose = was the purpose stated clearly; Literature = was relevant background literature reviewed; Participants = was the sample described in detail; Reliability = were the outcome measures reliable; Validity = were the outcome measures valid; Results = were the results reported in terms of statistical significance; Analysis = were the analysis method(s) appropriate; Clinical = was clinical importance reported; Conclusions = were the conclusions appropriate given study methods and results; Practical = were the implications of the results to practice provided. Total score is summed across each item where: Yes = 1; No = 0.

Table 2. Summary of studies reporting on the internal physiological responses to batting in cricket.

Study	Participants	Playing level	Format/protocol	Mean metabolic responses	Mean HR (b·min ⁻¹)	[BLa ⁻] (mmol·L ⁻¹)	RPE (AU)
Christie et al. ³	n = 10 22 ± 3 years	University (first team)	Simulated batting bout (7 overs)	Energy: 2,536 ± 302 kJ·h ⁻¹ V _E : 65.1 ± 7.9 L·min ⁻¹ VO ₂ : 26.7 ± 1.4 mL·kg ⁻¹ ·min ⁻¹ RER: 1.05 ± 0.05	145 ± 11	–	–
Houghton et al. ³²	n = 11 21 ± 2 years	Club (first and second teams)	BATEX	–	137 ± 11	3.2-4.5 across stages	10-17 across stages (6-20 scale)
	n = 11 19 ± 1 years	Club (third and fourth teams)	BATEX	–	143 ± 14	3.0-4.1 across stages	10-17 across stages (6-20 scale)
Houghton et al. ⁵	n = 9 20 ± 3 years	Club (first to fourth teams)	BATEX	–	130 ± 16	–	13 ± 3 (6-20 scale)
Petersen et al. ⁸	n = 16 22 ± 3 years	Second-tier international	Twenty20	–	149 ± 17	–	–
	n = 5		One Day	–	144 ± 13	–	–
Pote and Christie ⁶	n = 17 23 ± 2 years	Club (first and second teams)	BATEX	–	144 ± 15	–	12 ± 3 9-16 across stages (6-20 scale)
Vickery et al. ²⁸	n = 11 22 ± 3 years	Club (first team)	Battlezone	–	164 ± 12	3.2 ± 1.4	5 ± 2 (1-10 scale)
	n = 10	Club (first team)	One Day	–	159 ± 12	–	5 ± 2 (1-10 scale)
Vickery et al. ³³	n = 13 23 ± 4 years	Club (first and second teams)	Battlezone	–	–	1.8 ± 0.7	4 ± 1 (1-10 scale)

Note: HR = heart rate; RPE = rating of perceived exertion presented as mean or range; AU = arbitrary units; [BLa⁻] = blood lactate concentration presented as mean or range; BATEX = batting exercise simulation protocol consisting of 6 x 21-min stages typical of a One Day International-level score of 100 runs; Battlezone = small-sided cricket game-play consisting of 6-8-over bouts; Energy = energy expenditure; V_E = minute ventilation; VO₂ = oxygen uptake; RER = respiratory exchange ratio.

Table 3. Summary of studies reporting on the external physiological responses to batting in cricket.

Study	Format/protocol	Standing	Walking	Jogging	Running	Striding	Sprinting	Total
<i>Activity frequencies</i>								
Duffield and Drinkwater ²²	One Day							
	50-run innings	190 ± 40	216 ± 49	86 ± 34	–	50 ± 18	22 ± 10	–
	80-run innings	285 ± 53	332 ± 69	143 ± 49	–	81 ± 31	34 ± 14	–
	100-run innings	315 ± 70	367 ± 93	156 ± 43	–	95 ± 44	43 ± 32	–
	Multi-day							
	50-run innings	264 ± 66	267 ± 647	68 ± 17	–	41 ± 11	19 ± 6	–
80-run innings	438 ± 80	438 ± 74	107 ± 13	–	65 ± 9	35 ± 12	–	
100-run innings	527 ± 111	526 ± 96	139 ± 14	–	83 ± 9	39 ± 12	–	
Petersen et al. ^{42#}	Twenty20	–	–	–	–	–	12 ± 5	–
						38 ± 17 (high-intensity)		
Petersen et al. ⁸	Twenty20	–	–	–	–	–	15 ± 9·h ⁻¹	–
						45 ± 16 ·h ⁻¹ (high-intensity)		
	One Day	–	–	–	–	–	13 ± 9·h ⁻¹	–
						39 ± 16 ·h ⁻¹ (high-intensity)		
	Multi-day	–	–	–	–	–	8 ± 3·h ⁻¹	–
						28 ± 6 ·h ⁻¹ (high-intensity)		
Vickery et al. ²⁸	The Battlezone	–	–	–	–	–	23 ± 19·h ⁻¹	–
						224 ± 73 ·h ⁻¹ (high-intensity)		
	One Day	–	–	–	–	–	8 ± 8·h ⁻¹	–
						50 ± 21 ·h ⁻¹ (high-intensity)		
Vickery et al. ^{33*}	The Battlezone	–	–	–	–	–	3 ± 3	–
						39 ± 20 (high-intensity)		
<i>Activity durations</i>								
Duffield and Drinkwater ²²	One Day							
	50-run innings	50.8 ± 11.5 min	29.3+6.6 min	3.0 ± 1.3 min	–	1.4 ± 0.5 min	0.8 ± 0.3 min	–
	80-run innings	74.5 ± 13.7 min	41.4 ± 7.1 min	5.0 ± 1.7 min	–	2.3 ± 0.8 min	1.0 ± 0.5 min	–
	100-run innings	79.1 ± 12.1 min	45.5 ± 9.3 min	5.1 ± 1.3 min	–	2.6 ± 1.1 min	1.2 ± 0.9 min	–

	Multi-day							
	50-run innings	68.6 ± 20.3 min	35.1 ± 8.2 min	2.6 ± 0.8 min	–	1.1 ± 0.3 min	0.6 ± 0.2 min	–
	80-run innings	113.9 ± 22.0 min	55.6 ± 11.8 min	3.9 ± 0.8 min	–	1.7 ± 0.3 min	1.1 ± 0.4 min	–
	100-run innings	133.2 ± 29.5 min	65.1 ± 13.0 min	5.4 ± 1.0 min	–	2.3 ± 0.4 min	1.3 ± 0.5 min	–
Petersen et al. ^{42#}	Twenty20	–	28.43 ± 0.78 min (low-intensity)		1.35 ± 0.72 min (high-intensity)			–
<i>Activity distances</i>								
Houghton et al. ⁵	BATEX	–	1,359 ± 157 m·h ⁻¹	233 ± 33 m·h ⁻¹	99 ± 10 m·h ⁻¹	217 ± 31 m·h ⁻¹	261 ± 58 m·h ⁻¹	2,171 ± 157 m·h ⁻¹
Petersen et al. ^{42#}	Twenty20	–	1,644 ± 507 m	395 ± 114 m	80 ± 34 m	153 ± 91 m	161 ± 83 m	2,433 ± 450 m
Petersen et al. ⁸	Twenty20	–	1,638 ± 352 m·h ⁻¹	332 ± 103 m·h ⁻¹	97 ± 35 m·h ⁻¹	187 ± 70 m·h ⁻¹	175 ± 97 m·h ⁻¹	2,429 ± 606 m·h ⁻¹
	One Day		1,808 ± 400 m·h ⁻¹	279 ± 119 m·h ⁻¹	86 ± 37 m·h ⁻¹	154 ± 70 m·h ⁻¹	149 ± 94 m·h ⁻¹	2,476 ± 631 m·h ⁻¹
	Multi-day		1,604 ± 438 m·h ⁻¹	200 ± 90 m·h ⁻¹	67 ± 18 m·h ⁻¹	107 ± 33 m·h ⁻¹	86 ± 28 m·h ⁻¹	2,064 ± 630 m·h ⁻¹
Vickery et al. ²⁸	The Battlezone	–	2619 ± 1173 m·h ⁻¹ (low-intensity)		1235 ± 422 m·h ⁻¹ (high-intensity)			3,895 ± 1,236 m·h ⁻¹
	One Day		1632 ± 794 m·h ⁻¹ (low-intensity)		271 ± 12 m·h ⁻¹ (high-intensity)			1,919 ± 793 m·h ⁻¹
Vickery et al. ^{33*}	The Battlezone	–	566 ± 55 m	351 ± 46 m	104 ± 31 m	99 ± 67 m	21 ± 27 m	1,147 ± 175 m

Note: Activity intensities typically calculated as walking = ≤ 2 m·s⁻¹, jogging = 2.01-3.5 m·s⁻¹, running = 3.51-4 m·s⁻¹, striding = 4.01-5 m·s⁻¹, sprinting = >5 m·s⁻¹; # indicates data scaled to a 30 min inning; * indicates data collected across mean bout length of 18 ± 2 min; BATEX = batting exercise simulation protocol consisting of 6 x 21-min stages typical of a One Day International-level score of 100 runs; Battlezone = small-sided cricket game-play consisting of 6 bouts of 8-overs.