

Does acute exercise affect the performance of whole-body, psychomotor skills in an inverted-U fashion? A meta-analytic investigation.

Terry McMorris^{abc*}, Beverley J. Hale^a, Jo Corbett^c, Kevin Robertson^b, Christopher I. Hodgson^d

^a Department of Sport and Exercise Science, University of Chichester, College Lane, Chichester, West Sussex PO19 6PE, United Kingdom

^b Department of Psychology, Faculty of Health and Life Sciences, Northumbria University, Northumberland Road, Newcastle-upon-Tyne NE1 8ST, United Kingdom

^c Department of Sport and Exercise Science, Faculty of Science, University of Portsmouth, Guildhall Walk, Portsmouth PO1 2BF, United Kingdom

^d Department of Adventure Education, University of Chichester, College Lane, Chichester, West Sussex PO19 6PE, United Kingdom

* Address correspondence to: Terry McMorris, 63 Four Winds Court, Hartlepool TS26 0LP, United Kingdom. e-mail t.mcmorris@chi.ac.uk Tel. 44 1429 235894

Abstract

The primary purpose of this study was to examine, using meta-analytical measures, whether research into the performance of whole-body, psychomotor tasks following moderate and heavy exercise demonstrates an inverted-U effect. A secondary purpose was to compare the effects of acute exercise on tasks requiring static maintenance of posture versus dynamic, ballistic skills. Moderate intensity exercise was determined as being between 40% and 79% maximum power output (\dot{W}_{MAX}) or equivalent, while $\geq 80\% \dot{W}_{MAX}$ was considered to be heavy. There was a significant difference ($Z_{diff} = 4.29$, $p = 0.001$, $R^2 = 0.42$) between the mean effect size for moderate intensity exercise ($g = 0.15$) and that for heavy exercise size ($g = -0.86$). These data suggest a catastrophe effect during heavy exercise. Mean effect size for static tasks ($g = -1.24$) was significantly different ($Z_{diff} = 3.24$, $p = 0.001$, $R^2 = 0.90$) to those for dynamic/ballistic tasks ($g = -0.30$). The result for the static versus dynamic tasks moderating variables point to perception being more of an issue than peripheral fatigue for maintenance of static posture. The difference between this result and those found in meta-analyses examining the effects of acute exercise on cognition show that, when perception and action are combined, the complexity of the interaction induces different effects to when cognition is detached from motor performance.

Keywords: arousal; fatigue; perception; action; catecholamines; prefrontal cortex

1. Introduction

Yerkes and Dodson's [1] inverted-U theory, concerning the effect of arousal on performance, and theories developed from this early work [2-4] have generally been used as the rationales for studies examining the effect of acute exercise on cognition (e. g., [5-7]). It has normally been hypothesized that moderate intensity exercise equates to moderate levels of arousal and so optimal performance is expected, while heavy exercise equates to over-arousal and so performance returns to the same level as that at rest. A recent meta-analysis [8] supported this hypothesis. The primary purpose of the present study was to examine, using meta-analytical measures, whether research into the performance of whole-body, psychomotor tasks following moderate and heavy exercise also demonstrates an inverted-U effect. Meta-analytic methods were undertaken as they facilitate the use of a larger sample size than one normally finds in research on this topic. Moreover, the emphasis on effect sizes rather than probability allows for a better evaluation of those studies where failure to show a significant effect was due to sample size possibly resulting in Type II errors.

Whole-body, psychomotor skills require integrated control by the Central and Peripheral Nervous Systems (PNS). The decision to act is made by the higher centers of the brain, particularly the prefrontal cortex, and action is initiated by the premotor cortex and/or supplementary motor area, with the former being primarily concerned with movement in response to external events while the latter mainly controls voluntary movement, although both are active during any type of movement. The information is passed downwards to the PNS via several Central Nervous System (CNS) regions including the basal ganglia, brainstem, cerebellum and spinal cord. Information from the CNS is transmitted by efferent nerves to motor units in the musculature. These neurons activate the musculature. Once the action begins information from the PNS, about the movement, is feedback to the brain by

afferent neurons, situated in the muscles, joints and spinal cord. The spinal cord itself can make very fast (~ 30 ms), but very limited alterations to the movement, using the process of α - γ coactivation [9]. Feedback to the cerebellum, the so-called long loop feedback, is greater in scope than the α - γ coactivation process but takes ~ 80 ms to be activated [10]. The most important feedback is to the sensory regions of the brain, particularly the visual and somatosensory cortices, and, in some skills, the auditory cortex. The prefrontal cortex and the sensory association areas receive information from the sensory cortices and organize and interpret this information. These higher centers of the brain can initiate large alterations to the movement but take time (> 400 ms) [11]. This CNS-PNS interaction ensures that the movements are coordinated and smooth, and that motor unit recruitment allows for the production of the required power.

The processes outlined above can be affected by a number of stressors, including acute exercise. The most obvious effect of acute exercise on the performance of such skills is physiological, although the precise nature of these physiological effects will vary primarily as a function of the intensity (e.g. moderate vs. heavy) but may also be influenced by the duration; the environment under which the exercise is conducted; the fitness level of the exerciser and the elapsed time between the exercise and performance of the criterion task (see [12] for a review). Broadly speaking, relevant physiological effects could incorporate central processes related to alterations in the intrinsic motoneuron properties, sensory feedback, or descending drive [13] and peripheral processes occurring distal to the neuromuscular junction, including those related to sarcoplasmic reticulum calcium release and decreased myofibrillar force production [14]. More precisely, authors have previously emphasized the role of physiological factors related to the level of nervous system activation [6, 15]; the efficiency of the peripheral motor processes (i.e. better synchronisation of the motor units discharge); peripheral sensorial processes [16-17]; the involvement of different metabolic

systems [18] and associated effects related to metabolic acidosis, or the accumulation of metabolic waste products and humoral changes [19].

The physiological changes induced by acute exercise are accompanied by biochemical changes peripherally and centrally. During and even immediately before exercise, the hypothalamus and brainstem initiate action of the sympathoadrenal system. This results in the release of catecholamines at the postganglionic cells of those neurons that require activating or inhibiting. If exercise increases in intensity to a moderate level, there is also release of epinephrine and, to a lesser extent norepinephrine, into the blood from the adrenal medulla. As exercise intensity increases further to a level which we could describe as heavy, there are larger increases in plasma norepinephrine and epinephrine concentrations. Peripherally, norepinephrine and epinephrine aid lipolysis, stimulate receptors in muscle and activate receptors in the pancreas to suppress insulin release. Epinephrine plays a major role in glycogenolysis and control of the cardiovascular system by activating receptors responsible for increasing heart rate and contractile force (see [20] for a review).

Although catecholamines do not readily cross the blood brain barrier, rodent studies (see [21] for a review) have demonstrated significant increases in brain concentrations of dopamine and norepinephrine following acute exercise. This is most likely due to the fact that peripherally circulating epinephrine and norepinephrine activate β -adrenoreceptors on the afferent vagus nerve, which runs from the abdomen through the chest, neck and head, and terminates in the nucleus tractus solitarii (NTS) within the blood-brain barrier. Noradrenergic cells in the NTS, which project into the locus coeruleus, stimulate norepinephrine synthesis and release to other parts of the brain [22-23]. This may also affect brain dopamine concentrations, as Devoto et al. [24] showed that electrical stimulation of the rat locus coeruleus resulted in increased brain concentrations of dopamine and one of its metabolites, 3,4-dihydroxyphenylacetic acid. During moderate intensity exercise, there are moderate

increases in concentrations of brain catecholamines, which activate the prefrontal cortex, sensory cortices and their association areas. These increases lead to improved sensation and perception by increasing the signal to 'noise' ratio within the brain. Heavy exercise, however, leads to even greater increases in brain concentrations of catecholamines, which disrupts the signal to 'noise' ratio, hence inhibiting sensation and perception [25-26].

Catecholamines are not the only neurochemicals, activated during exercise, which may affect sensation and perception. There are several but the hypothalamic-pituitary-adrenal cortex (HPA) axis hormones are probably the most important. Peripherally, during exercise, the HPA axis hormone cortisol plays major roles in glucose production from proteins, the facilitation of fat metabolism and muscle function, and the maintenance of blood pressure [27]. However, this appears to only occur when exercise is heavy [28]. Cortisol readily crosses the blood brain barrier, so peripheral increases in concentrations will lead to central increases. Moreover, the synthesis and release of cortisol by the HPA axis is initiated by the synthesis and release of the protein corticotrophin releasing factor (CRF) and the hormone adrenocorticotrophin hormone (ACTH) [29]. Given that CRF is released in the brain and ACTH in the anterior pituitary, which lies within the CNS, it is not surprising to find that rodent studies have demonstrated acute exercise-induced increases in brain concentrations of the HPA axis neurochemicals [30-32]. These neurochemicals interact with catecholamines in the brain, resulting in increased synthesis and release of dopamine and norepinephrine, which should inhibit sensation and perception during heavy exercise [33-35].

The theory outlined above suggests that moderate, acute exercise will facilitate the performance of whole-body, psychomotor skills, due to moderate increases in concentrations of the brain neurotransmitters dopamine and norepinephrine inducing improved perceptual performance. Moreover, physiological mechanisms that could positively influence motor processes through mechanisms, including effects on nervous system activation, motor unit

coordination [13] or improved contractile function [14], will also be facilitated. During heavy exercise, changes related to metabolic acidosis and fatigue may elicit different physiological changes which have the potential to impair the motor process at a variety of central and peripheral sites, although the influence of acidosis on contractile function is controversial [36]. Also excessive brain concentrations of catecholamines should inhibit sensation and perception, meaning that we can hypothesize that there will be a significant difference between effect sizes during moderate and heavy exercise. In humans, when the stressor is psychological and the task cognitive, one tends to find that these changes in brain concentrations of catecholamines result in an inverted-U effect. However, we assert that it is possible that, when the task is physical, a combination of the central and peripheral changes might result in heavy exercise inducing poorer performance than at rest, thus demonstrating an inverted-J effect.

A secondary purpose of this study was to compare the effects of acute exercise on tasks requiring static maintenance of posture (e. g. static balance and shooting) and dynamic, ballistic skills (e. g. most sports skills). While both require the integration of the CNS and PNS, the nature of the movements and the integration of perceptual information differ. Maintenance of posture has been shown to be heavily dependent on central perception of balance, which appears to be negatively affected by exercise [37-39], while the perceptual and decision making aspects of many dynamic skills have been shown not to be affected even by heavy exercise and indeed in some cases are facilitated [7, 40]. However, the physiological demands of dynamic skills may result in deterioration in performance of such skills.

2. Materials and methods

A literature search using the computer data bases PsycArticles, PsycINFO, Pubmed, SPORTSDiscus and Web of Knowledge was undertaken. Key words used in the searches were combinations of “acute”, “exercise”, *psychomotor performance*”, “*psychomotor skills*”, “*physical activity*”, “*fatigue*” and the actual names of a large variety of whole-body, psychomotor skills. In addition, reference lists from empirical reports and reviews were examined. Studies were included if they were performed on healthy individuals and repeated measures, within-subject designs were used. In studies using pharmacological or nutritional treatments, the control or placebo groups’ data were included in the meta-analyses but not the experimental groups.

2.1. Definitions of moderate and heavy exercise.

When exercise was aerobic, Borer’s [20] classifications of moderate and heavy exercise formed the basis of our definitions, the same as those used by McMorris and colleagues [8, 41]. Moderate intensity exercise was determined as being between 40% and 79% maximum power output (\dot{W}_{MAX}) or equivalent, while $\geq 80\% \dot{W}_{MAX}$ was considered to be heavy. If \dot{W}_{MAX} values were not presented but percent volume of maximum oxygen uptake ($\dot{V}O_{2MAX}$) or percent maximum heart rate were given, the conversion formulae of Arts and Kuipers [42] were applied. For other indicators of intensity, e.g. percent heart rate reserve, percent maximum aerobic power, percent ventilatory threshold and percent lactate threshold power, the exercise physiology and exercise endocrinology literatures were examined to ascertain whether or not the intensity would be below, within or above the 40–79% \dot{W}_{MAX} limits. In those studies using isotonic or isometric exercise, contractions $\geq 80\%$ of the participants’ maximal number of contractions were considered to be heavy, while contractions $< 80\%$ but $> 40\%$ maximum were deemed moderate. Where exercise was intermittent anaerobic and aerobic, duration and time working at each of the intensities were

used to determine whether or not the overall intensity could be classed as moderate or heavy. Where exercise was to voluntary exhaustion or until the individual could not maintain the required intensity, it was deemed heavy.

2.3. Data analyses

Where means and SDs were available, effect sizes were calculated using the Cohen's d formula (Mean at rest–Mean during or following exercise/SD at rest, where rest acts as the control). Studies in which means and SDs were only provided graphically were not included as it was not possible to accurately determine the means and especially the SDs. Each individual d was then transformed to the bias-corrected standardized mean difference, Hedges' g , by applying the correction factor J $\{(J=1- [3/(4df-1)])\}$ [43] and this was used to calculate a mean effect size using the random-effects model. Results of the Q test for homogeneity were calculated and reported as was τ^2 , which is a measure of absolute variance whereas Q is a measure of normalized variance [43]. Orwin's [44] Fail-safe N was calculated when the mean effect size g was ≥ 0.20 . Where sub-group analyses were undertaken, effect sizes for each group were compared using a Z -test on the differences with a random-effects model, with separate estimates of τ^2 for each sub-group. The proportion of variance explained by the moderator variable, R^2 , was calculated [43]. Most studies provided more than one effect size. In order to control for one or more studies having an undue bias on the results, one effect size per intensity per study was calculated, with one exception ([45] see section 3). The data were analyzed using the computer package Comprehensive Meta Analysis version 2.0 [see 43].

3. Results

The literature reviewed showed that there were 89 articles which examined the effect of acute exercise on the performance of whole-body, psychomotor skills but only 28 which met the criteria for inclusion and provided sufficient statistical information. However, one study [45] provided data for novice and expert performers separately and so was treated as two separate studies. This meant that there were 23 studies in which effect sizes were calculated for one exercise intensity only and six where two intensities were included. In total, there were 35 effect sizes and 570 participants. The types of tasks and exercise intensities used in each study can be seen in Table 1.

Insert Table 1 about here

An initial overall analysis, including both the moderate and heavy exercise dependent variables, is necessary before comparisons can be undertaken [43]. This showed that effect sizes were heterogeneous $Q(34) = 155.68$ ($p < 0.001$), $\tau^2 = 0.46$. The mean effect size was significant, $g = -0.55$ ($Z = 4.14$, $p < 0.001$), variance 0.018, $SE = 0.13$, and 95% confidence interval (CI) -0.81 to -0.29. The fail-safe N was 61. Twenty-six effect sizes were negative and nine positive. Sub-group analyses for moderate ($k = 11$) and heavy ($k = 24$) exercise showed a significant difference between the two variables ($Mean_{diff} = 1.03$, $SE = 0.24$, $Z_{diff} = 4.29$, $p = 0.001$, $R^2 = 0.42$). Mean effect size for moderate intensity exercise was non-significant ($g = 0.15$, $SE = 0.12$), while heavy exercise demonstrated a significant mean effect size ($g = -0.86$, $Z = 5.85$, $p < 0.001$, variance = 0.02, $SE = 0.15$, CI = -1.14 to -0.57). Heterogeneity for moderate exercise was non-significant [$Q(10) = 14.83$ ($p > 0.05$), $\tau^2 = 0.05$] but significant for heavy exercise [$Q(23) = 78.97$ ($p < 0.001$), $\tau^2 = 0.35$].

Sub-group analyses for static and dynamic skills demonstrated a significant difference between the two variables ($Mean_{diff} = 0.94$, $SE = 0.29$, $Z_{diff} = 3.24$, $p = 0.001$, $R^2 = 0.90$). Mean effect size for static tasks ($k = 10$) was significant ($g = -1.24$, $Z = 4.84$, $p < 0.001$, variance = 0.07, $SE = 0.26$, CI = -1.75 to -0.74), as was the mean effect size for dynamic ($k =$

27) skills ($g = -0.30$, $Z = 2.33$, $p < 0.02$, variance = 0.02, SE = 0.13, CI = -0.55 to -0.05).

Both conditions demonstrated significant heterogeneity, $Q(9) = 75.91$ ($p < 0.001$), $\tau^2 = 0.44$ for static balance and $Q(26) = 79.61$ ($p < 0.001$), $\tau^2 = 0.27$ for ballistic/dynamic.

It was decided, a posteriori, to examine the effect of using counterbalanced/randomized designs compared to a pre-exercise/post-exercise design. Sub-group analyses showed no significant differences. Mean effect size for counterbalanced/randomized designs was $g = -0.42$ (SE = 0.19, $Z = 2.20$, $p < 0.03$) and for pre- followed by post-exercise $g = -0.62$ (SE = 0.18, $Z = 3.51$, $p = 0.01$).

4. Discussion

The overall analysis shows a moderate to high effect size. That it was negative is not too surprising, given that there were more studies measuring the effects of heavy exercise than moderate. Nevertheless, a regression towards zero was expected, as we thought that moderate exercise would induce positive effect sizes while heavy would result in negative effect sizes. This was not demonstrated as 45.45% of the of the moderate intensity results showed negative effects. This result is very different to those found in meta-analyses examining the effect of acute exercise on cognitive skills, including perception, when the skill is carried out either during exercise or immediately following cessation of the exercise. In those studies, the overall analyses with both moderate and heavy exercise included, have tended to show small to moderate, but significant effects, mostly positive [8, 46-48] but one negative [49].

4.1. Moderate intensity exercise effects

The effect size for moderate exercise was positive but small and non-significant, moreover effect sizes were somewhat homogeneous and the CI was relatively small. Taken together, these data suggest that there was very little effect of moderate exercise on performance of the skills. We expected a high effect size ($g \geq 0.70$) because of the strong theoretical rationale. One would expect the increase in body temperature during moderate intensity exercise to induce increased speed of nerve transmission [50], which would aid coordination and power production. Moreover, when brain catecholamines concentrations are increased to a moderate level, there is increased firing of α_{2A} -adrenoreceptors by norepinephrine [51], which increases the strength of the neural signal, and D1 dopaminergic receptors by dopamine [52], which dampens the 'noise' by inhibiting firing to non-preferred stimuli, thus improving the signal to 'noise' ratio [25-27]. This should strongly improve perceptual performance by optimizing activity in the reticular formation, which controls attention, alertness and vigilance, and the prefrontal cortex, which is responsible for integration and interpretation of information from the sensory cortices and their association areas.

Before attempting to look at possible physiological and neuroscientific reasons for these results, we must examine some possible methodological issues. As the sample size ($k = 11$) was small from the point of view of number of studies, the possibility of a lack of power resulting in a Type II error has to be taken into account. According to Clarke-Carter [53], to attain a power of 0.80, with $k = 11$, we would need to elicit an effect size of $g = 0.75$, very close to our expected $g = 0.70$. This may account for the failure of $g = 0.15$ to reach significance but does not explain why we failed to show the effect size we expected or at least one near to it. The possibility that study designs failed to properly control exercise intensity also needs to be addressed. All exercise intensities classified as moderate met the criteria set out in 2.1. The only questionable issue might be the time spent exercising in the

McRae et al. [54] study (2 hours), which could have resulted in increased brain cortisol concentrations as well as increased dopamine and norepinephrine concentrations. Moreover, three studies [54-56] failed to take into account individual differences in fitness. Given the mean effect sizes for these studies, it does not appear that this has been a contributor, certainly not a major contributor, to our results.

However, the failure of all but two studies [57-58] to take into account individuals' lactate and catecholamines thresholds may have affected results. When we plot plasma concentrations of epinephrine and norepinephrine against exercise intensity, concentrations rise exponentially [59-60]. Green et al. termed the points at which there is a significant rise in concentrations, the epinephrine threshold and the norepinephrine threshold. Although the two thresholds generally show moderate to high correlations, some individuals do not follow the normal pattern [61]. It would appear that exercise intensity needs to be moderate before the thresholds are reached but there are large inter-individual variations [62]. It is generally thought that, for aerobic exercise, intensity needs to be $\sim 75\% \dot{V}O_{2\text{MAX}}$ [61], which according to Arts and Kuipers [42] equates to $\sim 65\% \dot{W}_{\text{MAX}}$. Moreover, blood lactate concentrations follow a similar exponential profile and the lactate threshold shows moderate to high correlations with the catecholamines thresholds [6, 61, 63]. Chmura, Nazar and Kaciuba-Uścilko [6] argued that it is at or immediately following the catecholamines thresholds that a significant improvement in cognitive function will be induced due to increased brain catecholamines concentrations. This makes sense, as increased concentrations of circulating epinephrine and norepinephrine will activate the β -adrenoreceptors on the vagus nerve, thus initiating the action of the vagus/NTS pathway and increased synthesis and release of norepinephrine in the locus coeruleus. Improved cognitive performance at or following the catecholamines thresholds [6-7] and the lactate threshold [64-65] has been demonstrated. Also, improved cognition has been shown at the ventilatory threshold [66-70], the point at

which ventilatory carbon dioxide shows a greater increase than ventilatory oxygen and which occurs about the same time as the lactate threshold [71].

Participants exercising below their catecholamines thresholds would probably not induce increased brain catecholamines concentrations and hence not show improved performance. However, individuals exercising above their thresholds might synthesize and release too much, thus negatively affecting performance. This might account for the equivocal nature of the results and also points to the need for those wishing to use warm-up exercise to aid whole-body, psychomotor performance to individualize the exercise intensity. Another possible reason for the failure to demonstrate positive and significant results could be that, if exercise is above the individual's threshold, CNS and PNS integration are compromised due to increased blood and muscle lactate concentrations, and changes in the balance between ventilatory carbon dioxide and oxygen having detrimental effects on the motor aspects of the psychomotor task.

These results suggest that the CNS-PNS interaction during the performance of psychomotor skills, following moderate intensity exercise, may be very complex. While the catecholamines thresholds may be ideal for brain activation, especially in the prefrontal cortex, this intensity may have negative effects on the physiological aspects. Indeed it could be that different skills require different intensities to induce optimal performance. A great deal more research is necessary. Moreover, research should include more physiological and biochemical measurements than the research at present in the literature.

4.2. Heavy intensity exercise effects

Results for the effects of heavy exercise were as expected with a high effect size being demonstrated. Although heterogeneity of effect sizes remained high, the CI ranged from very high to moderate to high, negative effects. Neurophysiologically, research has

shown that reduced excitations of motoneurons resulting from afferent feedback from muscle spindles [72], reduced impulse frequency to muscle fibers at the neuromuscular junction [73] and failures in the calcium release process [74] significantly affect coordination and power. Metabolically, decreased adenosine triphosphate supply [75], decreased glycogen concentrations [76], decreased pH and increased concentrations of inorganic phosphate [77] all have inhibitory effects on motor control. From a neurochemical perspective, heavy exercise induces very large increases in brain concentrations of catecholamines. This, in turn, leads to the excess norepinephrine activating the lower affinity α 1- and β -adrenoreceptors [51]. α 1-adrenoreceptors can result in reduced neuronal firing in the prefrontal cortex by phosphatidylinositol-protein kinase C intracellular signaling pathway activation. Excessive stimulation of D1 receptors and β -adrenoreceptors can induce excess activity of the secondary messenger cyclic adenosine monophosphate which dampens all neuronal activity, thus weakening the signal to 'noise' ratio in the prefrontal cortex (see [25-26]). During high levels of stress, this is probably exacerbated by stimulation of D2 receptors [25]. Although, stimulation of α 1- and β -adrenoreceptors can improve the signal to 'noise' ratio in the sensory cortices [78-80] and can aid some prefrontal cortex activities, overall it has a negative effect on prefrontal cortex activity [25-26]. This is important because the prefrontal cortex is responsible for the integration and coordination of perceptual information from a variety of sensory regions of the brain [81]. Moreover, as well as integrating and coordinating sensory and perceptual feedback, the prefrontal cortex, particularly the right inferior prefrontal cortex, plays a major role in inhibition of inappropriate motor responses [82], which have a negative effect on the performance of psychomotor skills.

As we saw in section 1, heavy exercise also initiates the release of the HPA axis hormones, which exacerbate the negative effects of the catecholamines. These neurochemicals interact with catecholamines in the brain to affect perception and cognition.

In the locus coeruleus, CRF neurons innervate noradrenergic neurons and norepinephrine is released [83-84]. Similarly there is strong evidence for an interaction between corticosteroid concentrations and dopamine release [85-88]. Thus increased brain concentrations of CRF, ACTH and cortisol during heavy exercise add to the negative effects of catecholamines.

4.3 Effects on static versus dynamic/ballistic psychomotor skills

The results of the sub-groups analyses show that we were correct to expect differences but the proportion of variance was far greater than we had anticipated. We expected only a relatively low coefficient, e. g. $R^2 \simeq 0.40$, as both sets of skills require high levels of perpetual-action coupling. That the effect for the static skills was negative was as expected but that it was so high was surprising. Moreover, although heterogeneity was high, CI results showed that effect sizes ranged from very high to high, negative effects. These skills recruit muscles in the vicinity of the knees, ankles, calves, toes and hips but the range of movement and power required are comparatively small [89-91]. Centrally they require a large input from the dorsolateral prefrontal cortex to integrate information from the visual and somatosensory cortices, cerebellum and vestibular apparatus. Given that the dorsolateral prefrontal cortex is especially susceptible to disruption by excess catecholamines [25-26], inhibition of performance is not surprising. Furthermore, several researchers have argued that exercise affects proprioception more than the motor aspects of balance [37, 39] and it has been shown that the attentional demands of balance actually increase following heavy exercise [37].

Results for ballistic/dynamic skills showed that CI ranged from moderate, negative values to almost zero. The high level of heterogeneity and the large CI suggest the need for more research. The dynamic, ballistic skills also require central integration of perceptual information but even heavy exercise has been shown to have only a limited negative effect on

the perceptual and decision-making aspects of such skills [8, 40, 92]. Moreover, many of the skills utilized in the research covered by this analysis may well have been autonomic to the participants. This is especially so given that many of the skills were sports skills and the participants were often physical education and/or sports science majors. Autonomic skills may well be less negatively affected by stress, even physiological stress. This has been shown for well-learned cognitive skills and implicitly-learned motor skills [92-93]. However, the peripheral physiological adaptations to exercise probably have a negative effect on coordination and power, which results in a small but significant negative effect. If automaticity is a key moderator, it would appear that overlearning could help to lessen the problem of performing following heavy exercise.

4.4. Use of counterbalancing/randomization of testing

Observation of the raw data led us, a posteriori, to examine the use of counterbalancing/randomization of testing as opposed to the use of pre-exercise testing followed by post-exercise testing. Research methods texts recommend counterbalancing or randomization but many of the studies included in the analyses utilized a pre-exercise followed by post-exercise protocol. The possibility of a learning or habituation effect is obvious. Therefore, we decided to compare mean effect sizes for studies using counterbalanced/randomized designs with those using pre-exercise followed by post-exercise testing. That there was no significant difference between studies using counterbalance/randomization and those using a pre-exercise testing followed by post-exercise testing protocol was a little surprising. McMorris and Hale [8], examining the effect of acute exercise on cognition, showed that randomized/counterbalanced designs elicited higher effect sizes than pre-exercise followed by post-exercise testing. They claimed that testing pre-exercise meant that the individual's dopamine and norepinephrine, and possibly

cortisol, concentrations would show an increase pre-exercise due to anticipation of the exercise to come, a phenomenon which has been known for some time [94] and demonstrated recently [95]. This could lead to pre-exercise cognitive performance being better than that at a real baseline. With psychomotor skills one might expect the same pre-exercise increase due to anticipation but there is likely to be a fall in brain catecholamines concentrations once the participants begin the pre-test. Falls in peripheral catecholamines concentrations have been shown when individuals begin to perform a skill compared to pre-performance concentrations [96]. This is probably due to the perception of the stress being greater than the actual stress [94]. This would negate the pre-exercise levels affecting the following rest performance.

5. Conclusion

The results of this study failed to fully support either an inverted-U or an inverted-J effect of acute exercise on the performance of whole-body, psychomotor skills. Moderate intensity exercise demonstrated no significant effect, while heavy exercise showed a negative effect. That moderate intensity exercise failed to induce a significant improvement from rest questions the use of a moderate intensity warm-up for improving performance, a practice that is common particularly in sport. Moreover, the difference between this result and those found in meta-analyses examining the effects of acute exercise on cognition [8, 41, 46-49] show that, when perception and action are combined, the complexity of the interaction induces different effects to when cognition is detached from motor performance. The same appears to be the case with heavy exercise, following which the neurochemical and physiological stress appear to combine to induce a detrimental effect compared to not only moderate intensity exercise but also compared to at rest, baseline measures. The result for the static versus

dynamic tasks moderating variables possibly point to perception being more of an issue than peripheral muscular fatigue for maintenance of static posture.

There are several issues that future research needs to examine. Firstly, a lot more research is required and studies should include physiological and neurochemical measures. Such measures would allow for a better knowledge of the amount of physiological stress placed on the participants. Moreover, individual differences in participants' fitness levels need to be taken into account in the experimental designs. Comparison of tasks involving greater and less prefrontal cortex activation would allow for the investigation of whether the problems were mainly central or peripheral or equally both. Also the limited amount of studies including female participants makes gender comparisons impossible to examine at the moment.

References

*denotes study included in meta-analyses

- [1] Yerkes RM, Dodson JD.. The relation of strength of stimulus to the rapidity of habit formation. *J Comp Neurol Psychol* 1908;18: 459-82.
- [2] Easterbrook JA. The effect of emotion on cue utilization and the organization of behavior. *Psychol Rev* 1959;66: 183-201.
- [3] Kahneman D. *Attention and Effort*. Englewood Cliffs, NJ: Prentice Hall; 1973
- [4] Sanders AF. Towards a model of stress and human performance. *Acta Psychol* 1983 53, 61-97.
- [5] Brisswalter J, Arcelin R, Audiffren M, Delignières D. Influence of physical exercise on simple reaction time: effect of physical fitness. *Percept Mot Skills* 1997;85: 1019-27.

- [6] Chmura J, Nazar H, Kaciuba-Uścilko H. Choice reaction time during graded exercise in relation to blood lactate and plasma catecholamine thresholds. *Int J Sports Med* 1994;15: 172-6.
- [7] McMorris T, Myers S, MacGillivray WW, Sexsmith JR, Fallowfield J, Graydon J, Forster D. Exercise, plasma catecholamine concentration and decision-making performance of soccer players on a soccer-specific test. *J Sports Sci* 1999;17: 667-76.
- [8] McMorris T, Hale BJ. Differential effects of differing intensities of acute exercise on speed and accuracy of cognition: a meta-analytical investigation. *Brain Cognition* 2012;80: 338-51.
- [9] Kuo AD. The relative roles of feedforward and feedback in the control of rhythmic movements. *Motor Control* 2002;6: 129-45.
- [10] Dewhurst DS. Neuromuscular control system. *IEEE T Biomed-Eng* 1967;14: 167-71.
- [11] Brunamonti E, Chiricozzi FR, Clausi S, Olivito G, Giusti MA, Molinari M, Ferraina S, Leggio, M. Cerebellar damage impairs executive control and monitoring of movement generation. *PLoS ONE* 2014;9: e85997. doi:10.1371/journal.pone.0085997
- [12] Åstrand P-O, Rodahl K, Dahl HA, Strømme SB. *Textbook of Work Physiology*, 4th ed. Champaign, IL: Human Kinetics; 2003
- [13] Taylor JL, Gandevia SC. A comparison of central aspects of fatigue in submaximal and maximal voluntary contractions *J Appl Physiol* 2008;104: 542–50
- [14] Allen DG. Skeletal muscle function: role of ionic changes in fatigue, damage and disease. *Clin Exp Pharmacol Physiol*, 2004;31: 485-93.
- [15] McMorris T, Graydon J. The effect of incremental exercise on cognitive performance. *Int J Sport Psychol* 2000;31: 66-81.

- [16] Davranche K, Burle B, Audiffren M, Hasbroucq T. Information processing during physical exercise: A chronometric and electromyographic study. *Exp Brain Res* 2005;165: 532–40.
- [17] Davranche K, Audiffren M, Denjean A. A distributional analysis of physical exercise on a choice reaction time task. *J Sports Sci* 2006;24: 323–9.
- [18] Fleury M, Bard C. Effects of different types of physical activity on the performance of perceptual tasks in peripheral and central vision and coincidence timing. *Ergonomics* 1987;30: 945-58.
- [19] Brisswalter JB, Durand M, Delignieres D, Legros P. Optimal and nonoptimal demand in a dual task of pedaling and simple reaction time: Effects on energy expenditure and cognitive performance. *J Hum Movement Stud* 1995;29: 15–34
- [20] Borer KT. *Exercise Endocrinology*. Champaign, IL: Human Kinetics; 2003
- [21] Meeusen R, Piacentini MF, De Meirleir K. Brain microdialysis in exercise research. *Sports Med* 2001;31: 965-83.
- [22] McGaugh JL, Cahill L, Roozendaal B. Involvement of the amygdala in memory storage: Interaction with other brain systems. *Proc Natl Acad Sci USA*. 1996;93: 13508-14.
- [23] Miyashita T, Williams CL. Epinephrine administration increases neural impulses propagated along the vagus nerve: role of peripheral beta-adrenergic receptors. *Neurobiol. Learn. Mem* 2006;85: 116-24.
- [24] Devoto P, Flore G, Saba P, Fà M, Gessa GL. Co-release of noradrenaline and dopamine in the cerebral cortex elicited by single train and repeated train stimulation of the locus coeruleus. *BMC Neurosci* 2005; doi:10.1186/1471-2202-6-31
- [25] Arnsten AFT. Stress signalling pathways that impair prefrontal cortex structure and function. *Nat Rev Neurosci* 2009;10: 410–22.

- [26] Arnsten AFT. Catecholamine influences on dorsolateral prefrontal cortical networks *Biol Psychiat* 2011;69: e89–e99. doi:10.1016/j.biopsych.2011.01.027
- [27] Deuster PA, Chrousos GP, Luger A, De Bolt JE, Bernier IL, Trostman UH, Kyle SB, Montgomery LC, Loriaux DL. Hormonal and metabolic responses of untrained, moderately trained, and highly trained men to 3 exercise intensities. *Metabolism* 1989;38: 141-8.
- [28] De Vries WR, Bernards NTM, De Rooij MH, Koppeschaar HPF. Dynamic exercise discloses different time-related responses in stress hormones. *Psychosom Med* 2000;62: 866-72.
- [29] Vale W, Rivier C. Substances modulating the secretion of ACTH by cultural anterior pituitary cells. *Fed Proc* 1977;36: 2094-9.
- [30] Droste SK, Gesing A, Ulbricht S, Müller MB, Linthorst ACE, Reul JMHM. Effects of long-term voluntary exercise on the mouse hypothalamic-pituitary-adrenocortical axis. *Endocrinology* 2003;144: 3012-23.
- [31] Ke Z, Yip SP, Li L, Zheng XX, Tong K-Y. The effects of voluntary, involuntary, and forced exercises on brain-derived neurotrophic factor and motor function recovery: a rat brain ischemia model. *PLoS ONE*, 2011; 6: e16643. doi:10.1371/journal.pone.0016643.
- [32] Yanagita S, Amemiya S, Suzuki S, Kita I. Effects of spontaneous and forced running on activation of hypothalamic corticotropin-releasing hormone neurons in rats. *Life Sci* 2007;80: 356-63.
- [33] Asbach S, Schulz C, Lehnert H. Effects of corticotropin-releasing hormone on locus coeruleus neurons in vivo: a microdialysis study using a novel bilateral approach. *Eur J Endocrinol* 2001;145: 359-63.
- [34] Wiczorek M, Dunn A. Relationships among the behavioral, noradrenergic, and pituitary-adrenal responses to interleukin-1 and the effects of indomethacin. *Brain Behav Immun* 2006;20: 477-87.

- [35] Wand GS, Oswald LM, McCaul ME, Wong DF, Johnson E, Zhou Y, Kuwabarsa H, Kumar A. Association of amphetamine-induced striatal dopamine release and cortisol responses to psychological stress. *Neuropsychopharmacol* 2007;32: 2310-20.
- [36] Debold EP. Recent insights into the molecular basis of neuromuscular fatigue. *Med Sci Sports Exerc* 2012;44: 1440-52.
- *[37] Vuillerme N, Forestier N, Nougier V. Attentional demands and postural sway: the effect of the calf muscles fatigue. *Med Sci Sports Exerc* 2002;34: 1907–12.
- [38] Forestier N, Teasdale N, Nougier V. Alteration of the position sense at the ankle induced by muscular fatigue in humans. *Med Sci Sports Exerc* 2002;34: 117–22
- [39] Redfern MS, Jennings JR, Martin C, Furnam JM. Attention influences sensory integration for postural control in older adults. *Gait Posture* 2001;14: 211–6,
- [40] Fontana FE, Mazzardo O, Mokgothu C, Furtado O, Gallagher JD. Influence of exercise intensity on the decision-making performance of experienced and inexperienced soccer players. *J Sport Exerc Psychol* 2009;31: 135–51.
- [41] McMorris T, Sproule J, Turner A, Hale BJ. Acute, intermediate intensity exercise, and speed and accuracy in working memory tasks: a meta-analytical comparison of effects. *Physiol. Behav* 2011;102: 421-8.
- [42] Arts FJP, Kuipers H. The relation between power output, oxygen uptake and heart rate in male athletes. *Int J Sports Med* 1994;15:228–31
- [43] Borenstein M, Hedges LV, Higgins JPT, Rothstein HR. *Introduction to meta-analysis*. Chichester: John Wiley and Sons; 2009.
- [44] Orwin RG. A fail-safe N for effect size in meta-analysis. *J Educ Stat* 1983;8: 157–9.
- *[45] Lyons M, Al-Nakeeb Y, Nevill A. Performance of soccer passing skills under moderate and high-intensity localized muscle fatigue. *J Strength Cond Res* 2006;20: 197-202

- [46] Etnier JL, Salazar W, Landers DM, Petruzzello SJ, Han M, Nowell P. The influence of physical fitness and exercise upon cognitive functioning: A meta-analysis. *J Sport Exerc Psychol* 1997;19: 249–77
- [47] Chang YK, Labban JD, Gapin JI, Etnier JL. The effects of acute exercise on cognitive performance: A meta-analysis. *Brain Res* 2012;1453: 87-101
- [48] Roig M, Nordbrandt S, Geertsen SS, Nielsen JB. The effects of cardiovascular exercise on human memory: A review with meta-analysis *Neurosci Biobehav Rev* 2013;37: 1645–66
- [49] Lambourne K, Tomporowski PD. The effect of acute exercise on cognitive task performance: A meta-regression analysis. *Brain Res* 2010;1341: 12-24
- [50] Ingjer F, Strømme SB. Effects of active, passive or no warm-up on the physiological response to heavy exercise. *Eur J Appl Physiol* 1979;40: 272-82.
- [51] Wang G-J, Volkow ND, Fowler JS, Francheschi D, Logan J, Pappas NR, Wong CT, Netusil N. PET studies of the effects of aerobic exercise on human striatal dopamine release. *J Nucl Med* 2000;41: 1352-6.
- [52] Vijayraghavan S, Wang M, Birnbaum SG, Bruce CJ, Williams GV, Arnsten AFT. Inverted-U dopamine D1 receptor actions on prefrontal neurons engaged in working memory. *Nat Neurosci.* 2007; 10:376–84.
- [53] Clark-Carter D. *Doing quantitative psychological research*. Hove: Psychology Press; 2002.
- *[54] McRae KA, Galloway SDR. Carbohydrate-electrolyte drink ingestion and skill performance during and after 2 hr of indoor matchplay. *Int J Sport Nutr Exe* 2012;22: 38-46
- *[55] Bullock W, Panchuk D, Broatch J, Christian R, Stepto NK. An integrative test of agility, speed and skill in soccer: Effects of exercise. *J Sci Med Sport* 2012;15: 431–6

*[56] Lidor R, Arnon M, Hershkoj Y, Maayan G, Falk B. Accuracy in a volleyball service test in rested and physical exertion conditions in elite and near-elite adolescent players. *J Strength Cond Res* 2007;21: 937-42

*[57] McMorris T, Sproule J, Draper S, Child R. Performance of a psychomotor skill following rest, exercise at the plasma epinephrine threshold and maximal intensity exercise. *Percept Mot Skills* 2000;91: 553-62.

*[58] McMorris T, Swain J, Lauder M, Smith N, Kelly J. Warm-up prior to undertaking a dynamic psychomotor task: does it aid performance? *J Sport Med Phys Fit* 2006;46: 328-34

[59] Green HJ, Hughson RL, Orr GW, Ranney DA. Anaerobic threshold, blood lactate, and muscle metabolites in progressive exercise. *J Appl Physiol* 1983;54: 1032-8.

[60] Lehman M, Schmid P, Keul J. Plasma catecholamine and blood lactate cumulation during incremental exhaustive exercise. *Int J Sports Med* 1985;6: 78-81.

[61] Podolin DA, Munger PA, Mazzeo RS. Plasma-catecholamine and lactate response during graded-exercise with varied glycogen conditions. *J Appl Physiol* 1991;71: 1427-33.

[62] Urhausen A, Weiler B, Coen B, Kindermann W. Plasma catecholamines during endurance exercise of different intensities as related to the individual anaerobic threshold. *Eur J Appl Physiol* 1994;69: 16-20.

[63] McMorris T, Sproule J, Child R, Draper S, Sexsmith JR. The measurement of plasma lactate and catecholamine thresholds: a comparison of methods. *Eur J Appl Physiol* 2000;82: 262-7.

[64] Córdova C, Silva VC, Moraes CF, Simões HG, Nóbrega OT. Acute exercise performed close to the anaerobic threshold improves cognitive performance in elderly females. *Braz J Med Biol Res* 2009;42: 458-64

[65] Kashihara K, Nakahara Y. Short-term effect of physical exercise at lactate threshold on choice reaction time. *Percept Mot Skills* 2005;100: 275-91.

- [66] Collardeau M, Brisswalter J, Audiffren M. Effects of a prolonged run on simple reaction time of well trained runners. *Percept Mot Skills* 2001;93: 679-89.
- [67] Davranche K, McMorris T. Specific effects of acute moderate exercise on cognitive control. *Brain Cognition* 2009;69: 565–70.
- [68] Del Giorno JM, Hall EE, O’Leary KC, Bixby WR, Miller PC. Cognitive function during acute exercise: A test of the transient hypofrontality theory. *J Sport Exerc Psychol* 2010;32: 312-23
- [69] Ferris LT, Williams JS, Shen C. The effect of acute exercise on serum brain-derived neurotrophic factor levels and cognitive function. *Med Sci Sports Exerc* 2007;39: 728-34.
- [70] Hyodo K, Dan I, Suwabe K, Kyutoku Y, Yamada Y, Akahori M, Byun K, Kato M, Soya H. Acute moderate exercise enhances compensatory brain activation in older adults. *Neurobiol Aging* 2012;33: 2621–32
- [71] Yamamoto Y, Miyashita M, Hughson RL, Tmura S, Shiohara M, Mutoh Y. The ventilatory threshold gives maximal lactate steady state. *Eur J Appl Physiol* 1991;63: 55-9.
- [72] Hagbarth KE, Macefield VG. The fusimotor system: Its role in fatigue. *Adv Exp Med Biol* 1995;384: 259-70.
- [73] Sieck GC, Prakash YS. Fatigue at the neuromuscular junction: Bench point versus presynaptic versus postsynaptic mechanisms. *Adv Exp Med Biol* 1995;384: 83-100.
- [74] Westerblad H, Bruton JD, Allen DG, Lännergren J. Functional significance of Ca^{2+} in long-lasting fatigue of skeletal muscle. *Eur J Appl Physiol* 2000;83:, 166-74.
- [75] Vollestad NK. Measurement of human muscle fatigue. *J Neurosci Meth* 1997;74: 219-27.
- [76] Karlsson J, Saltin B. Diet muscle glycogen and endurance performance. *J Appl Physiol* 1971;21: 203-6.
- [77] Fitts RH. Cellular mechanisms of muscle fatigue. *Physiol Rev* 1994;74: 49-94.

- [78] Foote SL, Freedman FE, Oliver AP. Effects of putative neurotransmitters on neuronal activity in monkey auditory cortex. *Brain Res* 1975;86: 229–42.
- [79] Waterhouse BD, Moises HC, Woodward DJ. Noradrenergic modulation of somatosensory cortical neuronal responses to iontophoretically applied putative transmitters. *Exp Neurol* 1980;69: 30–49.
- [80] Waterhouse BD, Moises HC, Woodward DJ. Alpha-receptor-mediated facilitation of somatosensory cortical neuronal responses to excitatory synaptic inputs and iontophoretically applied acetylcholine. *Neuropharmacology* 1981;20: 907–20.
- [81] Goldman-Rakic, PS. The nervous system, higher functions of the brain. In: Plum F, editor. *Handbook of Physiology Vol. V*. Bethesda: American Physiological Society; 1987. p. 373-417.
- [82] Aron AR, Robbins TW, Poldrack RA. Inhibition and the right inferior frontal cortex. *Trends Cogn. Sci* 2004;8: 170–7
- [83] Asbach S, Schulz C, Lehnert, H. Effects of corticotropin-releasing hormone on locus coeruleus neurons in vivo: a microdialysis study using a novel bilateral approach. *Eur J Endocrinol* 2001;145: 359-63.
- [84] Van Bockstaele EJ, Colago EE, Valentino RJ. Corticotropin-relapsing factor-containing axon terminals synapse onto catecholamine dendrites and may presynaptically modulate other afferents in the rostral pole of the nucleus locus coeruleus in the rat brain. *J Comp Neurol* 1996;364: 523-34.
- [85] Marinelli M, Piazza PV. Interaction between glucocorticoid hormones , stress and psychostimulant drugs. *Eur J Neurosci* 2002;16: 387-94.
- [86] Oswald LM, Wong DF, McCaul M, Zhou Y, Kuwabara H, Choi L, Brasic J, Wand GS. Relationships among ventral striatal dopamine release, cortisol secretion, and subjective responses to amphetamine. *Neuropsychopharmacol* 2005;30: 821-32.

- [87] Wand GS, Oswald LM, McCaul ME, Wong DF, Johnson E, Zhou Y, Kuwabarsa H, Kumar A. Association of amphetamine-induced striatal dopamine release and cortisol responses to psychological stress. *Neuropsychopharmacol* 2007;32: 2310-20.
- [88] Pruessner JC, Champagne F, Meaney MJ, Dagher A. Dopamine release in response to a psychological stress in humans and its relationship to early life maternal care: a positron emission tomography study using [¹¹C]Raclopride. *J Neurosci* 2004;24: 2625-831.
- [89] Gribble PA, Hertel J. Effect of hip and ankle muscle fatigue on unipedal postural control. *J Electromyol Kinesiol* 2004;14: 641-6.
- [90] Gribble PA, Hertel J. Effect of lower extremity muscle fatigue on postural control. *Arch Phys Med Rehab* 2004;85: 589-92
- *[91] Yaggie J, Armstrong WJ. Effects of lower extremity fatigue on indices of balance. *J Sport Rehabil* 2004;13: 312-22.
- [92] McMorris T, Graydon, J. The effect of exercise on the decision-making performance of experienced and inexperienced soccer players. *Res Q Exerc Sport* 1996;67: 109-14.
- [93] Masters RSW, Poolton JM, Maxwell JP. Stable implicit motor processes despite aerobic locomotor fatigue. *Conscious Cogn* 2008;17: 335–8
- [94] Mason JW, Hartley LH, Kotchen TA, Mougey EH, Ricketts PT, Jones LG. Plasma cortisol and norepinephrine responses in anticipation of muscular exercise. *Psychosom Med* 1973;35: 406-14.
- [95] McMorris T, Davranche K, Jones G, Hall B, Minter C. Acute incremental exercise, performance of a central executive task, and sympathoadrenal system and hypothalamic-pituitary-adrenal axis activity. *Int J Psychophysiol* 2009;73: 334-40.
- [96] Peirce NS, Forster CD, Heptinstall S, MacDonald IA. Unexpected changes in the catecholamines content of platelets and plasma during exercise. *Platelets* 1999;10: 312-8.

- *[97] Ali A, Williams C, Nicholas CW, Foskett A. The influence of carbohydrate-electrolyte ingestion on soccer skill performance. *Med Sci Sports Exerc* 2007;39: 1969-76.
- [98] Nicholas CW, Nuttall FE, Williams C. The Loughborough Intermittent Shuttle Test: a field test that stimulates the activity pattern of soccer. *J Sports Sci* 2000;18: 97-104,
- [99] Ali A, Williams C, Hulse MA, Strudwick A, Reddin J, Howarth L, Eldred S, Hirst M, McGregor S.. Reliability and validity of two tests of soccer skill. *J Sports Sci* 2007;25: 1461-70.
- [100] Edwards, B. J., Lindsay, K., & Waterhouse, J. Effect of time of day on the on the accuracy and consistency of the badminton serve. *Ergonomics* 2005;15: 1488–98
- *[101] Anshel MH, Novak J. Effects of different intensities of fatigue on performing a sports skill requiring explosive muscular effort: a test of the specificity of practice principle. *Percept Mot Skills* 1989; 69: 1379-89.
- *[102] Bottoms L, Sinclair J, Taylor K, Polman R, Fewtrell D. The effects of carbohydrate ingestion on the badminton serve after fatiguing exercise. *J Sports Sci* 2012;30: 285–93
- *[103] Davey PR, Thorpe RD, Williams C. Fatigue decreases skilled tennis performance. *J Sports Sci* 2002;20: 311-8
- *[104] Dickin DC, Doan JB. Postural stability in altered and unaltered sensory environments following fatiguing exercise of lower extremity joints. *Scand J Med Sci Sports* 2008;18: 765–72
- *[105] Evans RK; Scoville C, Ito MA, Mello RP. Upper body fatiguing exercise and shooting performance. *Mil Med* 2003;168: 451–6.
- *[106] Gabbett TJ. Influence of fatigue on tackling technique in Rugby League players. *J Strength Cond Res* 2008;22: 625-32.

- *[107] Gros Lambert A, Gillot G, Davenne D, Rouillon JD. Influence de l'exercice physique sur la qualité du tir en biathlon (Influence of physical exercise on the quality of shooting in a biathlon). *Sci Sport* 1995;10: 47-8
- *[108] Impellizzeri FM, Rampinini E, Maffiuletti NA, Castagna C, Bizzini M, Wisløff U. Effects of aerobic training on the exercise-induced decline in short-passing ability in junior soccer players. *Appl Physiol Nutr Metab* 2008;33: 1192–8
- [109] Rampinini E, Impellizzeri FM, Castagna C, Azzalin A, Ferrari Bravo D, Wisløff U. Effect of match-related fatigue on short passing ability in young soccer players. *Med Sci Sports Exerc* 2008;40: 934–42.
- *[110] Johnston III RB, Howard ME, Cawley PW, Losse GM. Effect of lower extremity muscular fatigue on motor control performance. *Med Sci Sports Exerc* 1998;30: 1703-7.
- *[111] Lyons M, Al-Nakeeb Y, Nevill A. The impact of moderate and high intensity total body fatigue on passing accuracy in expert and novice basketball players. *J Sports Sci Med* 2006;5: 215-27
- *[112] McGregor SJ, Nicholas CW, Lakomy HKA, Williams C. The influence of intermittent high-intensity shuttle running and fluid ingestion on the performance of a soccer skill. *J Sports Sci* 1999;17: 895-903
- *[113] McMorris T, Gibbs C, Palmer J, Payne A, Torpey N. Exercise and performance of a motor skill. *Brit J Phys Ed Res Suppl* 1994;15: 23-7.
- *[114] McMorris T, Delves S, Sproule J, Lauder M, Hale B. Incremental exercise, perceived task demands and performance of a whole body motor task. *Brit J Sport Med* 2005;39: 537-41.
- *[115] McMorris T, Rayment T. Short-duration, high-intensity exercise and performance of a sports-specific skill: a preliminary study. *Percept Mot Skills* 2007;105: 523-30.

*[116] Sarshin A, Mohammadi S, Babael H, Shahrabad P, Sedighi M. The effects of functional fatigue on dynamic postural control of badminton players. *Biol Exerc* 2011;7: 25-34.

*[117] Sunderland C, Nevill ME. High-intensity intermittent running and field hockey skill performance in the heat. *J Sports Sci* 2005;23: 531–40

*[118] Tharion WJ, Montain SJ, O'Brien C, Shippee RL., Hoban JL. Effects of military exercise tasks and a carbohydrate-electrolyte drink on rifle shooting performance in two shooting positions. *Int J Ind Ergonom* 1997;19: 31-9.

*[119] Wikstrom EA, Powers ME, Tillman MD. Dynamic stabilization time after isokinetic and functional fatigue *J Athl Training* 2004;39:, 247–53.

[120] Ayalon A, Inbar O, Bar-Or O. Relationships among measurements of explosive strength and anaerobic power 1974 In: Nelson RC, Morehouse CA, editors. *Biomechanics IV International series on sport sciences 1*. Baltimore: University Press; 1974, p 572-7.

*[121] Young W, Gulli R, Rath D, Russell A, O'Brien B, Harvey J. Acute effect of exercise on kicking accuracy in elite Australian football players. *J Sci Med Sport* 2010;13: 85–9.

Figure captions

Figure 1. Forest plot for all studies.

Note: H heavy exercise; M moderate intensity exercise; D dynamic/ballistic skills; S static balance; CR counterbalanced or random design; SO (same order) pre-exercise test followed by post-exercise test.

Figure 2. Forest plot for heavy exercise studies.

Note: H heavy exercise; D dynamic/ballistic skills; S static balance; CR counterbalanced or random design; SO (same order) pre-exercise test followed by post-exercise test.

Figure 3. Forest plot for dynamic/ballistic skills studies.

Note: H heavy exercise; M moderate intensity exercise; D dynamic/ballistic skills; CR counterbalanced or random design; SO (same order) pre-exercise test followed by post-exercise test.

Figure 4. Forest plot for static balance studies.

Note: H heavy exercise; M moderate intensity exercise; S static balance; CR counterbalanced or random design; SO (same order) pre-exercise test followed by post-exercise test.