

Northumbria Research Link

Citation: Jones, Thomas, Howatson, Glyn, Russell, Mark and French, Duncan (2016) Performance and endocrine responses to differing rations of concurrent strength and endurance training. *Journal of Strength and Conditioning Research*, 30 (3). pp. 693-702. ISSN 1064-8011

Published by: Lippincott, Williams & Wilkins

URL: <http://doi.org/10.1519/JSC.0000000000001135>
<<http://doi.org/10.1519/JSC.0000000000001135>>

This version was downloaded from Northumbria Research Link:
<http://nrl.northumbria.ac.uk/25940/>

Northumbria University has developed Northumbria Research Link (NRL) to enable users to access the University's research output. Copyright © and moral rights for items on NRL are retained by the individual author(s) and/or other copyright owners. Single copies of full items can be reproduced, displayed or performed, and given to third parties in any format or medium for personal research or study, educational, or not-for-profit purposes without prior permission or charge, provided the authors, title and full bibliographic details are given, as well as a hyperlink and/or URL to the original metadata page. The content must not be changed in any way. Full items must not be sold commercially in any format or medium without formal permission of the copyright holder. The full policy is available online: <http://nrl.northumbria.ac.uk/policies.html>

This document may differ from the final, published version of the research and has been made available online in accordance with publisher policies. To read and/or cite from the published version of the research, please visit the publisher's website (a subscription may be required.)

www.northumbria.ac.uk/nrl



1 **PERFORMANCE AND ENDOCRINE RESPONSES TO DIFFERING RATIOS OF**
2 **CONCURRENT STRENGTH AND ENDURANCE TRAINING**

3

4 **PERFORMANCE RESPONSES TO DIFFERING RATIOS OF CONCURRENT**
5 **TRAINING**

6

7 THOMAS. W JONES¹, GLYN HOWATSON^{2,3}, MARK RUSSELL² and DUNCAN N.
8 FRENCH^{2,4}

9 *¹ASPIRE, Academy for Sports Excellence, Doha, Qatar*

10 *²Department Sport, Exercise and Rehabilitation, Northumbria University, Newcastle*
11 *upon Tyne, United Kingdom*

12 *³Water Research Group, School of Environmental Sciences and Development, North*
13 *West University, Potchefstroom, South Africa*

14 *⁴English Institute of Sport, Sportcity, Manchester, United Kingdom*

15

16 Dr Duncan N. French

17 Department of Sport and Exercise, and Rehabilitation

18 Northumbria University

19 Northumberland Building

20 Newcastle upon Tyne

21 NE1 8ST

22 duncan.french@northumbria.ac.uk

23

1 ABSTRACT

2 The present study examined functional strength and endocrine responses to varying
3 ratios of strength and endurance training in a concurrent training regimen. 30
4 resistance-trained men completed 6 weeks of 3 d·wk⁻¹ of i) strength training (ST), ii)
5 concurrent strength and endurance training ratio 3:1 (CT3), iii) concurrent strength
6 and endurance training ratio 1:1 (CT1) or iv) no training (CON). Strength training was
7 conducted using whole-body, multi-joint exercises, while endurance training
8 consisted of treadmill running. Assessments of maximal strength, lower body power,
9 and endocrine factors were conducted pre-training and following 3 and 6 weeks.
10 Following the intervention ST and CT3 elicited similar increases in lower body
11 strength; furthermore, ST resulted in greater increases than CT1 and CON (all $p <$
12 0.05). All training conditions resulted in similar increases in upper body strength
13 following training. ST group observed greater increases in lower body power than all
14 other conditions (all $p < 0.05$). Following the final training session, CT1 elicited
15 greater increases in cortisol than ST ($p = 0.008$). When implemented as part of a
16 concurrent training regimen, higher volumes of endurance training result in the
17 inhibition of lower body strength, whereas low volumes do not. Lower body power
18 was attenuated by high and low frequencies of endurance training. Higher
19 frequencies of endurance training resulted in increased cortisol responses to
20 training. These data suggest that if strength development is the primary focus of a
21 training intervention, frequency of endurance training should remain low.

22

23 **KEY WORDS** combined exercise, interference, cortisol, resistance training, training
24 frequency

25

1 INTRODUCTION

2 Various sports and events require contrasting physical performance phenotypes for
3 successful performance. Training for sports and events at the extremes of the
4 strength-endurance continuum, such as Powerlifting and ultra-endurance challenges,
5 is relatively straight-forward compared with sports and events that require a
6 combination of strength and endurance capabilities. In these situations athletes and
7 coaches are often forced to combine training methods which elicit contrasting and
8 even antagonistic physiological and performance responses (12). In the case of
9 '*concurrent training*', the divergent stimuli of strength and endurance training can
10 result in attenuated strength type adaptation when compared to strength training
11 performed in isolation. This divergent physiology is known as the interference effect
12 or phenomenon (17).

13
14 Research has indicated that any interference experienced during a concurrent
15 strength and endurance training regimen may be dependent in part on the volume of
16 training performed (1, 13, 24, 25, 33). Despite this, no study has specifically
17 examined the effects of whole body, multi-joint concurrent training interventions with
18 varying training volumes and the effect that it has on muscle force characteristics.
19 Previous work from our laboratory (20) has indicated that the magnitude of
20 interference experienced may be proportional to the frequency of endurance training
21 performed; indicating overall training volume and exercise stress may indeed
22 regulate the presence of any interference experienced.

23
24 Elevated training 'stress' has previously been proposed as a mechanism for
25 interference (10), and is perhaps attributable to the experimental design of some

1 published studies in this area. Often the concurrent training condition will perform
2 double the overall training volume and total work to that of the strength training alone
3 condition, which has previously resulted in muted strength development (6, 16, 20,
4 22). In contrast studies employing lower concurrent training volumes have reported
5 no inhibited strength development as a result of concurrent training (24, 25). These
6 findings may support the hypothesis that total work performed in a concurrent
7 programme influences both the presence and magnitude of any interference
8 experienced, although the underlying mechanisms are yet to be fully elucidated.

9

10 Previous research has reported a decreased testosterone:cortisol ratio following
11 concurrent training with no such decrease in participants who performed strength
12 training alone (2, 3, 22). This may implicate elevated endocrine responses and
13 catabolism as a contributing factor to interference. As such, it is reasonable to
14 suggest that the higher training volumes experienced in concurrent training regimens
15 can result in elevated physiological stress, which is reflected in the responses of
16 primary anabolic and catabolic hormones. This shift in the endocrine milieu in favour
17 of catabolism may contribute to attenuated strength and hypertrophic adaptation
18 associated with concurrent training.

19

20 Previous work from our laboratory (20) illustrates the value in exploring the role of
21 training frequency in a systematic fashion. Furthermore no research has assessed if
22 differing ratios of strength and endurance training can influence the degree of
23 interference experienced as a result of adaptations in the anabolic:catabolic
24 environment. Therefore, the purpose of this research was to investigate the strength,

1 anthropometric and endocrine responses to a variety of concurrent strength and
2 endurance training ratios, with incremental loads in a functional multi joint model.

3

4 **METHOD**

5 **Experimental Approach to the Problem**

6 A balanced, randomized, between-group study design was employed to examine the
7 effect of differing ratios of strength and endurance training in a concurrent regimen
8 on strength, anthropometric, and endocrine responses. A 6 week training
9 intervention was completed, during which participants were randomly assigned to
10 one of four experimental conditions: either i) strength training alone (ST), ii)
11 concurrent strength and endurance training at a ratio of 3:1 (CT3), iii) concurrent
12 strength and endurance training at a ratio of 1:1 (CT1), or iv) no training (CON).
13 Participants in the ST group were required to perform strength training alone on all
14 scheduled training sessions. The CT3 group completed strength training on every
15 scheduled session with every third session immediately followed by an endurance
16 training protocol. Elsewhere, participants designated CT1 completed an identical
17 strength training protocol immediately followed by endurance training at every
18 scheduled session. Those participants in the CON group performed no strength or
19 endurance training during the entire experimental period. Due to the requirements of
20 the separate training protocols, it was not possible to match total work performed in
21 the respective experimental conditions. All participants were instructed to abstain
22 from any other strength or endurance training throughout the experimental period
23 beyond that prescribed by the investigator.

24

1 Participants completed their respective intervention 3 d·wk⁻¹ with ~48 h between
2 sessions for 6 weeks resulting in a total of 18 separate training sessions in the micro
3 cycle. In order to assess whether the frequency and ratio of strength and endurance
4 training performed influenced strength and changes in body composition,
5 assessments of 1 repetition maximums (1RM), countermovement jump height
6 (CMJ), and body composition were assessed pre, mid and post-intervention. To assess
7 the effect of the designated training interventions on endocrine factors related to
8 strength and morphological adaptation, venous blood samples were taken and
9 subsequently analysed for circulating testosterone and cortisol concentrations.
10 During the investigation, venous blood samples were collected immediately before
11 (pre) and following the cessation of exercise (post) in the initial, mid and final
12 compound training sessions of the 18 sessions performed.

13

14 **Subjects**

15 Prior to all experimental procedures the study was approved by the Northumbria
16 University research ethics committee. All subjects were informed of the risks and
17 benefits of the investigation prior to signing an approved informed consent document
18 to participate in the study. Thirty healthy, recreationally resistance-trained men (age:
19 23 ± 4 y; body mass: 79.2 ± 6.7 kg; height: 179.2 ± 6.7 cm; % body fat: 16.2 ± 5.4 %;
20 sum of assessed 1RMs: 506.0 ± 11.4 kg; CMJ: 52.5 ± 7.3 cm; $\dot{V}O_{2max}$: 50.2 ± 5.8
21 ml·kg⁻¹·min) volunteered to participate in the study. Prior to commencing, participants
22 were matched for age, body mass, body fat % and 1RM (sum off all assessed 1RMs)
23 load (all $p > 0.05$), and then randomly assigned (via block randomisation) to one of
24 the four experimental conditions. Each participant had completed > 2 years of
25 strength training activities prior to the start of a study, and were considered

1 recreationally “resistance trained”; all participants were conducting strength training \geq
2 $2 \text{ d}\cdot\text{wk}^{-1}$, however none were involved in a sport-specific training programme. All
3 participants were non-smokers, free from any endocrine or metabolic
4 contraindications, and were not following any specialized dietary interventions. In all
5 cases participants were asked to refrain from nutritional supplementation or
6 pharmacological interventions for 30 days prior to and throughout the duration of any
7 experimental intervention.

8

9 **Procedures**

10 ***Strength training protocol***

11 Prior to the intervention all participants completed a familiarisation week involving
12 each respective training session in order to habituate themselves fully with the
13 exercise techniques employed. The strength training intervention was comprised of 3
14 sessions, and each was performed on separate days with ~ 48 h between sessions.
15 Each session was composed of differing exercises; as such each of the sessions
16 were designated “compound”, “pull” and “push” respectively, to best describe the
17 nature of exercises performed. Full details of each session are presented in Table 1.
18 The respective sessions were performed in the same order each week (i.e.,
19 compound, push then pull). Furthermore, the order of exercises within each session
20 was consistent throughout the intervention.

21

22 During familiarisation, training intensity was set at 70% of 1 repetition maximum
23 (1RM) for 3 sets of 10 repetitions. The first 3 weeks of the training intervention
24 required participants to complete all sessions and exercises at 80% 1RM for 4 sets
25 of 8 repetitions. The following and final 3 weeks of the intervention were completed

1 at an intensity of 85% 1RM for 5 sets of 6 repetitions. These loads, volumes and rest
2 intervals were selected as they are deemed appropriate for eliciting adaptations in
3 strength and hypertrophy in recreationally trained non-athletes (27, 28). Additionally,
4 strength training programmes of this nature involving exercises which stimulate large
5 muscle masses and shorter rest periods have been shown to elicit large increases in
6 the endocrine factors assessed within this study (21, 32). Full details of the
7 intervention are presented in Table 1.

8

9 All strength and/or endurance-based exercise commenced at the same time of day
10 (1000 h \pm 1 h) to avoid any diurnal performance or endocrine variations (15).
11 Participants were also advised to abstain from exercise for 24 h prior to a visit.
12 Training load was modified accordingly for each exercise if a participant's 1RMs
13 were observed to change at the mid-intervention assessments. Compliance was
14 100% for all participants.

15

16 *Table 1 about here*

17

18 ***Endurance training protocol***

19 In all instances endurance training was conducted immediately following strength
20 training. The endurance training protocol required participants to run on a treadmill
21 (hp Cosmos, Pulsar, Nussdorf-Traunstein, Germany) at 1% incline at 70% of their
22 pre-determined peak running velocity at $\dot{V}O_{2\max}$ ($v\dot{V}O_{2\max}$). Running velocity was
23 modified if participant's $v\dot{V}O_{2\max}$ was observed to change at the mid-intervention
24 assessments.

25

1 **Whole body strength assessments - 1 repetition maximum (1RM)**

2 1RM loads were established for all strength-training exercises prior to the
3 experimental intervention and following 3 and 6 weeks of training. For analysis
4 purposes lower body strength was assessed via back squat and deadlift 1RM total
5 load. To examine strength development in the upper body musculature, bench press,
6 bent over row and military press total 1RM load was analysed. These exercises were
7 chosen as they are considered gross motor movements that require all the major
8 joints and muscle groups involved in the strength training intervention. All
9 assessments were conducted in line with standardised procedures (29).

10

11 **Maximal aerobic capacity - $\dot{V}O_{2max}$**

12 Assessments of participant's maximal oxygen uptake and peak running velocity at
13 $\dot{V}O_{2max}$ were conducted at baseline, after 3 weeks of training and following the 6
14 week training intervention. All assessments were conducted in line with standardised
15 procedures reported elsewhere (34).

16

17 **Lower body power - countermovement jump assessment**

18 Lower body power was assessed via maximal countermovement jump height (CMJ)
19 and was conducted prior to and following 3 and 6 weeks of training. Maximal CMJ
20 was adopted as a proxy of lower body power, and was assessed using a contact mat
21 (Just Jump, Probotics, Huntsville, AL, USA). Following familiarization, independent
22 trials of CMJs were conducted with 3 min between each individual jump; the highest
23 jump being recorded for data analysis. When performing the test, participants
24 positioned themselves in the centre of the contact mat and place their hands on the
25 iliac crest where they were to remain throughout. CMJs began from an erect

1 standing position. When ready, participants squatted to a self-selected depth
2 perceived as their individual optimal depth, and immediately ascended to jump
3 vertically for maximal height.

4

5 ***Body composition - air displacement plethysmography***

6 All participants lean mass and % body fat was assessed prior to and following 3 and
7 6 weeks of training. Lean mass and % body fat were assessed using air
8 displacement plethysmography (BodPod, Life Measurements Instruments, CA, USA)
9 (11, 26, 30). Initially the device was calibrated using a metal cylinder of known and
10 standardised composition. Participants were asked to disrobe to minimal clothing
11 and place a tight fitting cap over their hair. Participants were then weighed on a
12 calibrated scale prior to entering the chamber. Once two consistent measures of
13 body composition were obtained % body fat and lean mass were calculated using
14 associated software (8).

15

16 ***Rate of perceived exertion***

17 To examine perception of physical exertion in response to the training intervention,
18 rate of perceived exertion (RPE) was recorded during strength training. Briefly,
19 participants were required to select a number from 6 to 20, corresponding to a
20 statement which best described their level of exertion at that particular moment (4, 7,
21 31).

22

23 ***Blood sampling and storage***

24 When blood samples were collected, participants arrived at the lab having refrained
25 from consuming food or caffeine for 2 h prior to assessment. Venous blood samples

1 were collected from the antecubital fossa in a branch of the basilica vein into
2 vacutainer tubes (BD Vacutainer, NJ, USA) coated with Ethylenediaminetetraacetic
3 acid (EDTA) to negate. Whole blood was subsequently centrifuged (accuSpin 3R,
4 Fisher Scientific, Loughborough, UK) at 4°C and 1509 g for 10 min, after which the
5 resultant plasma from each sample was then transferred to individual eppendorf
6 containers for subsequent storage at -80°C. Venous blood samples were collected
7 immediately before (pre) and following the cessation of exercise (post) in the initial,
8 mid and final compound training sessions (additional information presented in Table
9 1) of the 18 sessions performed.

10

11 ***Biochemical analysis***

12 Plasma testosterone and cortisol were measured in duplicate (testosterone; ICC =
13 0.89, $R = 0.89$, Cortisol: ICC = 0.92, $R = 0.95$) via commercially available enzyme-
14 linked immunosorbent assay (ELISA) kits (IBL International, Hamburg, Germany). In
15 all cases procedures were followed according to the manufacturer's instructions. For
16 both variables, 25 μ L of each standard, control and sample were pipetted into the
17 respective wells of the microtitre plate, after which 2000 μ L of enzyme conjugate was
18 then pipetted into each well and the plate was covered and left to incubate at room
19 temperature (18 - 25°C) for 60 min. After this period the incubation solution was
20 discarded and the microplate was washed 3 times with wash buffer and distilled
21 water solution diluted at a ratio of 1:10. 100 μ L of Tetramethylbenzidine (TMB)
22 substrate solution was then pipetted into each well prior to a 15 min incubation
23 period. Immediately following this incubation 100 μ L of TMB stop solution was
24 pipetted into each well and the contents were briefly mixed by gently agitating the
25 plate. The optical density was measured at 450 nm within 10 min of the stop solution

1 being added using an Anthos 2010 microplate reader (DAZDAQ LTD, Brighton, UK
2 (reference-wavelength 600 – 650 nm)). For testosterone there was a minimum
3 detection limit of 0.2 nmol·L⁻¹, inter-assay and intra-assay variation of 4.2 – 7.4 and
4 3.1 – 5.4 and the calibration curve revealed Pearson's correlation coefficients (r) =
5 0.99. For cortisol there was a minimum detection limit of 6.8 nmol·L⁻¹ with an inter-
6 assay and intra-assay variation of 2.1 – 5.0 and 2.6 – 3.5, the calibration curve
7 revealed $r = 0.99$, respectively.

8

9 **Statistical analysis**

10 Data are presented as mean \pm standard deviation. Values of RMs, CMJ and lean
11 mass were transformed to a percentage change ($\Delta\%$) from baseline and used for
12 analysis. Prior to analysis, dependant variables were verified as meeting required
13 assumptions of parametric statistics and changes in all assessed measures were
14 analysed using mixed model repeated measures ANOVA tests. ANOVA analysed
15 differences between 4 conditions (ST, CT3, CT1 and CON) and 3 time points
16 (baseline, mid-intervention and post-intervention). The alpha level of 0.05 was set
17 prior to data analysis. Assumptions of sphericity were assessed using Mauchly's test
18 of sphericity, if the assumption of sphericity was violated Greenhouse Gessier
19 correction was employed. If significant effects between conditions or over time were
20 observed *post-hoc* differences were analysed with the use of Bonferroni correction.
21 Statistical power of the study was calculated post-hoc using G*Power statistical
22 software (v3.1.3, Düsseldorf, Germany) using the effect size, group mean, SD and
23 sample size of the primary outcome measures, in this case being lower and upper
24 body maximal strength and endocrine factors. Power was calculated as between 0.8
25 and 1 indicating sufficient statistical power (5).

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25

RESULTS

Physical performance measures

Participant's baseline strength and endurance physical performance capabilities were similar between experimental conditions, these data are presented in Table 2.

Table 2 about here

Upper and lower body maximal strength

A significant group x time interaction was observed ($F_{(4, 36)} = 4.940, p = 0.003$) for lower body strength development, as was an effect of time ($F_{(1, 36)} = 45.042, p < 0.001$). All training conditions elicited increases in lower body strength at the mid-intervention time point following 3 weeks of training (ST; $9.0 \pm 4.5\%$, $p < 0.001$. CT3; $9.8 \pm 11.0\%$, $p = 0.024$. CT1; $5.8 \pm 3.2\%$, $p < 0.001$). Similarly lower body strength improved in all training conditions from baseline to post-intervention (ST; $17.2 \pm 7.2\%$, $p < 0.001$. CT3; $15.0 \pm 11.8\%$, $p = 0.003$. CT1; $10.1 \pm 4.9\%$, $p < 0.001$). ST was the only condition to significantly increase lower body strength from mid to post-intervention ($8.3 \pm 2.8\%$, $p = 0.016$, Figure 1).

Figure 1 about here

All training conditions improved lower body strength to a greater extent than CON at both mid and post-intervention (all $p < 0.05$). Post-training ST improved lower body strength $7.1 \pm 2.4\%$ more than CT1 ($p = 0.036$, Figure 1).

1 A significant group x time interaction ($F_{(5, 41)} = 2.895, p = 0.027$) and an effect of time
2 ($F_{(2, 36)} = 31.510, p < 0.001$) were observed for upper body strength development.
3 CT3 and CT1 both improved upper body strength between baseline to mid-
4 intervention ($6.2 \pm 6.9\%, p = 0.024$ and $7.8 \pm 4.5\%, < 0.001$ respectively, Figure 2).
5 All training conditions increased upper body strength from pre to post-training (all $p <$
6 0.05). Upper body strength improved in all training conditions following training
7 interventions (ST; $10.5 \pm 5.2\%, p < 0.001$. CT3; $10.6 \pm 10.7\%, p = 0.014$. CT1; 12.1
8 $\pm 6.9\%, p < 0.001$). ST was the only condition to improve upper body strength from
9 mid to post-training ($6.9 \pm 0.1\%, p = 0.019$).

10

11

Figure 2 about here

12

13 All training conditions elicited significantly greater increases in upper body strength
14 than CON at mid- and post intervention (all $p < 0.05$, Figure 2).

15

16 **Lower body power**

17 A significant group x time interaction ($F_{(6, 52)} = 3.236, p = 0.009$) and effect of time
18 ($F_{(2, 52)} = 26.086, p < 0.001$) were observed for lower body power development. Both
19 ST and CT1 increased CMJ from baseline to mid-intervention (ST; $8.7 \pm 7.0\%, p =$
20 0.003 . CT1; $3.0 \pm 2.3\%, p = 0.002$). Post-intervention all training conditions elicited
21 significant increases in CMJ from baseline (ST; $13.1 \pm 7.3\%, p < 0.001$. CT3; $7.1 \pm$
22 $3.7\%, p < 0.001$. CT1; $4.8 \pm 2.3\%, p < 0.001$; Figure 3).

23

24

Figure 3 about here

25

1 Participants in the ST condition achieved significantly higher CMJ than those
2 following CT1 ($7.0 \pm 3.5\%$) and CON ($5.7 \pm 4.7\%$) conditions after 3 weeks of
3 training (i.e. mid-intervention) (both $p = 0.04$). Following training (i.e. post-
4 intervention), ST elicited $6.0 \pm 3.6\%$ greater increases in CMJ than CT3, $8.3 \pm 5.0\%$
5 greater than CT1 and $10.9 \pm 2.3\%$ greater than CON (all $p < 0.05$).

7 **Strength training performance**

8 During the first 3 weeks of the training intervention all groups ability to maintain the
9 required training intensity was similar ($F_{(3, 30)} = 1.063$, $p = 0.548$) and did not change
10 significantly over time ($F_{(1, 30)} = 4.295$, $p = 0.062$). Similar results were observed in
11 the final 3 weeks of the intervention as ability to maintain designated training load
12 was not different between conditions ($F_{(3, 28)} = 1.301$, $p = 0.293$) or over time ($F_{(1, 28)}$
13 $= 3.777$, $p = 0.052$).

15 **Testosterone**

16 No group x time interaction was reported for circulating basal testosterone
17 concentrations ($F_{(6, 52)} = 1.820$ $p = 0.113$, Table 3). A significant group x time
18 interaction was however observed for the testosterone response to strength training
19 ($F_{(3, 26)} = 11.466$, $p < 0.001$). Testosterone responses to the respective training
20 interventions also changed significantly over time ($F_{(1, 26)} = 130.683$, $p < 0.001$).
21 Following the initial and mid sessions ST was the only condition to increase
22 testosterone levels greater than CON ($30.7 \pm 5.0\%$, $p = 0.04$ and $37.1 \pm 12.9\%$ $p =$
23 0.005 respectively). CT3 was the only condition to elicit a greater increase in
24 testosterone than CON post the final session ($42.2 \pm 10.5\%$, $p = 0.002$). ST and CT3
25 elicited significant increases from pre training in both the mid and final sessions (all p

1 < 0.05). Testosterone was also increased post training in the CT3 condition following
2 the final session ($p = 0.01$). No other increases were observed.

3

4 **Cortisol**

5 No group x time interaction was observed for circulating basal cortisol concentrations
6 ($F_{(6, 52)} = 1.540$, $p = 0.184$, Table 3). A significant a group x time interaction ($F_{(3, 26)} =$
7 7.592 , $p = 0.001$) and an effect of time ($F_{(1, 26)} = 101.852$, $p < 0.001$) were observed
8 for cortisol responses to the respective training interventions. Following the initial
9 session ST was the only condition to increase cortisol levels to a greater extent than
10 CON ($84.7 \pm 22.1\%$, $p = 0.014$). Post training after the mid-intervention session CT1
11 was the only condition which resulted in significantly greater cortisol increases than
12 CON ($49.2 \pm 3.1\%$, $p < 0.001$). Following the final session, CT1 elicited $26.6 \pm 8.4\%$
13 greater cortisol increases than ST ($p < 0.008$). All training conditions elicited
14 significant increases in cortisol post training on all assessed sessions (all $p < 0.05$).

15

16 **Testosterone-cortisol ratio**

17 No group x time interactions were present for basal testosterone:cortisol ratio (T:C
18 ratio) ($F_{(6, 52)} = 1.903$, $p = 0.098$) nor the T:C ratio response to training ($F_{(6, 52)} =$
19 1.124 , $p = 0.361$).

20

21 *Table 3 about here*

22

23 **Lean mass**

1 Participant's baseline lean mass was similar between experimental conditions, these
2 data are presented in Table 4. No group x time interaction was observed for changes
3 in participant's lean mass.

4

5 *Table 4 about here*

6

7 **Body fat %**

8 A significant group x time interaction was observed for body fat % ($F_{(6, 52)} = 4.616$, p
9 = 0.001). Following the 6 week training intervention, CT1 resulted in $2.65 \pm 0.04\%$
10 greater decreases in body fat % than CON ($p < 0.001$) at the post-intervention time
11 point. No other significant effects of time or group were observed for changes in
12 body fat %.

13

14

15 **Rate of perceived exertion**

16 A significant group x time interaction was present for RPE ($F_{(5, 52)} = 2.744$, $p =$
17 0.029). At week 5 and 6 of the training intervention RPE was significantly lower in the
18 ST group than CT1 (both $p < 0.05$) (Figure 4). No other interactions or effects were
19 present.

20

21 *Figure 4 about here*

22

23 **DISCUSSION**

24 The present study sought to prioritise strength development in concurrent training
25 regimens with varying volumes of endurance training. The primary finding of this

1 study was that an increase in the frequency of endurance training and total training
2 volume within the concurrent training paradigm resulted in the attenuated
3 development of lower body strength when compared to strength training alone.
4 Following 6 weeks of training, ST and CT3 conditions resulted in similar increases in
5 lower body strength, whereas the improvements of those performing both strength
6 and endurance training collectively 3 times per week (CT1) were muted (Figure 1).
7 These findings reflect data presented in our previous work (20), in which ST and CT3
8 resulted in similar increases in maximal voluntary contraction (MVC), whereas
9 increases in the CT1 condition were significantly lower. Although no other published
10 research has examined differing frequencies of strength and endurance training on
11 strength-related adaptation, studies employing concurrent training frequencies of ≥ 3
12 $\text{d}\cdot\text{wk}^{-1}$ have typically reported some manifestation of interference characteristics (2,
13 14, 19, 22). Lower concurrent training frequencies ($\leq 2 \text{ d}\cdot\text{wk}^{-1}$) have however
14 resulted in similar development of strength related phenotypes following both
15 concurrent and strength training programmes (24, 25). When combined, the findings
16 of these studies are consistent with those of the present study. Concurrent training
17 conducted $3 \text{ d}\cdot\text{wk}^{-1}$ (CT1) resulted in inhibited gains in maximal lower body strength,
18 whereas performing concurrent training once per week with 2 strength alone
19 sessions (CT3; concurrent training frequency of $1 \text{ d}\cdot\text{wk}^{-1}$) elicited similar lower body
20 strength increases than strength-training in isolation. The findings of this study and
21 those of previous research indicate higher training volumes and elevated
22 physiological stress may contribute to the presence of the interference phenomenon.
23
24 In addition to the inhibition of lower body strength development lower body power
25 development was also inhibited following 3 and 6 weeks of training in the CT1

1 condition when compared with strength training alone (Figure 3). Furthermore, lower
2 volumes of endurance training also resulted in attenuated increases in lower body
3 power, as post-intervention participants who performed strength and endurance
4 training at a ratio of 3:1 (CT3) exhibited improvements which were $6.0 \pm 3.6\%$ ($p =$
5 0.04 , smallest worthwhile change = 1.2% (18)) lower than those who performed
6 strength training alone. As previously stated, maximal lower body strength
7 development was not different between ST and CT3 conditions (Figure 1), which
8 may indicate that power phenotypes are more susceptible to interference than
9 maximal strength indices. This suggestion is supported by previous research
10 indicating that development of variables including CMJ, rate of force development
11 (RFD) and peak torques at high velocities have been inhibited as a result of
12 combining strength and endurance training, yet maximal strength development
13 remained uninhibited (6, 9, 14).

14

15 Unlike lower body strength and power development, increases in upper body
16 strength were similar following both strength training alone and both concurrent
17 training conditions (CT3 and CT1). Furthermore, following 3 weeks of training CT1
18 resulted in $4.2 \pm 0.8\%$ greater increases than strength training alone (Figure 2),
19 although this was not statistically significant ($p = 0.09$). Previous research has also
20 reported concurrent training does not result in the inhibition of upper body maximal
21 strength (1, 3). Unlike the present study, which employed steady state running,
22 previous research involved rowing (3) and arm cranking (1) as the endurance
23 training modalities. It may be argued that whilst aerobically demanding the stimuli of
24 arm cranking and rowing are further towards the strength end of the strength-
25 endurance continuum than steady state running. As such, it is reasonable to suggest

1 that concurrent training may not differently affect the upper body musculature, but
2 rather for interference to occur the assessed musculature must experience divergent
3 contractile activity (i.e. strength and endurance stimulus) of contrasting intensities
4 and durations. It is reasonable to suggest that the lower body musculature was
5 placed in a greater state of conflict than the upper body, as both training stimuli
6 directly affected hip dominant and lower limb muscle groups and only the strength
7 training protocol required noteworthy contributions from the upper body musculature.
8 Due to the relatively low number of high force contractions involved in strength
9 training and the continuous lower force contractions experienced during endurance
10 training, different patterns of motor unit activation are required. It is possible that the
11 divergent demands placed on the neuromuscular system by strength and endurance
12 training elicited differing alterations in motor unit recruitment in the musculature of
13 the lower limbs, previous research has also implicated altered neural activation
14 during high force contractions as a potential mechanism for impaired strength
15 development (22, 23). Moreover, the potential altered neural recruitment during
16 rapid and high force contractions may have contributed to the inhibition of lower body
17 power development as a result of both high and low frequencies of concurrent training
18 (Figure 3).

19

20 Following the final training session of the intervention CT1 elicited greater cortisol
21 levels than ST which is consistent with previous research (2, 3). This may indicate
22 higher frequencies of concurrent training can result in elevated physiological stress,
23 which was also reflected in participant's perceived exertion during training (Figure 4).
24 In addition to enhanced training stress elevations in cortisol have been implicated
25 in catabolism and impaired hypertrophic development with concurrent training (22).

1 However, in the present study increases in lean mass were similar between training
2 conditions, as such it is unlikely the observed elevations in cortisol influenced muscle
3 morphological adaptation. The variance in the findings of the present study and
4 those of Kraemer *et al.* (22) are perhaps due to the differing lengths of the respective
5 training programmes. Kraemer *et al.* (22) employed a 12 week intervention whereas
6 in the present study participants trained for 6 weeks. As the CT1 condition resulted in
7 the inhibition of strength development following 6 weeks of training it may be
8 speculated that had the interventions been longer CT1 may have also resulted in
9 impaired increases in lean mass.

10

11 **PRACTICAL APPLICATIONS**

12 The findings of this study build on the understanding of concurrent training
13 developed in the isolated limb model discussed in our previous work (20). The data
14 presented here indicate that if strength development is the primary goal of an training
15 programme, endurance-training frequency should be kept to a minimum. It should
16 however be noted, that this minimal dose of endurance training should be sufficient
17 to maintain any necessary endurance performance characteristics. Also the
18 elevations in post exercise cortisol concentrations observed only in participants
19 conducting strength and endurance training 3 times weekly indicate that overall
20 training stress likely plays a key role in the inhibition of strength development.
21 Therefore if a concurrent training programme must be performed it is imperative that
22 appropriate monitoring strategies are employed to ensure training stress doesn't
23 become too great and result in the plateau of strength development. Furthermore if
24 development of power type characteristics is required then it appears that frequency
25 and volume of endurance training should be minimized or omitted from the

- 1 programme all together. This may be achieved via appropriate programme
- 2 construction and periodization to allow power development to occur in periods in
- 3 which endurance type training can be kept to a minimum.

1 REFERENCES

- 2 1. Abernethy PJ and Quigley BM. Concurrent strength and endurance training of the elbow
3 extensors. *The Journal of Strength & Conditioning Research* 7: 234-240, 1993.
- 4 2. Bell G, Syrotuik D, Martin T, Burnham R, and Quinney H. Effect of concurrent strength and
5 endurance training on skeletal muscle properties and hormone concentrations in humans.
6 *European Journal of Applied Physiology* 81: 418-427, 2000.
- 7 3. Bell G, Syrotuik D, Socha T, Maclean I, and Quinney HA. Effect of strength training and
8 concurrent strength and endurance training on strength, testosterone, and cortisol. *The*
9 *Journal of Strength & Conditioning Research* 11: 57-64, 1997.
- 10 4. Borg G. Perceived exertion as an indicator of somatic stress. *Scandinavian Journal of*
11 *Rehabilitation Medicine* 2: 92-98, 1970.
- 12 5. Cohen J. Statistical power analysis. *Current Directions in Psychological Science* 1: 98-101,
13 1992.
- 14 6. Craig BW, Lucas J, Pohlman R, and Stelling H. The effects of running, weightlifting and a
15 combination of both on growth hormone release. *The Journal of Strength & Conditioning*
16 *Research* 5: 198-206, 1991.
- 17 7. Day ML, McGuigan MR, Brice G, and Foster C. Monitoring exercise intensity during resistance
18 training using the session RPE scale. *The Journal of Strength & Conditioning Research* 18:
19 353-358, 2004.
- 20 8. Dempster P and Aitkens S. A new air displacement method for the determination of human
21 body composition. *Medicine & Science in Sports & Exercise* 27: 1692-1697, 1995.
- 22 9. Dudley GA and Djamil R. Incompatibility of endurance-and strength-training modes of
23 exercise. *Journal of Applied Physiology* 59: 1446-1451, 1985.
- 24 10. Dudley GA and Fleck SJ. Strength and endurance training are they mutually exclusive? *Sports*
25 *Medicine* 4: 79-85, 1987.
- 26 11. Fields DA, Goran MI, and McCrory MA. Body-composition assessment via air-displacement
27 plethysmography in adults and children: a review. *The American Journal of Clinical Nutrition*
28 75: 453-467, 2002.
- 29 12. García-Pallarés J and Izquierdo M. Strategies to optimize concurrent training of strength and
30 aerobic fitness for rowing and canoeing. *Sports Medicine* 41: 329-343, 2011.
- 31 13. Gravelle BL and Blessing DL. Physiological adaptation in women concurrently training for
32 strength and endurance. *The Journal of Strength & Conditioning Research* 14: 5-13, 2000.
- 33 14. Häkkinen K, Alen M, Kraemer W, Gorostiaga E, Izquierdo M, Rusko H, Mikkola J, Häkkinen A,
34 Valkeinen H, and Kaarakainen E. Neuromuscular adaptations during concurrent strength and
35 endurance training versus strength training. *European Journal of Applied Physiology* 89: 42-
36 52, 2003.
- 37 15. Hayes LD, Bickerstaff GF, and Baker JS. Interactions of cortisol, testosterone, and resistance
38 training: influence of circadian rhythms. *Chronobiology International* 27: 675-705, 2010.
- 39 16. Hennessy LC and Watson AWS. The interference effects of training for strength and
40 endurance simultaneously. *The Journal of Strength & Conditioning Research* 8: 12-19, 1994.
- 41 17. Hickson RC. Interference of strength development by simultaneously training for strength
42 and endurance. *European Journal of Applied Physiology and Occupational Physiology* 45:
43 255-263, 1980.
- 44 18. Hopkins WG. How to interpret changes in an athletic performance test. *Sportscience* 8: 1-7,
45 2004.
- 46 19. Izquierdo M, Häkkinen K, Ibanez J, Kraemer WJ, and Gorostiaga EM. Effects of combined
47 resistance and cardiovascular training on strength, power, muscle cross-sectional area, and
48 endurance markers in middle-aged men. *European Journal of Applied Physiology* 94: 70-75,
49 2005.

- 1 20. Jones TW, Howatson G, Russell M, and French DN. Performance and neuromuscular
2 adaptations following differing ratios of concurrent strength and endurance training. *J*
3 *Strength Cond Res* 27: 3342-3351, 2013.
- 4 21. Kraemer WJ, Gordon S, Fleck S, Marchitelli L, Mello R, Dziados J, Friedl K, Harman E, Maresh
5 C, and Fry A. Endogenous anabolic hormonal and growth factor responses to heavy
6 resistance exercise in males and females. *International Journal of Sports Medicine* 12: 228-
7 235, 2008.
- 8 22. Kraemer WJ, Patton JF, Gordon SE, Harman EA, Deschenes MR, Reynolds K, Newton RU,
9 Triplett NT, and Dziados JE. Compatibility of high-intensity strength and endurance training
10 on hormonal and skeletal muscle adaptations. *Journal of Applied Physiology* 78: 976-989,
11 1995.
- 12 23. Leveritt M and Abernethy PJ. Acute effects of high-intensity endurance exercise on
13 subsequent resistance activity. *The Journal of Strength & Conditioning Research* 13: 47-51,
14 1999.
- 15 24. McCarthy J, Agre JC, Graf B, Pozniak MA, and Vailas AC. Compatibility of adaptive responses
16 with combining strength and endurance training. *Medicine & Science in Sports & Exercise* 27:
17 429-436, 1995.
- 18 25. McCarthy J, Pozniak MA, and Agre JC. Neuromuscular adaptations to concurrent strength
19 and endurance training. *Medicine & Science in Sports & Exercise* 34: 511-519, 2002.
- 20 26. McCrory MA, Gomez TD, Bernauer EM, and Molé PA. Evaluation of a new air displacement
21 plethysmograph for measuring human body composition. *Medicine & Science in Sports &*
22 *Exercise* 27: 1686-1691, 1995.
- 23 27. Peterson MD, Rhea MR, and Alvar BA. Maximizing strength development in athletes: a meta-
24 analysis to determine the dose-response relationship. *The Journal of Strength & Conditioning*
25 *Research* 18: 377-382, 2004.
- 26 28. Peterson MD, Rhea MR, and Alvar BA. Applications of the Dose-Response for Muscular
27 Strength Development: A review of Meta-Analytic Efficacy and Reliability for Designing
28 Training Prescription. *The Journal of Strength & Conditioning Research* 19: 950-962, 2005.
- 29 29. Rønnestad BR, Hansen EA, and Raastad T. High volume of endurance training impairs
30 adaptations to 12 weeks of strength training in well-trained endurance athletes. *European*
31 *Journal of Applied Physiology* 112: 1457-1466, 2011.
- 32 30. Siri WE. Body composition from fluid spaces and density: analysis of methods, in: *Techniques*
33 *for Measuring Body Composition*. WE Siri, ed., 1961, pp 223-244.
- 34 31. Sweet TW, Foster C, McGuigan MR, and Brice G. Quantitation of resistance training using the
35 session rating of perceived exertion method. *The Journal of Strength & Conditioning*
36 *Research* 18: 796-802, 2004.
- 37 32. Volek JS, Kraemer WJ, Bush JA, Incledon T, and Boetes M. Testosterone and cortisol in
38 relationship to dietary nutrients and resistance exercise. *Journal of Applied Physiology* 82:
39 49-54, 1997.
- 40 33. Volpe SL, Walberg-Rankin J, Rodman KW, and Sebolt DR. The effect of endurance running on
41 training adaptations in women participating in a weight lifting program. *The Journal of*
42 *Strength & Conditioning Research* 7: 101-109, 1993.
- 43 34. Walshe I, Robson-Ansley P, Gibson ASC, Lawrence C, Thompson KG, and Ansley L. The
44 reliability of the IL-6, sIL-6R and sgp130 response to a preloaded time trial. *European journal*
45 *of applied physiology* 110: 619-625, 2010.

46

47

1 **ACKNOWLEDGEMENTS**

2 The authors would like to thank all individuals who volunteered to participate in the
3 study. Additional thanks go to Luke Dopson, Jordan Heath, Ashkan Hakimian, Scott
4 Keeling and Sean Armstrong for their assistance with data collection. The results of
5 the present study do not constitute any endorsement from the NSCA.

1 Figure Legends

2
3 **Figure 1.** Mean relative changes in lower body strength (as assessed by back squat and deadlift) in response to respective training interventions in the ST (n = 8), CT3 (n = 8), CT1 (n = 8) and CON (n = 6) conditions. ST, strength training alone performed every session; CT3, strength performed every session, strength and endurance training performed every third session; CT1, strength and endurance training performed every session; CON, no strength or endurance training performed during experimental period. * significant increases from baseline in all training conditions ($p < 0.05$). ** significant increase from mid-intervention in ST ($p = 0.016$). † significantly greater increases than CON in training conditions ($p < 0.05$). ‡ ST significantly greater than CT1 ($p = 0.036$).

13
14
15 **Figure 2.** Mean relative changes in upper body strength (as assessed by bench press, bent over row and military press) in response to respective training interventions in the ST (n = 8), CT3 (n = 8), CT1 (n = 8) and CON (n = 6) conditions. ST, strength training alone performed every session; CT3, strength performed every session, strength and endurance training performed every third session; CT1, strength and endurance training performed every session; CON, no strength or endurance training performed during experimental period. * significant increases from baseline in CT3 and CT1 ($p < 0.05$). ** significant increases from baseline in all training conditions ($p < 0.05$). † Significant increase from mid-intervention in ST ($p = 0.019$). ‡ all training conditions greater than CON ($p < 0.05$).

25
26
27 **Figure 3.** Mean relative changes in countermovement jump height in response to respective training interventions in the ST (n = 8), CT3 (n = 8), CT1 (n = 8) and CON (n = 6) conditions. ST, strength training alone performed every session; CT3, strength performed every session, strength and endurance training performed every third session; CT1, strength and endurance training performed every session; CON, no strength or endurance training performed during experimental period. * ST and CT1 significantly greater than baseline ($p < 0.05$). ** ST, CT3 and CT1 significantly greater than baseline ($p < 0.001$). † ST significantly greater than CT1 and CON ($p < 0.05$). ‡ ST significantly greater than CT3, CT1 and CON (all $p < 0.05$).

36
37
38 **Figure 4.** Mean RPE experienced in the ST (n = 8), CT3 (n = 8) and CT1 (n = 8) conditions. ST, strength training alone performed every session; CT3, strength performed every session, strength and endurance training performed every third session; CT1, strength and endurance training performed every session. * ST significantly lower than CT1 ($p < 0.05$).

43

1 **Table 1.** Programme variables within periodized resistance training intervention.
2

Week 1		Pre-intervention assessments	
Week 2		Familiarisation	
	Sets		3
	Repetitions		10
	% 1RM		70
	Rest (s)		90
Weeks 3 – 5		Training	
	Sets		4
	Repetitions		8
	% 1RM		80
	Rest (s)		120
Week 6		Mid-intervention assessments	
Week 7 – 9		Training	
	Sets		5
	Repetitions		6
	% 1RM		85
	Rest (s)		120
Week 10		Post-intervention assessments	
Sessions	Compound	Pull	Push
	back squat, bench press, bent over row, dead lift and military press	high pull, lat pull down, seated row, standing dumbbell reverse fly and seated hamstring curls	incline bench press, front squat, push press, seated leg press and dumbbell chest flys

1 **Table 2.** Participant's baseline maximal strength, lower body power and maximal
 2 aerobic capacity.
 3

Lower body maximal strength – 1RMs (kg)				
	ST	CT3	CT1	CON
Back squat	117.8 ± 7.7	120.3 ± 11.8	122.4 ± 8.9	118.5 ± 12.5
Deadlift	136.3 ± 7.9	142.6 ± 12.4	139.7 ± 6.7	136.9 ± 9.5
Total	254.1 ± 11.5	262.9 ± 14.2	262.1 ± 10.6	255.4 ± 11.4
Upper body maximal strength – 1RMs (kg)				
	ST	CT3	CT1	CON
Bench press	99.1 ± 9.2	105.9 ± 7.1	107.4 ± 12.4	101.6 ± 8.8
Bent over row	80.0 ± 5.3	77.5 ± 6.6	82.5 ± 5.8	80.5 ± 7.4
Military press	61.6 ± 6.1	67.5 ± 5.8	65.5 ± 7.9	60.3 ± 5.1
Total	240.6 ± 11.9	250.9 ± 12.8	255.4 ± 14.0	242.4 ± 13.6
Lower body power – CMJ (cm)				
	ST	CT3	CT1	CON
	52.7 ± 10.3	52.8 ± 7.7	50.7 ± 7.5	53.9 ± 5.1
Maximal aerobic capacity – $\dot{V}O_{2max}$ (ml·kg·min)				
	ST	CT3	CT1	CON
	52.1 ± 7.0	47.4 ± 4.9	49.5 ± 6.3	51.9 ± 7.8

4 **Note:** ST, strength training alone performed every session; CT3, strength performed
 5 every session, strength and endurance training performed every third session; CT1,
 6 strength and endurance training performed every session; CON, no strength or
 7 endurance training performed during experimental period.

1 **Table 3.** Effects of respective training interventions on testosterone, cortisol and testosterone:cortisol (T:C) ratio.

Condition	Training Session					
	Initial		Mid		Final	
	Pre	Post	Pre	Post	Pre	Post
ST						
Testosterone (nmol·L ⁻¹)	17.2 ± 4.0	23.4 ± 5.4*†	16.4 ± 2.7	23.7 ± 4.2*†	19.6 ± 10.0	27.3 ± 13.6
Cortisol (nmol·L ⁻¹)	262.6 ± 86.6	495.6 ± 150.0*†	254.4 ± 124.3	408.5 ± 145.3*	269.5 ± 116.0	389.1 ± 99.7*
T:C Ratio (x10 ³)	76.2 ± 46.3	53.5 ± 28.0	77.5 ± 36.0	63.4 ± 20.8	88.7 ± 69.7	77.67 ± 50.1
CT3						
Testosterone (nmol·L ⁻¹)	13.0 ± 1.6	17.6 ± 2.2*	15.4 ± 3.7	20.1 ± 4.2*	19.5 ± 4.6	27.1 ± 5.6*†
Cortisol (nmol·L ⁻¹)	260.5 ± 114.6	522.0 ± 325.7*	284.7 ± 103.6	460.5 ± 134.6*	262.8 ± 90.9	428.7 ± 137.0*
T:C Ratio (x10 ³)	60.7 ± 31.5	50.3 ± 42.1	58.4 ± 16.6	48.4 ± 13.6	81.0 ± 28.5	68.6 ± 22.7‡
CT1						
Testosterone (nmol·L ⁻¹)	18.7 ± 7.5	24.4 ± 11.7	19.5 ± 5.2	24.5 ± 8.0	17.3 ± 4.3	21.9 ± 4.6
Cortisol (nmol·L ⁻¹)	278.2 ± 64.9	471.6 ± 186.9*	331.4 ± 17.1	499.9 ± 48.3*†	368.3 ± 51.8	507.8 ± 45.2*† ^{AE}
T:C Ratio (x10 ³)	71.4 ± 33.9	57.6 ± 30.4	59.0 ± 15.1	49.1 ± 14.9	47.8 ± 12.5	43.0 ± 6.4
CON						
Testosterone (nmol·L ⁻¹¹)	16.1 ± 1.4	16.8 ± 1.1	16.2 ± 1.5	17.6 ± 1.4	18.5 ± 3.0	17.7 ± 2.1
Cortisol (nmol·L ⁻¹)	291.6 ± 65.0	311.5 ± 47.8	305.5 ± 91.1	320.6 ± 96.2	306.6 ± 115.8	330.2 ± 101.9
T:C Ratio (x10 ³)	58.1 ± 16.4	55.0 ± 9.0	57.0 ± 18.1	59.1 ± 17.8	58.4 ± 20.8	63.5 ± 26.0

2 * significantly greater than pre ($p < 0.05$), † significantly greater than CON ($p < 0.05$), ‡ significantly greater than post mid-session
3 ($p < 0.05$), ^{AE} increase significantly greater than ST ($p < 0.05$).


1 **Table 4.** Participant's basal lean mass.
2

Lean mass (kg)			
ST	CT3	CT1	CON
68.4 ± 6.8	66.1 ± 8.1	70.2 ± 3.7	66.9 ± 8.7

3 **Note:** ST, strength training alone performed every session; CT3, strength performed
4 every session, strength and endurance training performed every third session; CT1,
5 strength and endurance training performed every session; CON, no strength or
6 endurance training performed during experimental period

7


1 Figure 1



2

3


1 Figure 2



2

3


1 Figure 3



2

3

1 Figure 4



2