



### Scoping the application of primary care advanced clinical practice roles in England

Journal:	<i>International Journal of Health Governance</i>
Manuscript ID	ijhg-03-2020-0015.R1
Manuscript Type:	Original Article
Keywords:	Emerging healthcare delivery structures < Health economics, Primary care < Health Professions, Continuous quality improvement < Health Service Quality Assurance, Health care quality < Health Service Quality Assurance, Professional education & development < Health Service Quality Assurance, Clinical effectiveness < Health Service Quality Assurance

1  
2  
3 **Title: Scoping the application of primary care advanced clinical practice roles in**  
4 **England.**  
5

6  
7  
8 **Abstract**

9 **Purpose**

10 To scope the profile and application of advanced clinical practitioner (ACP) roles in primary  
11 care in the North of England, and how these roles meet the requirements of Health  
12 Education England's (HEE) ACP workforce capability framework.  
13  
14

15 **Design**

16 A 2 stage design was used. Stage 1 analysed health and social care workforce intelligence  
17 reports to inform scoping of numbers of ACPs working in primary care. Stage 2 used 2  
18 surveys. Survey 1 targeted ACP leads and collected strategic level data about ACP  
19 application. Survey 2 targeted staff who perceived themselves to be working as ACPs.  
20 Survey 2 was in 3 parts. Part 1 collected demographic data. Part 2 required participants to  
21 record their perceived competence against each of the HEE ACP framework capability  
22 criteria. Part 3 required respondents to identify facilitators and barriers to ACP practice.  
23  
24

25 **Results**

26 Despite the introduction of HEE's ACP capability framework, there is inconsistency and  
27 confusion about the ACP role. Results indicated a need for standardisation of role definition,  
28 and educational and practice requirements. Results also suggested that some ACPs are not  
29 working to their full potential, while some staff who are employed as 'gap-fillers' to provide  
30 routine clinical services perceive themselves as ACPs despite not working at ACP level.  
31  
32

33 **Originality/value**

34 Although previous research has explored the application of ACP practice in primary care,  
35 few studies have considered ACP application in the light of the introduction of workforce  
36 capability frameworks aimed at standardising ACP practice.  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

## Introduction

The increasing pressures on primary care services in England are well documented. Challenges include increasing demand on services to support an ageing population (European Commission, 2015; NHS England, 2019), escalating budgetary and organisational pressures (Fawdon and Adams, 2013), increased patient expectations (Williams, 2017), and staff shortages, particularly general practitioner (GP) shortages arising from struggles to recruit new partners and salaried GPs, and an increase in practising GPs leaving to work abroad (Barton *et al.*, 2012a; Peckham *et al.*, 2016). Imison *et al.*'s (2016) report for NHS Employers, the *General Practice Forward View* (NHS England, 2016), and the *NHS Long Term Plan* (NHS England, 2019) acknowledge these challenges, setting the strategic direction for an effective primary health care system with the aims of mitigating GP shortages, improving efficiency and improving patient care. Strategies include investing in the generation of 5,000 additional GPs, extending the skills of registered professionals, and developing advanced clinical practitioner (ACP) roles. The recent *Update to the GP Contract Agreement 2020/21-2023/24* (NHS England, 2020) confirms GP shortages are an ongoing challenge, and recommends further expansion of advanced multi-disciplinary team roles to release GP capacity.

Within this context, Health Education England (HEE) commissioned a study to gain insight into how ACP is specifically applied in primary care within the North of England. This paper presents the results of phase 1 of this study - scoping the profile and application of ACP in primary care.

HEE is an executive non-government departmental public body sponsored by England's Department of Health and Social Care. Its function is to provide national leadership, planning, coordination and commissioning for education and training, within the health and public health workforce in England.

## Literature review

Barton *et al.* (2012a) and Williams (2017) track the development of ACP in the United Kingdom (UK), suggesting that the origins lie with the introduction of a nurse practitioner role in primary healthcare in the late 1980s. Williams (2017) proposes evolution of ACPs in the UK is associated with medical staff shortages resulting from difficulties in the recruitment and retention of GPs. These authors note that ACPs provide a set of services that might otherwise be performed by doctors (for example, being the first contact for people with minor illness, providing routine follow-up of patients with chronic conditions, prescribing drugs or ordering tests). To a large extent, this involves a substitution of tasks from doctors to ACPs, with the main aim being to reduce demands on doctors' time, that in turn, improves access

1  
2  
3 to care, and reduces costs. Participants in Clay and Stern's (2015) study estimated that 27%  
4 of GP appointments were potentially avoidable if operational systems were transformed, for  
5 example, by using ACP consultations instead of GP consultations where appropriate.  
6  
7  
8

9  
10 A number of systematic reviews have been undertaken that investigate the effectiveness of  
11 the ACP role in primary care. Laurant *et al.* (2018) suggest ACP care improves patients'  
12 functional, health and psychological status; improves rates of patients' goal achievements,  
13 and increases levels of family-expressed satisfaction. Begley *et al.* (2013) found a clear  
14 difference between clinical specialist and advance practitioners with advanced practice roles  
15 providing improved service delivery, and greater clinical and professional leadership. Swan  
16 *et al.* (2015) found that ACPs in primary care settings perform as well as medical staff in  
17 terms of clinical outcomes and patient satisfaction, but at a lower cost. Donald *et al.* (2013)  
18 found ACP care improves the health status of older adults living in long-term care settings,  
19 and family satisfaction with care.  
20  
21  
22  
23  
24  
25

26  
27 In spite of the advantages ACP can bring to primary care, the development of the role has  
28 been largely reactionary. This has led to difficulties in defining, and further developing the  
29 role. The Royal College of Nursing (RCN) (2010) recognised that title variation hindered the  
30 public's understanding about what levels of care ACP nurses can deliver. In addition,  
31 variation in titles impedes judgements, scrutiny and understanding of practitioners'  
32 knowledge and competency to practice at an advanced level. The Department of Health's  
33 (2010) report *Advanced level nursing: A position statement* agreed, acknowledging that the  
34 terminology associated with advanced level practice had been applied inconsistently to a  
35 number of different roles, which has led to confusion about the scope and competence  
36 required at this level of practice. Surveys and studies exploring ACP and advanced nursing  
37 practice (ANP) job titles, job descriptions and levels of competency have identified  
38 considerable variation (Begley *et al.*, 2012; Elliot *et al.*, 2015; East *et al.*, 2015). These  
39 authors argue that such role inconsistency and confusion leads to inefficiencies in care,  
40 inconsistencies in levels of competency, duplication in care activities, and ineffective  
41 professional relationships.  
42  
43  
44  
45  
46  
47  
48  
49  
50

51  
52 In order to address inconsistencies in ACP roles and competence, suggestions have been  
53 made to develop a standard definition of the role, describe expected practice levels, and  
54 determine minimum educational standards, although these definitions have tended to focus  
55 on nursing practice (for example; Department of Health, 2010; Pearce and Breen, 2018;  
56 RCN, 2018). In 2017, Health Education England (HEE) published *Multi-professional*  
57 *framework for advanced clinical practice in England* (2017) - a workforce capability  
58  
59  
60

1  
2  
3 framework designed to guide development of ACP roles in a consistent way. The framework  
4 offers a standard definition of ACP that is applicable across all professions, provides clarity  
5 about the nature of expert knowledge and skills, and directs the educational requirements  
6 and governance processes required for the effective planning and development of the ACP  
7 workforce:  
8  
9

10  
11 *Advanced clinical practice is delivered by experienced, registered health and*  
12 *care practitioners. It is a level of practice characterised by a high degree of*  
13 *autonomy and complex decision making. This is underpinned by a master's level*  
14 *award or equivalent that encompasses the four pillars of clinical practice,*  
15 *leadership and management, education and research, with demonstration of core*  
16 *capabilities and area specific clinical competence.*  
17

18  
19 Despite the publication of ACP definitions and standards, some professional staff propose  
20 that consistency will be difficult to achieve without registration of the ACP role. From a  
21 nursing perspective, the RCN (2010) called for consultation a decade ago, and the recent  
22 Blake Stevenson (2019) review of standards of specialist post registration education,  
23 commissioned by the Nursing and Midwifery Council (NMC), acknowledged the need for  
24 further consultation. However, advanced practitioner level as part of the NMC and Health  
25 and Care Professions Council registers has remained unrealised. Barton *et al.* (2012b)  
26 propose advanced practitioners represent no greater public risk than new registrants,  
27 therefore, a separate part of the register would hold little benefit. Nevertheless, concerns  
28 about lack of registration remain. This has resulted in suggestions about how to regulate the  
29 role in the absence of national registration. East *et al.* (2015) suggest that UK NHS Trusts  
30 should develop registers. However, it would be difficult to include staff working outside of the  
31 NHS. The RCN (2018) has developed the notion of 'credentialing', where nurses can apply  
32 to be recognised as ACPs via an on-line application, but this is not universally recognised as  
33 a means of regulation.  
34  
35

36  
37 Although previous research has explored the application of ACP practice in primary care,  
38 few studies have considered ACP application in the light of the HEE ACP capability  
39 framework. HEE commissioned this study to gain insight into how ACP is specifically applied  
40 in primary care within the North of England. The objectives of the study were to: phase 1)  
41 scope the profile and application of ACP roles in primary care and how they meet the  
42 requirements of HEE's ACP framework; phase 2) identify any specific developments  
43 required to support ACP is to be effectively maximised 'at scale' within primary care. This  
44 paper reports on phase 1.  
45  
46

## 47 48 49 50 51 52 53 54 55 56 **Methods**

57  
58 The study location was the three HEE regions in the North of England (North West, Central  
59 North, and North East). For the study as a whole, a mixed methods approach was used. As  
60

1  
2  
3 phase 1 focused on scoping the profile and application of ACP, two approaches were used:  
4 stage 1 was an analysis of primary care services tables relating to staff numbers; stage 2  
5 used a survey approach.  
6  
7

### 8 **Stage 1 method: Analysis of service tables**

9  
10 The following public records were accessed:

- 11 • NHS Digital's GP medical service tables
- 12 • NHS service tables for nurses and direct patient care
- 13 • Skills for Care social care workforce intelligence reports

14  
15 Information from these documents relating to the 3 study locations was used to inform the  
16 scoping of the application of the ACP role in primary care.  
17

### 18 **Stage 2 method: Surveys**

#### 19 *Sample*

20  
21 Due to the potential for variability in definitions and perceptions about what constitutes  
22 advanced clinical practice, a mixture of purposive and snowball sampling was used.  
23 Purposive sampling was used to target personnel with knowledge of the scope of ACP  
24 practice within their work areas, and staff perceived by themselves or colleagues to be  
25 ACPs. Snowball sampling allowed these respondents to refer others to the study. Although  
26 this approach can have negative impacts on research in terms of selection bias, it an  
27 appropriate method of identifying participants in circumstances where the target population  
28 is not clearly defined.  
29

30  
31 *Survey data collection:* Online survey tools were developed by the research team. The tools  
32 were adapted from workforce development survey tools developed by McNall (2012), and  
33 used in a number of workforce scoping and application studies (for example, Thompson *et*  
34 *al.*, 2018 workforce caring for older people with complex needs; McNall *et al.*, 2016  
35 workforce caring for people with learning disabilities; McNall and Atkinson, 2014 primary  
36 care workforce). Before circulation, the adapted tools were piloted by two ACPs working in  
37 the study location (a clinical quality lead and an advanced nurse practitioner lead) for  
38 appropriateness of content, structure and clarity. Survey 1 was distributed via a weblink to  
39 clinical commissioning group (CCG) leaders, training hub leaders, ACP leads, directorates of  
40 nursing, allied health profession (AHP) service leads, care home and home care providers,  
41 and voluntary sector service leads working into primary care services across the HEE  
42 (North) region. The survey collected data on number, background and roles of ACPs in their  
43 region; use of the ACP framework to inform job descriptions; professional development  
44 opportunities and support provided for ACPs; barriers to developing ACP roles.  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 The invitation to participate in survey 1 included a weblink to survey 2, and survey 1  
4 participants were requested to cascade survey 2 to nurses and AHPs in their locations. In  
5 addition, permission was sought from the Queen's Nursing Institute to circulate the weblink  
6 for survey 2 via their social media and twitter sites. To maximise response rates, reminder  
7 emails were sent on a weekly basis. Survey 2 was a 3-part survey. Part 1 collected data on  
8 demographic, experience and educational backgrounds of participants; current role;  
9 professional development opportunities and support available. Part 2 required participants to  
10 record their perceived competence and confidence against each of the capability criteria  
11 within the HEE ACP framework. Responses were captured using a 4-point Likert scale: 1=  
12 not at all; 2 = not very; 3 = somewhat; 4 = very. Part 3 requested respondents to identify  
13 facilitators and barriers to ACP practice, and suggest recommendations about future role  
14 development.

15  
16  
17  
18  
19  
20  
21  
22  
23 *Data analysis:* Data from the completed surveys were imported into Statistical Package for  
24 the Social Sciences (SPSS) software, and data from the hardcopy observation tools were  
25 entered manually into SPSS in preparation for analysis.

26  
27  
28 Descriptive frequency analysis was used to provide an overview of the data. Additional  
29 analysis was carried out as follows:

- 30  
31  
32
- 33 • Survey 2 required participants to self-rate their capabilities against each individual  
34 capability of the ACP framework. Mean capability ratings were then calculated for  
35 each individual capability, and total capability for each of the four pillars.
  - 36 • Spearman rho correlation calculations were used to determine possible relationships  
37 between highest academic level and capability, and band (job grade) and capability.
- 38  
39  
40

41  
42 Only results addressing the scope and application of ACP are reported in this paper.

43  
44  
45 Research ethics approval to undertake the study was secured from the Faculty of Health and  
46 Life Sciences, xxx University.

### 47 48 49 **Results: analysis of service tables**

50  
51 NHS Digital's (2018a) GP medical services tables for nurses and direct patient care were  
52 accessed. Information from electronic staff records informs these tables. The tables showed  
53 that 1,287 GP ANPs were employed across the 3 HEE North of England regions. The basis  
54 for these numbers was labelling staff according to staff employment records i.e. job titles  
55 determined by employers.  
56  
57  
58  
59  
60

GP service tables also showed numbers of AHPs working in GP practices. The tables did not indicate how many of these were working as ACPs. In order to capture information about numbers of ACPs working in primary care, but not in GP practices, NHS Digital's (2018b) NHS services tables for nurses and direct patient care were accessed. With regard to nurses, the tables showed that across the three HEE northern regions 40 community nurse consultants and 342 community matrons were employed by the NHS. These results may indicate the number of senior nurses that may be working at ACP level in primary care outside GP practice. However, numbers for senior AHPs show the total number employed by NHS services. Information about whether these staff work in or into primary care was not provided. The Skills for Care (2018) workforce intelligence report provided detailed information on the state of the social care workforce. Numbers and demographics for registered health and social care professionals was provided, but the report did not indicate level of practice.

In summary, available information gives some indication of nurses working at ACP level in GP practices and the NHS but results rely on assumptions. Results show there is insufficient information to calculate numbers of AHPs working as ACPs, or nurses in social care working as ACPs.

### **Results: analysis of surveys**

Participants were drawn from a wide range of locations across the North of England. In total, there were 116 respondents to the surveys, 45 responding to survey 1. As survey 1 was primarily a scoping exercise to determine numbers of ACPs working throughout the North of England, service leads completed the survey, as these individuals were most likely to have access to this information. For the purposes of this study, these respondents are entitled 'ACP leads'. Survey 1 ACP lead respondents were as follows: 67% ANP leads; 6.7% directors of quality and safety (nurses); 4.4% CCG lead nurses; 4.4% GP federation managers; 4.4% education lead (nurse); 4.4% community matron leads; the remaining were service lead nurses for frailty, respiratory, dementia or workforce services. As ACP leads were commenting on the workforce, rather than on their own experiences as ACPs, it was unnecessary to present any further demographic data.

Table 1 provides details of survey 2 respondents' characteristics.

### **INSERT TABLE 1 HERE**

#### **Table 1: Survey 2 respondents**

Survey 1 asked ACP lead respondents to provide details of numbers and professional backgrounds of ACPs working in primary care in their area of practice. The valid response rate for this questions was 84.4%. Of these, 26.3% provided definitive numbers, and 73.7%



1  
2  
3 were unsure. Of respondents that were unsure, 17.9% provided explanatory comments on  
4 the survey, or emailed members of the research team to explain why these questions could  
5 not be answered fully. After analysis, 100% of these comments were coded: 'there is no  
6 registration or standard definition of an ACP, so it is not possible to ascertain numbers of  
7 staff working as ACPs'.  
8  
9  
10

11  
12 Total mean capability per pillar for all survey 2 respondents was calculated from valid  
13 responses. Valid response rates were: clinical skills capability questions 79.3%; L&M  
14 capability questions 72.4%; education capability questions 71.5%; research capability  
15 questions 71.5%. Total mean capability for clinical skills was: clinical skills 3.58 (SD 0.61);  
16 L&M 3.24 (SD 0.74); education 3.34 (SD 0.69); research 2.86 (SD 0.81). If a mean capability  
17 of 3 is taken as a minimum ACP capability level (3 = somewhat capable), then the results  
18 suggest that the ACP workforce in general has highest levels of capability in clinical skills  
19 practice, followed by education, then leadership and management. The workforce in general  
20 does not meet minimum capability levels in research.  
21  
22  
23  
24  
25

26  
27 Table 2 shows total mean capability per pillar by job. Intermediate care leads (occupational  
28 therapists), community emergency care practitioners (paramedics), and extended scope  
29 practitioners (physiotherapists) were not included as numbers were too small to provide  
30 meaningful results. Standard deviations (SDs) highlight that capability variation occurs  
31 between individual practitioners within the same job group. Results suggest that, if a mean  
32 capability of 3 is taken as a minimum ACP capability level (3= somewhat capable), then care  
33 home manager and district nurse respondents are not working at ACP level; trainee ACP  
34 respondents are only working at ACP level in clinical practice and education, and only nurse  
35 consultant respondents are working at ACP level in research. Total mean capability per pillar  
36 rankings for total participants were (from highest to lowest) clinical practice 3.58 (SD 0.61),  
37 education 3.34 (SD 0.69), leadership and management 3.24 (SD 0.74), and research 2.86  
38 (SD 0.81). This is reflected in all jobs except for care home managers, for whom rankings  
39 were (from highest to lowest) leadership and management, education, and clinical  
40 practice/research. This is to be expected because this staff group are employed specifically  
41 in a management role, are responsible for ensuring their staff are adequately trained, and  
42 delegate clinical practice to clinical lead nurses and registered nurses in their employ.  
43  
44  
45  
46  
47  
48  
49  
50  
51

52 **INSERT TABLE 2 HERE**

53 **Table 2: Mean capability per pillar by job**

54  
55  
56 Figure 1 and table 3 show total mean capability with regard to the ACP framework by band.  
57

58 **INSERT FIGURE 1 HERE**

59 **Figure 1: Total mean capability (by pillar) by band**  
60

**INSERT TABLE 3 HERE****Table 3: Mean capability per pillar by band**

Findings suggest that, if a mean capability of 3 is taken as a minimum ACP capability level (3= somewhat capable), then band 6 practitioner respondents are not working at ACP level. Only respondents of band 8b and above are working at ACP level in research. Capability level rankings for total participants are (from highest to lowest) clinical practice, education, leadership and management and research. This is reflected in all bands except for band 6, for whom rankings are (from highest to lowest) research, clinical practice, education, and leadership and management. Research capability level for band 6 is higher than band 7 and band 8a. It is unclear why this is the case.

Spearman rho correlation calculations were used to examine relationships between band and mean capability in the 4 pillars. Significant correlation was found between band and capability levels for all 4 pillars, suggesting that higher band is associated with higher capability levels:

- Clinical skills –  $\rho(54)=0.583, p<0.05$
- Leadership and management -  $\rho(54)=0.548, p<0.05$
- Education -  $\rho(54)=0.530, p=0.05$
- Research -  $\rho(54)=0.432, p<0.05$

Analysis of survey responses showed inconsistencies in minimum education qualifications required for the role, and qualifications attained by ACPs. Survey 1 asked ACP leads whether a minimum academic qualification was required for ACP roles. The valid response rate to this question was 68.9%. While 80% of respondents said yes, 20% said there was no minimum requirement. Of those that said a minimum qualification was required, 30.5% (24.4% of the total valid responses) required a Master's degree; 43.4% (34.7% of the total valid responses) a Bachelor's degree, and 4.4% (3.5% of the total valid responses) a diploma/certificate. The remaining did not specify a degree/diploma level, but required non-medical prescriber (NMP) module accreditation (13%; 10% of the total valid responses) or advanced clinical skills module accreditation (8.7%; 6.9% of the total valid responses). In survey 2, ACP participants were asked what their highest academic qualification was. The valid response rate to this question was 86.2%. Responses were: 4.7% doctorate; 41.8% Master's degree; 48.8% Bachelor's degree; 4.7% diploma/certificate.

Findings suggest that, if a mean capability of 3 is taken as a minimum ACP capability level (3= somewhat capable), then practitioner respondents with diploma/certificate education

level were not working at ACP level in clinical practice or research. Respondents with doctorates were most likely to work at ACP level in research. Capability level rankings for total participants were (from highest to lowest) clinical practice, education, leadership and management and research. This is reflected in Bachelor's and Master's degree levels. Certificate and diploma level competency rankings are leadership and management, education, clinical practice and research. One explanation is the inclusion of care home managers in certificate/diploma category. Research capability for staff educated to certificate/diploma level is higher than that for those educated to Bachelor degree level. It is unclear why this is the case. Doctorate level competency rankings are research, clinical practice, leadership and management and education. This is to be expected as staff qualified to doctoral level are likely to be research active.

Spearman rho correlation calculations were used to examine relationships between highest academic level and mean capability in the 4 pillars. These correlational comparisons showed positive relationships in all areas, and significant correlation was found between highest academic level and clinical skills, leadership and management, and research, suggesting that having high academic qualifications is associated with higher capability levels in clinical skills, leadership and management, and research:

- Clinical skills –  $\rho(53)=0.395, p<0.05$
- Leadership and management -  $\rho(53)=0.369, p<0.05$
- Education -  $\rho(53)=0.31, p=0.055$
- Research -  $\rho(46)=0.488, p<0.05$

**INSERT FIGURE 2 HERE**

**Figure 2: Mean capability (by pillar) by highest qualification**

**INSERT TABLE 4 HERE**

**Table 4: Mean capability per pillar by highest qualification**

After completing their self-reported capability scores, survey 2 asked participants to report any challenges and facilitators to practicing at ACP level for each pillar. The valid response rate to this question was 52.3%. This was an open question, and responses were analysed and coded. None of the responses identified facilitators. Identified challenges were: no requirement or opportunity to practice in the current role as other staff in the workplace carry out those roles/tasks; lack of interorganisational support of the role (for example, primary care providers requiring ACPs to make referrals to secondary care, but secondary care

providers not accepting referrals from ACPs); lack of understanding about the ACP role remit (for example, employers not understanding the advanced level of practice ACP roles are capable of, which results in restricted or limited opportunities to practice at an advanced level); demands of the routine clinical role restricting opportunities for advanced level practice and development; limited access to appropriate and relevant study programmes and courses to support ACP capability development and maintenance; lack of funding for study programmes and courses; lack of access to clinical mentorship and supervision to support ACP capability development and maintenance. Table 5 shows the percentage of valid responses per ACP practice challenge for each pillar and as a total.

### INSERT TABLE 5 HERE

#### **Table 5: Percentage of responses per ACP practice challenge for each pillar and as a total**

The first four challenges listed in the table refer to job and organisational factors relating to work allocation and interprofessional and interorganisational relationships. In total, 73% of challenges to ACP practice emanate from these. The final 3 challenges listed, totalling 27%, relate to access to education to support ACP practice development and maintenance.

### **Discussion**

Workforce intelligence documents for the NHS and social care do not record ACPs with the exception of GP ANPs. However, the recording of GP ANPs may not be reliable as some ACP lead respondents suggested as there is no registration or standard definition of ACP, it is not possible to accurately determine the scope and application of ACP, or which staff are working at ACP level. Staff from a range of job groups responded to the invitation to participate, demonstrating that either they or their employers identify them as ACPs. Results show differences in capability between job groups, and standard deviations highlight that capability variation occurs between individual practitioners within the same job group. Despite the existence of the HEE capability framework, these results show that a standard definition and agreed standards for ACP practice are not widely acknowledged or implemented, which means the title of ACP, or being identified as ACP, may offer limited insight into the capability level and practice of the practitioner. The inconsistency and confusion about ACP practice highlighted by Begley et al. (2013) and Elliot *et al.* (2016) appears to remain. This suggests that in order to achieve consistent and standardised ACP practice, regulation or registration of the role may be required because guidance in the form of definitions and capability frameworks are not universally implemented. This is in accord

1  
2  
3 with the Blake Stevenson (2019) review. However, other studies and consultation reviews  
4 propose that much more analysis is required to ensure the benefits of registration outweigh  
5 the challenges arising from costs, staff requiring multiple registrations, and the potential for  
6 causing public confusion about professional registration.  
7  
8  
9

10 Results show positive correlation between mean capability levels and band, and mean  
11 capability levels and qualifications, demonstrating that the capabilities of senior staff and  
12 staff with higher academic qualifications are more likely to align with the HEE capability  
13 framework. Staff with qualifications less than Master's degrees have a mean capability level  
14 for research that is less than 3. Staff working at lower levels than band 8a have a mean  
15 capability level for research that is less than 3. This is reflected in the HEE definition of ACP,  
16 which describes ACPs as 'experienced practitioners...characterised by a high degree of  
17 autonomy and complex decision making...underpinned by a master's level award'. This is  
18 demonstrated in the findings regarding capability by job group. The nurse consultant job  
19 group had the highest mean capability score than other job groups in all pillars, and was the  
20 only job group to have a mean capability score over 3 in research. The NHS job specification  
21 for nurse consultants requires post holders to work at band 8b level or above, and be  
22 qualified to Master's degree level or above (NHS Employers, 2020). It must be  
23 acknowledged however, that the data is collected from self-reported capability. There may  
24 be an expectation that senior, highly educated staff are more likely to practice at ACP level,  
25 and this expectation may have influenced their self-reported scores. Nevertheless, these  
26 results demonstrate that a number of staff working below the HEE ACP standards perceive  
27 themselves as, and/or are called ACPs/ANPs, or are perceived, and/or called this by their  
28 employers.  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40

41 Capability level rankings for total participants in relation to job group, band, and qualifications  
42 are (from highest to lowest) clinical practice, education, leadership and management, and  
43 research. Results regarding challenges to practice may go some way to explaining this  
44 overall ranking. While 27% of valid responses referred to difficulties in accessing education  
45 to support development or maintenance of ACP capability, 73% referred to organisational  
46 factors. A major challenge was 'no requirement to practice in the current role', particularly  
47 with regard to leadership and management, but also the other pillars; another major  
48 challenge being 'demands of the routine clinical role' which hindered practice in all pillars,  
49 but particularly in education and research. Lack of understanding about the role remit and  
50 lack of interorganisational support were also challenges. Phase 2 of this study (reported in  
51 XXX *et al.*, 2019), in which 22 practitioners were interviewed about their experiences as  
52 ACPs suggested that the absence of a standard definition of ACP, or what the remit of ACP  
53 actually is, can lead the role being used as a 'gap filler' i.e. used by employers to address  
54  
55  
56  
57  
58  
59  
60

specific local gaps due to shortages of medical staff. While, for example, this acknowledges the role of ANPs in 'freeing up GPs', as intended by the *GP Forward View*, findings from XXX *et al.*'s (2019) study suggested ANP participants often took on routine clinical activities, while leadership, education, research and complex clinical practice become the remit of GPs. This, together with the results from phase 1 reported in this paper, suggest either ACPs are not being used to their full potential, which restricts their scope of practice, or they are perceived/called ACPs although they are not actually employed to, or capable of, working at ACP level.

These explanations are reinforced by differences between the qualifications required by employers regarding ACP, and what is achieved. Results demonstrate that participants were more highly qualified than required by employers. Of note is that 20% of valid ACP lead responses did not require a minimum qualification level, while 21% of valid ACP lead responses requiring a minimum qualification (16.9% of total valid responses) required module accreditation in NMP or advanced clinical skills. These results suggest that ACPs and employers value qualifications differently. Where employers are looking for 'gap filling', they may value experience, and/or qualifications that specifically focus on clinical activities, more highly than higher academic attainment (so that ACPs can act as substitute clinicians for routine clinical activities), while ACPs themselves regard higher academic attainment as integral to ACP practice.

### Limitations

The study has a number of limitations. The small sample size recruited in one area of England limits generalisability. Snowball sampling may have introduced selection bias. Respondents to survey 2 self-assessed their competency against the HEE framework. As such expectations about competency regarding higher band staff and staff with higher qualifications may have influenced their self-reported scores.

### Conclusion

Despite the introduction of HEE's ACP capability framework, inconsistency and confusion about the role remain. Results indicate that in order to enable scoping of the application of ACP, and ensure standardisation of educational and practice requirements and consistency of ACP capability, **a standard role definition is required**. Results also suggest there is a mismatch between the HEE ACP capability framework requirements and the requirements of some employers. This can lead to ACPs not working to their full potential, or staff being employed as 'gap-fillers' to provide routine clinical services perceiving themselves and being perceived as ACPs despite not working at ACP level in some or all pillars. Standardisation of

1  
2  
3 ACP role definition and capabilities is required to distinguish ACPs to prevent  
4 inconsistencies in practice and confusion about ACP role, and to maximise the effectiveness  
5 of the role. As guidance via capability frameworks does not appear to achieve this,  
6 regulation or registration may be required, **although further analysis with regard to the**  
7 **benefits/challenges of registration is necessary.**  
8  
9  
10

## 11 **References**

12  
13 Barton, T.D., Bevan, L., and Mooney, G. (2012a), "The development of advanced nursing  
14 roles", *Nursing Times*, Vol.108 No.24, pp.18-20.

15  
16 Barton, T.D., Bevan, L., and Mooney, G. (2012b), "A governance framework for advancing  
17 nursing", *Nursing Times*, Vol.108 No.25, pp.22-24.

18  
19 Begley, C., Elliot, N., Lalor, J., Coyne, I., Higgins, A., and Comiskey, C.M. (2013),  
20 "Differences between clinical specialist and advanced practitioner clinical practice,  
21 leadership, and research roles, responsibilities, and perceived outcomes (the SCAPE  
22 study)", *Journal of Advanced Nursing*, Vol.69 No.6, pp.323–1337.

23  
24 Blake Stevenson Ltd. (2019), "Evaluation of post-registration standards of proficiency for  
25 specialist community public health nurses and the standards for specialist education and  
26 practice standards", available at:  
27 [https://www.nmc.org.uk/globalassets/sitedocuments/education-programme/evaluation-](https://www.nmc.org.uk/globalassets/sitedocuments/education-programme/evaluation-post-registration-scphn-and-spg-standards.pdf)  
28 [post-registration-scphn-and-spg-standards.pdf](https://www.nmc.org.uk/globalassets/sitedocuments/education-programme/evaluation-post-registration-scphn-and-spg-standards.pdf) (accessed 5 March 2020).

29  
30 Clay, H., and Stern, R. (2015), "Making time in general practice", available at:  
31 [https://www.primarycarefoundation.co.uk/images/PrimaryCareFoundation/Downloading\\_](https://www.primarycarefoundation.co.uk/images/PrimaryCareFoundation/Downloading_Reports/PCF_Press_Releases/Making-Time_in_General_Practice_FULL_REPORT_28_10_15.pdf)  
32 [Reports/PCF\\_Press\\_Releases/Making-Time-](https://www.primarycarefoundation.co.uk/images/PrimaryCareFoundation/Downloading_Reports/PCF_Press_Releases/Making-Time_in_General_Practice_FULL_REPORT_28_10_15.pdf)  
33 [in\\_General\\_Practice\\_FULL\\_REPORT\\_28\\_10\\_15.pdf](https://www.primarycarefoundation.co.uk/images/PrimaryCareFoundation/Downloading_Reports/PCF_Press_Releases/Making-Time_in_General_Practice_FULL_REPORT_28_10_15.pdf) (accessed 18 January 2020).

34  
35 Department of Health. (2010), *Advanced Level Nursing: A Position Statement*, Stationery  
36 Office, London.

37  
38 Donald, F., Martin-Misener, R., Carter, N., Sonald, E., Kaasalainen, S., Wickson-Griffiths,  
39 A., Lloyd, M., Akhtar-Danesh, N. and DiCenso, A. (2013), "A systematic review of the  
40 effectiveness of advanced practice nurses in long term care", *Journal of Advanced*  
41 *Nursing*, Vol.69 No.10, pp.2148–2161.

42  
43 East, L., Knowles, K. and Pettman, M. (2015), "Advanced level nursing in England:  
44 Organisation challenges and opportunities", *Journal of Nursing Management*, Vol.23 No.8,  
45 pp.1011-1019.

46  
47 Elliot, N., Begley, C., Sheaf, G. and Higgins, A. (2016), "Barriers and enablers to advanced  
48 practitioners' ability to enact their leadership role: A scoping review", *International Journal*  
49 *of Nursing Studies*, Vol.60, pp.24-45.

50  
51 European Commission. (2015), *The Ageing Report: Economic and Budgetary Projections*  
52 *for the 28 EU Member States (2013-2060)*. Brussels, European Commission.

53  
54 Fawdon, H. and Adams, J. (2013), "Advanced clinical practitioner role in the emergency  
55 department", *Nursing Standard*, Vol.28 No.16-18, pp.48-51.

56  
57 Health Education England. (2017), "Multi-professional framework for advanced clinical  
58 practice in England", available at:  
59  
60

1  
2  
3 <https://www.hee.nhs.uk/sites/default/files/documents/Multi-professional%20framework%20for%20advanced%20clinical%20practice%20in%20England.pdf> (accessed 20 October 2019).

7 Imison, C., Castle-Clarke, S., and Watson, R. (2016), "Reshaping the workforce to deliver the care patients need: Research Report", available at:  
8 <https://www.nuffieldtrust.org.uk/files/2017-01/reshaping-the-workforce-web-final.pdf>  
9 (accessed 20 October 2019)

12 Laurant. M., van-der-Biezen, M., Wijers, N., Watananirun, K., Kontopantelis, E. and van-  
13 Vught, A. (2018), "Nurses as substitutes for doctors in primary care". Cochrane Database  
14 of Systematic Reviews, Vol.2018 No.7, Art.CD001271.

16 McNall, A. (2012), *An emancipatory practice development study: Using critical discourse  
17 analysis to develop the theory and practice of sexual health workforce development*. PhD  
18 Thesis. Northumbria University, Newcastle.

20 McNall, A. and Atkinson, J. (2014), *Report of Primary Care Nursing Workforce  
21 Development Project*. Northumbria University, Newcastle.

23 McNall, A., McKenzie, K., & Branch, A. (2016), *Scoping the Workforce Development  
24 Needs of Health and Social Care Providers Delivering Positive Behavioural Support for  
25 those with Learning Disabilities across the North East and Cumbria*. Northumbria  
26 University, Newcastle.

28 NHS Digital. (2018a), "General and personal medical services, England: Final 31  
29 December 2017 and Provisional 31 March 2018, experimental statistics", available at:  
30 [https://digital.nhs.uk/data-and-information/publications/statistical/general-and-personal-  
31 medical-services/final-31-december-2017-and-provisional-31-march-2018-experimental-  
32 statistics](https://digital.nhs.uk/data-and-information/publications/statistical/general-and-personal-medical-services/final-31-december-2017-and-provisional-31-march-2018-experimental-statistics) (accessed 14 January 2019).

36 NHS Digital. (2018b), "NHS workforce statistics – September 2018", available at:  
37 [https://digital.nhs.uk/data-and-information/publications/statistical/nhs-workforce-  
38 statistics/september-2018](https://digital.nhs.uk/data-and-information/publications/statistical/nhs-workforce-statistics/september-2018) (accessed 14 January 2019).

40 NHS Employers (2020), "National job profiles". Available at:  
41 [https://www.nhsemployers.org/pay-pensions-and-reward/job-evaluation/national-job-  
42 profiles](https://www.nhsemployers.org/pay-pensions-and-reward/job-evaluation/national-job-profiles) (accessed 26 February 2020).

44 NHS England. (2016), "General practice forward view", available at:  
45 <https://www.england.nhs.uk/wp-content/uploads/2016/04/gpfv.pdf> (accessed 16 January  
46 2019).

48 NHS England. (2019), "The NHS long term plan", available at:  
49 <https://www.longtermplan.nhs.uk/wp-content/uploads/2019/01/nhs-long-term-plan.pdf>  
50 (accessed: 18 January 2019).

52 NHS England. (2020), "Update to the GP contract agreement 2020/21-2023/24", available  
53 at: [https://www.england.nhs.uk/wp-content/uploads/2020/03/update-to-the-gp-contract-  
54 agreement-v2-updated.pdf](https://www.england.nhs.uk/wp-content/uploads/2020/03/update-to-the-gp-contract-agreement-v2-updated.pdf) (accessed: 5 March 2020).

56 Pearce, C. and Breen, B. (2018), "Advanced clinical practice and nurse-led clinics: A time  
57 to progress", *British Journal of Nursing*, Vol.27 No 8, pp.444-448.



1  
2  
3 Peckham, S., Marchand, C. and Peckham, A. (2016), "General practitioner recruitment and  
4 retention: An evidence synthesis", available at:  
5 [http://blogs.lshtm.ac.uk/prucomm/files/2016/11/PRUComm-General-practitioner-  
7 recruitment-and-retention-review-Final-Report.pdf](http://blogs.lshtm.ac.uk/prucomm/files/2016/11/PRUComm-General-practitioner-<br/>6 recruitment-and-retention-review-Final-Report.pdf) (accessed 20 October 2019).

8 Royal College of Nursing. (2010). *Advanced Nurse Practitioners: A RCN Guide to the  
9 Advanced Nurse Practitioner Role, Competencies and Programme Accreditation*, RCN,  
10 London.

11  
12 Royal College of Nursing. (2018), "RCN credentialing", available at:  
13 <https://www.rcn.org.uk/professional-development/professional-services/credentialing>  
14 (accessed 12 October 2019).

15  
16 Skills for care.(2018), "Workforce intelligence: expert adult social care insight", available at:  
17 [https://www.skillsforcare.org.uk/adult-social-care-workforce-data/Workforce-  
19 intelligence/publications/regional-information/Regional-information.aspx](https://www.skillsforcare.org.uk/adult-social-care-workforce-data/Workforce-<br/>18 intelligence/publications/regional-information/Regional-information.aspx) (accessed 14  
20 January 2019).

21  
22 Swan, M., Ferguson, S., Chang, A., Larson, E. and Smaldone, A. (2015), "Quality of  
23 primary care by advanced practice nurses: A systematic review", *International Journal for  
24 Quality in Health Care*, Vol.27 No.5, pp.396–404.


25  
26 Thompson, J., Tiplady, S., McNall, A., Cook, G. and Courtney, L. (2018), *An Integrated  
27 System Based Approach to Workforce Development for Enhanced Care for Older People  
28 with Complex Needs*. Northumbria University, Newcastle.

29  
30 Williams, K. (2017), "Advanced practitioners in emergency care: A literature review",  
31 *Emergency Nurse*, Vol.25 No.4, pp.36-41.


32  
33 XXX. (2019), Paper published by the authors in the *Journal of Health Organization and  
34 Management*.

35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60



International Journal of Health Governance



International Journal of Health Governance

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

**Table 1: Survey 2 respondents**

Participant characteristics	Percentage	Mean (SD)
Gender:		
female	91.5	
male	8.5	
Age		47.1 years (SD=8.4)
Job title:		
care home manager	5.6	
GP ANP	59.2	
trainee GP ANP	8.5	
nurse consultant	7	
community matron	5.6	
district/community nurse	8.5	
intermediate care lead (occupational therapist)	1.4	
community emergency care practitioner (paramedic)	1.4	
extended scope practitioner (physiotherapist)	2.8	
Years as a qualified professional		24.5 years (SD=9.4)
Years working as an ACP since qualification		10.5 years (SD=6.2)
Band (role grade):		
6	9.9	
7	28.2	
8a	31	
8b	23.9	
>8b	7	

**Table 2: Mean capability per pillar by job**

ACP pillar	Care home manager	GP ANP	Trainee ANP	Nurse consultant	Community matron	District/ community nurse
Clinical practice: mean (SD)	2 (0)	3.78 (0.35)	3.15 (0.36)	3.94 (0.05)	3.73 (0.72)	2.39 (1.21)
Leadership & management; mean (SD)	3 (0.71)	3.39 (0.6)	2.82 (0.42)	3.87 (0.11)	3.42 (0.7)	2.33 (1.32)
Education: mean (SD)	2.5 (0)	3.46 (0.5)	3.04 (0.26)	3.88 (0.13)	3.58 (0.72)	2.29 (1.12)
Research: mean (SD)	2 (0)	2.96 (0.82)	2.67 (0.94)	3.59 (0.61)	2.92 (0.5)	2.19 (0.44)

**Table 3: Mean capability per pillar by band**

ACP pillar	Band 6	Band 7	Band 8a	Band 8b	>band 8b
Clinical practice: mean (SD)	2.6 (1.33)	3.31 (0.65)	3.71 (0.26)	3.89 (0.16)	4 (0)
Leadership & management; mean (SD)	2.3 (1.17)	3.06 (0.65)	3.2 (0.76)	3.69 (0.31)	3.91 (0.16)
Education: mean (SD)	2.33 (1.2)	3.15 (0.54)	3.3 (0.72)	3.73 (0.5)	3.96 (0.07)
Research: mean (SD)	3 (0.66)	2.39 (0.45)	2.81 (0.88)	3.38 (0.8)	3.42 (0.4)

**Table 4: Mean capability per pillar by highest qualification**

ACP pillar	ACP pillar	Diploma & certificate	Bachelors degree	Masters degree	Doctorate
Clinical practice: mean (SD)	CS Mean	3 (1.41)	3.53 (0.36)	3.64 (0.74)	3.96 (0.06)
Leadership & management; mean (SD)	LM Mean	3.2 (0.62)	3.09 (0.59)	3.35 (0.89)	3.78 (0.06)
Education: mean (SD)	Ed Mean	3.13 (0.88)	3.22 (0.51)	3.39 (0.88)	3.75 (0)
Research: mean (SD)	Res Mean	2.69 (0.98)	2.5 (0.63)	2.95 (0.82)	4 (0)

**Table 5: Percentage of responses per ACP practice challenge for each pillar and as a total**

Challenge to ACP practice	Clinical practice %	Leadership & management %	Education %	Research %	Total %
No requirement or opportunity to practice in the current role	10.5	58.8	20	13	25.5
Lack of interorganisational support of the role	10.5	16.4			6.7
Lack of understanding about the ACP role remit	15.8				3.9
Demands of the routine clinical role	21.1	23.5	46.7	56.5	36.9
Limited access to appropriate and relevant study programmes and courses	15.8		23.3		9.8
Lack of funding for study programmes and courses	10.5	11.8		30.4	13.2
Lack of access to clinical mentorship and supervision	15.8				3.9