

Northumbria Research Link

Citation: Voulgaridou, Georgia-Persephoni, Kiziridou, Magdalini, Mantso, Theodora, Chlichlia, Katerina, Galanis, Alex, Koukourakis, Michael, Franco, Rodrigo, Panagiotidis, Mihalis and Pappa, Aglaia (2016) Aldehyde dehydrogenase 3A1 promotes multi-modality resistance and alters gene expression profile in human breast adenocarcinoma MCF-7 cells. The International Journal of Biochemistry & Cell Biology, 77 (A). pp. 120-128. ISSN 1357-2725

Published by: Elsevier

URL: <http://dx.doi.org/10.1016/j.biocel.2016.06.004>
<<http://dx.doi.org/10.1016/j.biocel.2016.06.004>>

This version was downloaded from Northumbria Research Link:
<http://nrl.northumbria.ac.uk/29181/>

Northumbria University has developed Northumbria Research Link (NRL) to enable users to access the University's research output. Copyright © and moral rights for items on NRL are retained by the individual author(s) and/or other copyright owners. Single copies of full items can be reproduced, displayed or performed, and given to third parties in any format or medium for personal research or study, educational, or not-for-profit purposes without prior permission or charge, provided the authors, title and full bibliographic details are given, as well as a hyperlink and/or URL to the original metadata page. The content must not be changed in any way. Full items must not be sold commercially in any format or medium without formal permission of the copyright holder. The full policy is available online: <http://nrl.northumbria.ac.uk/policies.html>

This document may differ from the final, published version of the research and has been made available online in accordance with publisher policies. To read and/or cite from the published version of the research, please visit the publisher's website (a subscription may be required.)

www.northumbria.ac.uk/nrl



1 **Aldehyde dehydrogenase 3A1 promotes multi-modality resistance and alters**
2 **gene expression profile in human breast adenocarcinoma MCF-7 cells**

3

4 Georgia-Persephoni Voulgaridou¹, Magdalini Kiziridou¹, Theodora Mantso^{1,4},
5 Katerina Chlichlia¹, Alex Galanis¹, Michael I. Koukourakis², Rodrigo Franco³,
6 Mihalis I. Panayiotidis⁴ and Aglaia Pappa^{1*}

7

8 Departments of ¹Molecular Biology & Genetics, and ²Radiotherapy/Oncology,
9 Radiobiology & Radiopathology Unit, Department of Medicine, School of Health
10 Sciences, Democritus University of Thrace, Alexandroupolis 68100, Greece; ³Redox
11 Biology Center, School of Veterinary Medicine & Biomedical Sciences, University of
12 Nebraska-Lincoln, Lincoln, NE 68583, USA, ⁴School of Life Sciences, Heriot-Watt
13 University, Edinburgh, EH14 4AS, Scotland, UK

14

15

16 *Corresponding author:

17 Aglaia Pappa, Ph.D.
18 Department of Molecular Biology & Genetics
19 Democritus University of Thrace
20 University Campus, Dragana,
21 68100 Alexandroupolis, GREECE
22 Phone: +30-25510-30625
23 Fax: +30-25510-30625
24 E-mail: apappa@mbg.duth.gr
25

26 **Abstract**

27 Aldehyde dehydrogenases participate in a variety of cellular homeostatic mechanisms
28 like metabolism, proliferation, differentiation, apoptosis, whereas recently, they have
29 been implicated in normal and cancer cell stemness. We explored roles for ALDH3A1
30 in conferring resistance to chemotherapeutics/radiation/oxidative stress and whether
31 ectopic overexpression of ALDH3A1 could lead to alterations of gene expression
32 profile associated with cancer stem cell-like phenotype. MCF-7 cells were stably
33 transfected either with an empty vector (mock) or human aldehyde dehydrogenase
34 3A1 cDNA. The expression of aldehyde dehydrogenase 3A1 in MCF-7 cells was
35 associated with altered cell proliferation rate and enhanced cell resistance against
36 various chemotherapeutic drugs (4-hydroxyperoxycyclophosphamide, doxorubicin,
37 etoposide, and 5-fluorouracil). Aldehyde dehydrogenase 3A1 expression also led to
38 increased tolerance of MCF-7 cells to gamma radiation and hydrogen peroxide-
39 induced stress. Furthermore, aldehyde dehydrogenase 3A1-expressing MCF-7 cells
40 exhibited gene up-regulation of cyclins A, B1, B2, and down-regulation of cyclin D1
41 as well as transcription factors p21, CXR4, Notch1, SOX2, SOX4, OCT4, and JAG1.
42 When compared to mock cells, no changes were observed in mRNA levels of ABCA2
43 and ABCB1 protein pumps with only a minor decrease of the ABCG2 pump in the
44 aldehyde dehydrogenase 3A1-expressing cells. Also, the adhesion molecules EpCAM
45 and CD49F were also found to be up-regulated in aldehyde dehydrogenase
46 3A1-expressing cells. Taken together, ALDH3A1 confers a multi-modality resistance
47 phenotype in MCF-7 cells associated with slower growth rate, increased clonogenic
48 capacity, and altered gene expression profile, underlining its significance in cell
49 homeostasis.

50

51 **Keywords**

52 ALDH, ALDH3A1, MCF-7, cancer stem cells, oxidative stress, CD49F, EpCAM,
53 breast cancer, chemoresistance.

54

55 **1. Introduction**

56 Aldehyde dehydrogenase 3A1 (ALDH3A1) belongs to the broad family of aldehyde
57 dehydrogenases (ALDHs). It is an NADP⁽⁺⁾-dependent enzyme, responsible for
58 oxidizing medium chain saturated and unsaturated aldehydes to their corresponding
59 carboxylic acids (Kim et al., 2014, Vasiliou et al., 2004, Vasiliou et al., 2000).
60 Because of its ability to detoxify toxic aldehydes, by-products of lipid peroxidation
61 like 4-hydroxy-2-nonenal (4-HNE), ALDH3A1 is considered an important component
62 of cellular anti-oxidant defense (Black et al., 2012, Jang et al., 2014, Pappa et al.,
63 2003a, Pappa et al., 2003b, Voulgaridou et al., 2011). Apart from its essential
64 metabolic function, it has been suggested that ALDH3A1 may have additional roles in
65 cellular homeostasis (Kim, Lee, 2014, Voulgaridou et al., 2013) including those of
66 cell cycle regulation and protection against apoptosis and DNA damage (Chen et al.,
67 2013, Estey et al., 2007, Jang, Bruse, 2014, Lassen et al., 2007, Pappa et al., 2005,
68 Pappa et al., 2001, Stagos et al., 2010). However, ALDHs have gained even more
69 attention, after their correlation with normal and cancer stem cell (CSC) populations
70 (Gasparetto et al., 2012). In particular, the aldehyde dehydrogenase 1 (ALDH1)
71 isoform was found to be critical for the isolation of cancer cells with stem-like
72 features like self-renewal capacity, low proliferation rate, chemo-/radioresistance and
73 enhanced clonogenic and tumorigenic potential (Calderaro et al., 2014, Croker and
74 Allan, 2012, Deng et al., 2010, Lee et al., 2011, Sullivan et al., 2010, Yan et al.,
75 2014). Moreover, increased expression of ALDH was also used as an index for the
76 isolation of tumor cell subpopulations with stem-like characteristics in addition to
77 being associated with poor clinical outcome (Lee, Kim, 2011, Sullivan, Spinola,
78 2010). In this context, ALDH3A1 has been described as “tumor-associated aldehyde
79 dehydrogenase” (T-ALDH) (Lin et al., 1988) and has been shown to be upregulated in

80 several cancer types (Parajuli et al., 2014, Patel et al., 2008). Finally, only recently, it
81 has been postulated to possess additional functional roles in stem cell biology in
82 respect to self-protection, differentiation and cellular expansion (Ma and Allan, 2011).
83 However, there are not many studies suggesting how exactly the over-expression of
84 ALDH is utilized as a CSC marker and in particular what might be the underlying
85 mechanism(s) of such involvement. For these reasons, we established an isogenic
86 MCF-7 cell line pair (differing only in the expression of human ALDH3A1) with the
87 aim to (i) investigate into the effects of ALDH3A1 on cell viability and colony
88 formation efficiency under various exogenous stresses, like chemotherapeutics,
89 hydrogen peroxide (H₂O₂) and gamma-irradiation) and (ii) to identify specific gene
90 profiles attributed to such acquired CSC-like traits.

91

92 **2. Materials and Methods**

93 **2.1 Materials**

94 Human breast adenocarcinoma cell line MCF-7 was purchased from ATCC
95 (Manassas, VA, USA). All of the standard culture media, fetal bovine serum (FBS),
96 antibiotics and trypsin were either from Gibco (Life Technologies, Carlsbad, CA,
97 USA), Biosera (East Sussex, UK), Biochrome (Berlin, Germany) or Sigma-Aldrich
98 Co. (Taufkirchen, Germany). Lipofectamine and related transfection reagents were
99 obtained from Life Technologies (Carlsbad, CA, USA) while hygromycin and
100 protease inhibitors were from Carl Roth GmbH (Karlsruhe, Germany). Polyvinylidene
101 difluoride (PVDF) membranes were purchased from Millipore (Bedford, MA, USA)
102 and chemiluminescence reagents and BCA Protein assay kit were from Thermo
103 Scientific (Rockford, IL, USA). Autoradiography films were obtained from Genesee
104 Scientific (San Diego, CA, USA). All chemotherapeutic agents were from Sigma-

105 Aldrich Co. (Taufkirchen, Germany) except 4-hydroxyperoxycyclophosphamide
106 which was obtained from SantaCruz (Santa Cruz, California). Primers, dNTPs, Trizol
107 and Platinum SYBR Green, were purchased from Invitrogen (Life Technologies
108 Carlsbad, CA, USA) while random hexamers and PrimeScript Reverse Transcriptase
109 were from Takara (Shiga, Japan). Rabbit polyclonal antibody against human
110 ALDH3A1 was obtained from Abgent (San Diego, CA, USA). Mouse monoclonal
111 antibody against EpCAM was from Cell Signalling Technology (Danvers, MA,
112 USA). Goat anti-rabbit and mouse IgG horseradish peroxidase conjugated antibodies
113 were obtained from Millipore (Bedford, MA, USA). CF488A goat anti-mouse IgG for
114 immunofluorescence was from Biotium (Hayward, CA, USA). Unless stated
115 otherwise, all other chemicals were purchased from Sigma-Aldrich Co. (Taufkirchen,
116 Germany), Carl Roth GmbH (Karlsruhe, Germany) or Applichem (Darmstadt,
117 Germany).

118

119 **2.2 Cell Culture**

120 Human breast cancer cell line MCF-7 was maintained in Dulbecco's modified Eagle's
121 medium (DMEM) supplemented with 10% FBS, 100µg/ml streptomycin, and
122 100units/ml penicillin. MCF-7 stable transfected cell lines were cultured in the same
123 medium in the presence of 0.2mg/ml hygromycin. Cells were cultivated at 37°C with
124 5% CO₂ in a humidified incubator.

125

126 **2.3 Stable Transfection**

127 The full-length human ALDH3A1 was subcloned into a suitable mammalian
128 expression vector constructed as previously described (Bunting and Townsend,
129 1996a,b, Pappa, Chen, 2003a). MCF-7 cells (10^6) were transfected with 16 μ g
130 ALDH3A1/vector or control vector using the Lipofectamine 2000 reagent. Stably
131 transfected cells were selected in the presence of 0.2mg/ml of hygromycin in the
132 culture medium 48h post transfection. Selected clones were isolated, expanded and
133 maintained in the presence of hygromycin.

134

135 **2.4 Immunoblot analysis**

136 Cell lysates were prepared in 50mM Tris-HCl, pH 8.0 containing NaCl, 1% Nonidet
137 P₄₀, and the protease inhibitors: 100 μ g/ml PMSF, 0.5 μ g/ml leupeptin, 0.5 μ g/ml
138 aprotinin and 1 μ g/ml pepstatin A. Protein concentration was determined by the BCA
139 assay. Cell lysates (30 μ g of total protein) were separated by SDS-PAGE
140 electrophoresis and transferred to 0.2 μ M PVDF membranes. The blots were blocked
141 with 5% (w/v) BSA in TBST buffer (100mM Tris, pH 7.5, containing 150mM NaCl,
142 and 0.1% v/v Tween-20) (blocking buffer) for 2 hours. Primary antibodies were used
143 at different dilutions as follows: Polyclonal anti-ALDH3A1 and monoclonal anti-
144 EpCAM were used at dilutions of 1:500 and 1:5000 in blocking buffer respectively
145 (overnight, 4°C). Secondary horseradish peroxidase conjugated goat anti-rabbit and
146 mouse antibodies were used at a dilution of 1:5000 in blocking buffer (1-hour
147 incubation, RT). Signals were detected using the Supersignal West Pico
148 Chemiluminescent Substrate.

149

150 **2.5 Aldehyde dehydrogenase enzymatic activity assay**

151 The enzymatic activity of ALDH3A1 was estimated as described previously (Pappa,
152 Estey, 2003b). Briefly, a mixture of 75mM Na-pyrophosphate, pH 8.0 containing
153 1mM pyrazole, 2.5mM NADP⁺ and 50µl of cell lysates was prepared and used as a
154 blank. The reaction was initiated by the addition of 0.5mM benzaldehyde. NADPH
155 production was monitored for 5 min by the increase in the absorbance at 340nm with
156 a Biochrom Libra S22 UV/visible spectrophotometer (Biochrom, Cambridge, UK).
157 Finally, ALDH3A1 enzymatic activity was expressed as nanomoles of NADPH
158 produced per minute, per mg of protein by taking into consideration the molar
159 extinction coefficient of NADPH (6.22mM⁻¹/cm⁻¹).

160

161 **2.6 Colony Formation Assay**

162 Approximately 600 cells were plated in 10-cm culture dishes and subjected to various
163 doses (0 to 10 Gray) of gamma radiation (Cobalt 60). Subsequently, cells were placed
164 in a humidified incubator (37°C, 5% CO₂) and were monitored on a daily basis up to
165 the formation of visible colonies (usually two weeks later). Cells were then fixed and
166 stained with 0.5% of crystal violet solution diluted in 25% methanol. Colonies
167 containing ≥50 of cells were counted using a stereomicroscope and digital images
168 were obtained by camera or scanner and counted using ImageJ software.

169

170 **2.7 Sulforhodamine B (SRB) assay**

171 SRB assay was conducted as described earlier (Lassen et al. , 2006). Briefly, MCF-
172 7/mock and MCF-7/ALDH3A1 cells were seeded in 96-well culture plates and then

173 were treated, in triplicates, with 4-hydroxyperoxycyclophosphamide, etoposide,
174 doxorubicin, 5-fluorouracil, and H₂O₂. All chemotherapeutic agents were initially
175 prepared in DMSO (or water in the case 4-hydroxyperoxycyclophosphamide) (as
176 stock solutions of 50mM) and subsequently diluted (in cell culture medium) into
177 various working concentrations: 4-hydroxyperoxycyclophosphamide (0-1600μM),
178 etoposide (0-500μM), doxorubicin (0-1000μM), 5-fluorouracil (0-175μM). The
179 working concentrations of H₂O₂ were 0-1000μM, and water was used as a vehicle.
180 After a72-h incubation, cells were fixed with 50% (w/v) trichloroacetic acid (TCA)
181 for 1h at 4°C, washed 5 times with water and stained with 0.4% (w/v) SRB diluted in
182 1% acetic acid for 30 min. The excess dye was removed by washing with 1% (v/v)
183 acetic acid. Plates were dried overnight, and the protein-bound dye was dissolved in
184 10mM Tris base solution. Optical density was determined at 492nm by using a
185 microplate reader (Tecan Xflour 4). Controls were vehicle-treated cells. Sigma Plot
186 software (version 10) was used for estimating the EC₅₀ values through the regression
187 analysis via the four-parameter logistic curve as previously described (Anestopoulos
188 et al., 2013).

189

190 **2.8 Real-time PCR**

191 Total RNA was extracted using Trizol reagent according to the manufacturer's
192 instructions. For cDNA synthesis, 4.5μg of total RNA with 1 mM dNTPs and 50pmol
193 of random hexamers were used. For real-time PCR analysis, Platinum SYBR Green
194 was used according to the manufacturer's instructions. Reactions were carried out on
195 an Applied Biosystems Step One Instrument. The sequences of the primers are
196 provided in Table 1. Reactions were run in triplicate in three independent

197 experiments. Expression data were normalized to beta-actin using the $2^{-\Delta\Delta CT}$ method
198 described by Livak and Schmittgen, 2001.

199

200 **2.9 Immunofluorescence**

201 Cells (1.5×10^5) grown in a monolayer on the surface of coverslips were fixed 24-h
202 post plating with 4% formaldehyde in phosphate-buffered saline (PBS) (for 20 min)
203 and washed three times with PBS. Formaldehyde was neutralized by the addition of
204 1M of Glycine (pH 8.5). Cells were permeabilized with 0.1% Triton X-100 followed
205 by blocking with 5% BSA in PBS. The primary anti-EpCAM antibody was used at a
206 dilution of 1:800 (1h, RT) whereas the secondary (CF488A goat anti-mouse) was
207 used at a dilution of 1:250, in PBS, for 30 min. Nuclei were counterstained with 4'-6-
208 diamidino-2-phenylindole (DAPI) (1 $\mu\text{g/ml}$) and washed three times with PBS.
209 Finally, cells were mounted with MOWIOL (Calbiochem, Bad Soden, Germany) and
210 imaged with a 60x/NA 1.45 oil immersion objective and an Andor Ixon+885 digital
211 camera on a customized Andor Revolution Spinning Disk Confocal System built
212 around an IX81; Olympus stand (CIBIT Facility, MBG-DUTH). Andor IQ 2.7.1
213 software was used for image acquisition and analysis.

214

215 **2.10 Statistical analysis**

216 At least three independent experiments were conducted per sample for each
217 condition tested. All values were expressed as mean \pm S.E. Comparison of results
218 between two groups was performed by Student's t-test. Differences between

219 individual groups were assessed by a Dunnett post hoc test. Prism software (version
220 5) was used for all statistical analyses. A value of $p < 0.05$ was considered significant.

221

222 **3. Results**

223 **3.1 Generation and characterization of the MCF-7 isogenic cell line pair**

224 Stable transfection of the human ALDH3A1 cDNA in MCF-7 cells resulted in the
225 selection of two ALDH3A1/MCF-7 clones (Figure 1). Clone #2 with the highest
226 ALDH3A1 expression levels (confirmed by western blot analysis; Figure 1A) was
227 chosen for all subsequent experiments and thus designated as ALDH3A1/MCF-7.
228 Furthermore, expression of ALDH3A1 was also confirmed by real-time PCR (>100 -
229 fold in mRNA levels compared to mock/ALDH3A1 cells; Figure 1B). Enzymatic
230 activity, in ALDH3A1/MCF-7 cells, was estimated to be 535 ± 16 units/min/mg
231 whereas Mock/ALDH3A1 cells exhibited negligible activity (Figure 1C). Regular
232 monitoring of the enzymatic activity confirmed the maintenance of stable ALDH3A1
233 expression. Finally, it was observed that ALDH3A1/MCF-7 cells had considerably
234 slower cycling capacity when compared to mock ones and estimated that their colony
235 formation efficiency was approximately 57% of that of control cells (Figure 1D).

236

237 **3.1 Expression of ALDH3A1 confers chemoresistance to MCF-7 cells**

238 Next, we sought to determine the response of this isogenic cell line pair to various
239 chemotherapeutic agents characterized by different modes of actions. Mock/ and
240 ALDH3A1/MCF-7 cells were incubated for 72 h with increasing concentrations of 4-
241 hydroxyperoxycyclophosphamide (an active derivative of cyclophosphamide),
242 doxorubicin, etoposide, 5-fluorouracil and SRB-based cell viability curves were

243 plotted (Figures 2A-D respectively). Our data demonstrate that ALDH3A1 was
244 associated with a chemoresistant phenotype as indicated by the cell viability curves in
245 ALDH3A1-expressing cells compared to the non-expressing (mock) cells, under all
246 treatments. ALDH3A1/MCF-7 cells exhibited approximately 2-fold resistance to 4-
247 hydroxyperoxycyclophosphamide, (Figure 1A), ~11-fold resistance to doxorubicin
248 (Figure 2B), 8-fold resistance to etoposide (Figure 2C), and 2-fold resistance to 5-
249 fluorouracil (Figure 2D) when compared to mock cells.

250

251 **3.2 Expression of ALDH3A1 confers resistance to radiation- and H₂O₂-induced** 252 **cytotoxicity**

253 Next, we investigated on the response of the isogenic cell line pair to other cytotoxic
254 agents like H₂O₂ and gamma radiation. ALDH3A1 expression was associated with
255 increased tolerance to H₂O₂-induced cytotoxicity (Figure 3A). Interestingly, following
256 72 h incubation with a range of H₂O₂ concentrations (up to 1mM) viability in
257 ALDH3A1/MCF-7 cells did not fall below 60% when compared to control (untreated)
258 cells. On the contrary, mock/MCF-7 cells sustained roughly 10% viability under the
259 same experimental conditions (Figure 3A). Although the average EC₅₀ value for mock
260 cells was estimated around 92μM, we were unable to calculate an accurate EC₅₀ value
261 for ALDH3A1/MCF-7 cells in the same range of H₂O₂ concentrations (Figure 3A).
262 Data from colony formation collected up to two weeks post-irradiation with a range of
263 gamma irradiation (e.g. up to 10 Gy) revealed that ALDH3A1 contributed
264 significantly to the maintenance of colony formation under radiation stress (Figure
265 3B).

266

267 **3.3 ALDH3A1 alters gene expression profile in MCF-7 cells**

268 The resistant phenotype of ALDH3A1/MCF-7 cells together with the observation of
269 being slow cycling cells led to the evaluation of whether ALDH3A1 expression
270 caused any alterations in the genetic make-up of MCF-7 cells. Thus, we analyzed the
271 expression profile of several cell cycle regulatory proteins together with proteins-
272 pumps that modulate drug import/export processes in the cell. Because slow cycling
273 and chemotherapy/radiation resistance have been described as common traits for
274 cancer stem cells (Alison et al., 2011, Ghaffari, 2011), we investigated the gene
275 expression levels of those potentially relevant cancer stem cell markers including
276 CXCR4, Notch1, SOX2, Oct4, JAG1, EpCAM, and CD49f. qRT-PCR experiments
277 showed that the gene expression levels of cell cycle regulatory proteins (e.g. cyclins
278 A, B1, and B2) were up-regulated while cyclin D and p21 were down-regulated. No
279 significant changes were observed for cyclin E and p53 (Figure 4A). We also
280 examined the effects of ALDH3A1 on the expression of the ATP-binding cassette
281 (ABC) transporters ABCA2, ABCB1 (P-glycoprotein 1 or Multidrug Resistant
282 Protein 1) and ABCG2 (Breast Cancer Resistance Protein 1). ALDH3A1 expression
283 did not affect the expression levels of ABCA2 and ABCB1, whereas a slight decrease
284 was observed for ABCG2 (Figure 4B). Significant changes were observed for all
285 cancer stem cell markers tested in a manner where CXCR4, Notch1, SOX2, Oct4, and
286 JAG1 were significantly down-regulated whereas the epithelial cell adhesion
287 molecules EpCAM and CD49f (integrin subunit alpha 6) were up-regulated in
288 ALDH3A1/MCF-7 cells (Figure 4C). To further validate the RT-PCR results, we
289 selected the epithelial adhesion molecule EpCAM to confirm its up-regulation by both
290 immunofluorescence and immunoblotting. Indeed, Figure 4D depicts enhanced

291 immunofluorescent localization of EpCAM in the ALDH3A1/MCF-7 cells while
292 Western blotting also confirmed previous findings (Figure 4E).

293

294 **4. Discussion**

295 ALDHs represent a family of proteins implicated in cellular homeostasis in addition
296 to their metabolic role (Pappa, Estey, 2003b). Indeed, a variety of ALDH isoforms are
297 referred to as (i) corneal/lens crystallins (structural and protective components of
298 cornea/lens) (Estey, Piatigorsky, 2007), (ii) cell protectors against ischemia-induced
299 cardiac damage (Budasz et al., 2010, Luo et al., 2014), (iii) modulators of cell
300 proliferation rates (Lassen, Pappa, 2006, Liu et al., 2014, Pappa, Brown, 2005, Pappa,
301 Chen, 2003a, Zhang et al., 2014) and (iv) mediators of differentiation in normal and
302 cancer cells asserting to be markers of cell “stemness” (Balber, 2011, Dolle et al.,
303 2015). In particular, correlation of ALDHs with normal/cancer stem cells is not recent
304 with reports dating back to 1980s describing an association between leukemic cells
305 overexpressing ALDHs and resistance to cyclophosphamide (Russo and Hilton, 1988,
306 Tsukamoto et al., 1998). At the time and while studies were focused on the enzymatic
307 activity specificities of ALDHs (capable of detoxifying cyclophosphamide), it was
308 soon discovered that ALDHs expression was also a characteristic of healthy
309 progenitor hematopoietic cells but was gradually lost during the maturation process to
310 lymphocytes (Kastan et al., 1990). Since then, ALDHs (alone or in combination with
311 other known markers) were considered a valuable marker for isolating hematopoietic
312 progenitor populations (Armstrong et al., 2004, Fallon et al., 2003, Hess et al., 2004,
313 Storms et al., 1999). Furthermore, their usage as a putative stem cell marker was also
314 extended to the neuronal system (Balber, 2011, Cai et al., 2004, Corti et al., 2006a,
315 Corti et al., 2006b). Less than a decade ago, ALDHs were studied more extensively

316 and thus were proposed as CSC markers, initially in leukemias and later in cases of
317 solid tumours (Cheung et al., 2007, Pearce et al., 2005). Until now, ALDHs utilization
318 as CSC markers have been investigated in a broad range of different cancers and in
319 most cases, ALDHs expression was found to be a promising marker for the
320 discrimination of sub-populations with stem-like characteristics (Chen et al., 2010,
321 Deng, Yang, 2010, Emmink et al., 2011, Gong et al., 2010, Liang and Shi, 2012,
322 Marcato et al., 2011, Shien et al., 2012, Sullivan, Spinola, 2010, Wang et al., 2012).
323 On the other hand, there is still a long way to identifying specific ALDHs isoforms
324 responsible for different types of cancer in addition to determining variable potential
325 cancerous stem cell sub-population properties and qualities. Thus, elucidating the
326 underlying mechanisms of ALDHs over-expression in CSCs is of crucial importance
327 in tumor biology.

328 On another note, ALDH3A1 exhibits a distinct expression pattern. It is inducible by
329 xenobiotics in the liver and constitutively expressed in certain epithelial tissues like
330 lung, stomach, skin, and cornea. In the latter, its constitutive expression can reach up
331 to 40% of the water-soluble proteins thus classifying ALDH3A1 as a corneal
332 crystallin (Estey et al., 2010, Lassen, Bateman, 2007, Reisdorph and Lindahl, 2007).
333 In fact, ALDH3A1 is a characteristic example of the multi-functional nature of the
334 ALDH family as its expression has been associated with an apparent cell survival
335 advantage under various stress conditions thus implicating ALDH3A1 as being a
336 significant element in major homeostatic mechanisms including cell regulation and
337 apoptosis (Estey, Piatigorsky, 2007, Pappa, Chen, 2003a). To examine the putative
338 role of ALDH3A1 in the development of CSCs properties, we established MCF-7
339 cells are over-expressing ALDH3A1 and studied its impact on stem cell-like
340 properties. CSCs are relatively resistant to radiation as well as chemotherapeutic

341 agents like carboplatin, etoposide, fluorouracil, paclitaxel, daunorubicin,
342 mitoxantrone, cyclophosphamide, temozolomide, and gemcitabine (Dylla et al., 2008,
343 Hermann et al., 2007, Liu et al., 2006, Ma et al., 2008, Todaro et al., 2007, Wulf et al.,
344 2001). Interestingly, our results indicated that ALDH3A1 protects MCF-7 cells from
345 the cytotoxic effects of a wide variety of commonly used chemotherapeutic agents
346 like 4-hydroxyperoxycyclophosphamide, etoposide, doxorubicin and 5-fluorouracil
347 (Horak et al., 2013, Lekakis et al., 2012, Loi et al., 2013, Moitra et al., 2012). Indeed,
348 previous studies have documented up-regulation of ALDHs with enhanced
349 chemoresistance in breast cancer cells both in vitro and in vivo (Cioce et al., 2014,
350 Croker and Allan, 2012, Lee, Kim, 2011). The results are in accordance with previous
351 studies that have shown that overexpression of ALDH3A1 results in resistance to 4-
352 hydroxyperoxycyclophosphamide and other active metabolites of cyclophosphamide
353 [Bunting et al. J Biol Chem. 1994, 269: 23197-23203, Moreb et al., 2007).
354 Interestingly, increased resistance to doxorubicin has also been associated with the
355 ectopic expression of other ALDH members (Moreb et al., Chem. Biol. Interact.,
356 2012, 195: 52-60), which is possibly mediated through indirect mechanisms by
357 modulating oxidative stress response as previously reported for ALDH3A1 in
358 relation to resistance to mitomycin C and etoposide (Pappa A et al., J. Biol. Chem.
359 2005, 280: 27998–28006). Moreover, ALDH3A1/MCF-7 cells exhibited enhanced
360 survival and colony formation capacities in the presence of additional stress factors
361 like gamma radiation and exposure to H₂O₂. Certainly, the specificity of ALDH3A1
362 for the metabolism and detoxification of cyclophosphamide (Bunting and Townsend,
363 1996b) and 4-HNE (Pappa, Estey, 2003b) is an important contributing factor
364 underlining resistance, but its ability to protect adequately against a variety of other
365 stressors supports the notion for an overall, multi-mode resistance phenotype

366 characteristic of ALDH3A1/MCF-7 cells. One possible mechanism accountable for
367 the apparent resistance of these cells would be their slow-growing rate. This is in
368 accordance with another study where ALDH3A1 led to inhibition of proliferation,
369 slower cell cycling rates, and lower colony formation efficiency expression in human
370 corneal epithelial cells (Estey, Piatigorsky, 2007, Pappa, Brown, 2005). This anti-
371 proliferative action of ALDH3A1 was also observed in our study where the
372 ALDH3A1/MCF-7 cells had the capacity to form only about 57% of the colonies
373 formed in mock/MCF-7 cells. In general, CSCs are slow-growing cells in the
374 quiescent state and consequently resistant to drugs designed to target fast-growing
375 cancer cells (Dalerba et al., 2007, Tirino et al., 2013, Vinogradov and Wei, 2012). To
376 characterize the molecular mechanisms responsible for the slow proliferation rates
377 observed, we analyzed the gene expression profile of key cell cycle regulatory
378 proteins. We noticed that ALDH3A1-expressing MCF-7 cells exhibited an (i) up-
379 regulation of cyclins A, B1, B2 and (ii) down-regulation of cyclin D1 and
380 transcription factor p21. Previous studies demonstrated that protein levels of cyclins
381 A, B, E, E2F1, and p21, as well activities of cyclin A- and cyclin B- dependent
382 kinases were all decreased in ALDH3A1/HCE cells (Pappa, Brown, 2005). While it is
383 true that the comparative qPCR method used in this study detects differences only at
384 the transcriptional level, the differential expression pattern of major cell cycle
385 regulatory proteins (also previously reported for ALDH3A1-expressing HCE cells)
386 may account for the slow proliferation phenotype observed. On the other hand, there
387 are also reports associating knock down of ALDH3A1 in lung cancer cells with slower
388 growth (Moreb et al., 2008). To this end, findings so far appear contradictory, and
389 although they may reflect tissue-specific issues or differences in biology between

390 normal and cancer cells, they urge the need for further investigations towards the
391 clarification of the role of ALDH3A1 in cell proliferation.

392 The possibility that drug resistance displayed by the ALDH3A1-expressing cells is
393 likely due to enhanced expression of transporters that mediate chemotherapeutic drug
394 efflux (Gottesman et al., 2002, Ween et al., 2015) was excluded. In general, several
395 types of ABC transporters are known to be over-expressed in a variety of cancers
396 where they are responsible for the development of chemoresistance (Chang et al.,
397 2009, Doyle and Ross, 2003, Gottesman, Fojo, 2002, Mack et al., 2008). However, no
398 detectable changes were observed in mRNA levels of ABCA2 and ABCB1 protein
399 pumps. On the contrary, only a minor decrease observed for the ABCG2 pump in the
400 ALDH3A1-expressing cells compared to mock. Another possible mechanism for the
401 observed chemo-/radioresistance, in the presence of ALDH3A1, would be through
402 mediating DNA damage checkpoint response. Indeed, increased activation of the
403 DNA damage checkpoint response has been associated with expression of ALDH3A1
404 in corneal epithelial cells and preliminary data (obtained in our lab) certainly points
405 towards this direction (data not shown). Similarly, the resistance of glioblastoma
406 CSCs to irradiation has been attributed to increased activation of the DNA damage
407 checkpoint (Bao et al., 2006).

408 To better characterize the changes caused by ALDH3A1 on gene expression, we
409 investigated the presence of presumed protein markers found to be up-regulated in
410 CSCs. The gene expression profile was significantly differentiated between the two
411 MCF-7 isogenic cell lines. The mRNA levels of CXCR4, Notch1, SOX2, SOX4,
412 OCT4, and JAG1, displayed down-regulation whereas EpCAM and CD49F were
413 significantly up-regulated in the ALDH3A1/MCF-7 cells. We further validated the
414 expression of the epithelial cell adhesion molecule (EpCAM) by immunofluorescence

415 and immunoblotting and showed that EpCAM protein levels were substantially
416 elevated in the ALDH3A1 expressing MCF-7 cells. EpCAM together with CD49F
417 have been studied extensively for their functional roles and usage as potential CSCs
418 markers (Cariati et al., 2008, Deng et al., 2015, Guo et al., 2012, Guo et al., 2014,
419 Wang et al., 2011). EpCAM is suggested to provide a sustained proliferative signal to
420 cancer-initiating and normal stem cells where it is overexpressed. Cancer cells appear
421 to benefit from the constitutive expression of EpCAM for proliferation, self-renewal,
422 and anchorage-independent growth and invasiveness (Munz et al., 2009). On the other
423 hand, CD49F (also known as $\alpha 6$ integrin) plays a significant role in cell adhesion. Its
424 high expression in mammary epithelial cells is associated with progenitor and stem
425 cell activity (Goel et al., 2014). This integrin acts as an adhesion receptor for the
426 mammary epithelial cells mediating developmental signals and assisting cells in
427 sensing growth factor and hormonal signals (Kaimala et al., 2012). It appears to play a
428 major role in sustaining the survival of mammary carcinoma cells especially under
429 stress conditions such as those existing in the tumor microenvironment (Chung and
430 Mercurio, 2004).

431 In conclusion, MCF-7 cells over-expressing ALDH3A1 demonstrated low
432 proliferation rates associated with a resistant phenotype against various sources of cell
433 stress including exposure to various chemotherapeutics, gamma radiation, and H₂O₂
434 insult. Furthermore, they displayed differential expression of proteins involved in cell
435 cycle regulation and increased expression of the cell adhesion molecules CD49f and
436 EpCAM. Although the precise mechanisms remain unclear, our findings provide
437 considerable implications on defining the biological significance of ALDH3A1 in cell
438 homeostasis.

439

440 **Acknowledgments**

441 This research has been co-financed by the European Union (European Social Fund –
442 ESF) and Greek national funds through the Operational Program "Education and
443 Lifelong Learning" of the National Strategic Reference Framework (NSRF) -
444 Research Funding Program: Heracleitus II. Investing in knowledge society through
445 the European Social Fund.

446

447

448

449

450 **FIGURE LEGENDS**

451 **Figure 1: Characterization of the MCF-7 isogenic cell line pair. A.** Western blot
452 analysis of ALDH3A1 expression: Lane 1: recombinant ALDH3A1 (1 µg), lanes 2-7:
453 30 µg cell extracts, 2; parental MCF-7, 3-4; mock-transfected ALDH3A1, 6-7:
454 ALDH3A1/MCF-7 transfected clones. **B.** ALDH3A1 gene expression levels detected
455 by real-time PCR in mock/ and ALDH3A1/MCF7. **C.** Enzymatic activity of
456 ALDH3A1 in mock/MCF-7 and ALDH3A1/MCF-7 cells. Results are expressed as
457 means of a minimum of three independent experiments ± SE. **D.** Colony formation
458 efficiency of mock and ALDH3A1/MCF-7 cells. Cells (600) were seeded in 10 cm
459 culture dishes and were allowed to form colonies for two weeks in a humidified
460 incubator that were subsequently counted following crystal violet staining by using
461 Image J. Results are expressed as mean ±S.E of three independent experiments. ***
462 p<0.001.

463 **Figure 2: Effect of various chemotherapeutic agents on cell viability of mock/
464 and ALDH3A1/MCF-7 cells.**

465 Viability curves of mock/ and ALDH3A1/MCF-7 cells along with the calculated half
466 maximal effective concentrations (EC₅₀ values) of **(A)** 4-
467 hydroxyperoxycyclophosphamide, **(B)** doxorubicin, **(C)** etoposide, and **(D)** 5-
468 fluorouracil are represented. Viability curves of the ALDH3A1/MCF-7 cells are
469 shifted to the right indicating increased tolerance of the cells to the cytotoxic effect of
470 the agents used. Results are shown as mean ± S.E. At least three independent
471 experiments were performed for each condition. * p<0.05, ** p<0.01, *** p<0.001

472

473 **Figure 3: Effect of H₂O₂ and gamma radiation on the viability of mock/ and**
474 **ALDH3A1/MCF-7 cells.** The viability curves of mock/ and ALDH3A1/MCF7 cells
475 along with the half maximal effective concentrations (EC₅₀ values) of (A) H₂O₂ and
476 (B) gamma radiation are presented. ALDH3A1 expression is associated with
477 increased tolerance to the cytotoxic effects of H₂O₂ and gamma radiation. Results are
478 presented as mean ± SE of three independent experiments. * p<0.05, ** p<0.01, ***
479 p<0.001

480

481 **Figure 4: Expression of ALDH3A1 alters gene profiling in ALDH3A1/MCF-7**
482 **cells.** Effect of ALDH3A1 on the gene expression of (A) cycle cell regulatory
483 proteins (B) Membrane ABC transporters (C) Cancer stem cell markers. The
484 comparative quantification ΔΔCt method was utilized for analyzing the fold change of
485 gene expression. Beta-actin gene was used as endogenous control for the
486 normalization of samples. **D:** Immunofluorescence for EpCAM (green) in
487 ALDH3A1/MCF-7 (i) and mock/MCF-7 (ii) cells. No secondary antibody for
488 EpCAM was used in the negative control (iii), whereas nuclei were stained with DAPI
489 (4',6-diamino-2-phenylindole) (blue). **E.** Western blotting analysis for EpCAM in
490 mock and ALDH3A1/MCF-7 cells. Results are shown as mean ± S.E. At least three
491 independent experiments were performed for each condition. * p<0.05, ** p<0.01,
492 *** p<0.001

493

494
495
496

Table 1. Primers used for the real-time PCR comparative quantification

GENE	FORWARD PRIMER	REVERSE PRIMER
β-actin	GCGCGGCTACAGCTTCA	CTTAATGTCACGCACGATTTCC
ALDH3A1	CAGCGGCATGGGATCCTA	GCGGCGGTGAGAGAAAGTC
Cyclin A	ACGGGTTGCACCCCTTAAG	CCAAGGAGGAACGGTGACA
Cyclin B1	GGCCTCTACCTTTGCACTTCCT	GCTCGACATCAACCTCTCCAA
Cyclin B2	AAGCTTTTTCTGATGCCTTGCT	AGGGTTCTCCAATCTTCGTTAT
Cyclin D	AGACCTTCGTTGCCTCTTGTG	ATGGAGGGCGGATTGGAA
Cyclin E	GGCCTTGTATCATTCTCGTCAT	CGCACCCTGATACCCTGAA
p53	TCTGTCCCTTCCCAGAAAACC	CAAGAAGCCCAGACGGAAAC
p21	GGCGGGCTGCATCCA	AGTGGTGTCTCGGTGACAAAGTC
ABCA2	AGATGGACAAGATGATCGAG	GCTTGTACTTCAGGATGAGG
ABCB1	GAGGAAGACATGACCAGGTA	CTGTTCGATTATAGCATGAA
ABCG2	ACCTGAAGGCATTTACTGAA	TCTTTCCTTGCAGCTAAGAC
CXCR4	GGCCGACCTCCTCTTTGTC	TTGCCACGGCATCAACTG
Notch1	GCACCTCAGCCTGCACAGT	CTGTGTTGCTGGAGCATCTTCT

SOX2	TGCGAGCGCTGCACAT	TCATGAGCGTCTTGGTTTTCC
SOX4	CTGCGCCTCAAGCACATG	TTCTTCCTGGGCCGGTACT
Oct4	CGACCATCTGCCGCTTTG	GCCGCAGCTTACACATGTTCT
JAG1	TGAAGTAGAAGAGGACGACATGGA	CGGCTGCTTGGCAAACC
EpCAM	TTATGATCCTGACTGCGATGAGA	GGTGCCGTTGCACTGCTT
CD49F	GATCCCGGCCTGTGATTAATATT	CTGGCGGAGGTCAATTCTGT

497

498

499

500

501

502

503

504

505

506

507

508

509

510

511

512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565

References

- Alison MR, Lim SM, Nicholson LJ. Cancer stem cells: problems for therapy? *J Pathol.* 2011;223:147-61.
- Anestopoulos I, Kavou A, Tentis I, Kortsaris A, Panayiotidis M, Lazou A, et al. Silibinin protects H9c2 cardiac cells from oxidative stress and inhibits phenylephrine-induced hypertrophy: potential mechanisms. *J Nutr Biochem.* 2013;24:586-94.
- Armstrong L, Stojkovic M, Dimmick I, Ahmad S, Stojkovic P, Hole N, et al. Phenotypic characterization of murine primitive hematopoietic progenitor cells isolated on basis of aldehyde dehydrogenase activity. *Stem Cells.* 2004;22:1142-51.
- Balber AE. Concise review: aldehyde dehydrogenase bright stem and progenitor cell populations from normal tissues: characteristics, activities, and emerging uses in regenerative medicine. *Stem Cells.* 2011;29:570-5.
- Bao S, Wu Q, McLendon RE, Hao Y, Shi Q, Hjelmeland AB, et al. Glioma stem cells promote radioresistance by preferential activation of the DNA damage response. *Nature.* 2006;444:756-60.
- Black W, Chen Y, Matsumoto A, Thompson DC, Lassen N, Pappa A, et al. Molecular mechanisms of ALDH3A1-mediated cellular protection against 4-hydroxy-2-nonenal. *Free Radic Biol Med.* 2012;52:1937-44.
- Budas GR, Disatnik MH, Chen CH, Mochly-Rosen D. Activation of aldehyde dehydrogenase 2 (ALDH2) confers cardioprotection in protein kinase C epsilon (PKC ϵ) knockout mice. *J Mol Cell Cardiol.* 2010;48:757-64.
- Bunting KD, Townsend AJ. De novo expression of transfected human class 1 aldehyde dehydrogenase (ALDH) causes resistance to oxazaphosphorine anti-cancer alkylating agents in hamster V79 cell lines. Elevated class 1 ALDH activity is closely correlated with reduction in DNA interstrand cross-linking and lethality. *J Biol Chem.* 1996a;271:11884-90.
- Bunting KD, Townsend AJ. Protection by transfected rat or human class 3 aldehyde dehydrogenase against the cytotoxic effects of oxazaphosphorine alkylating agents in hamster V79 cell lines. Demonstration of aldophosphamide metabolism by the human cytosolic class 3 isozyme. *J Biol Chem.* 1996b;271:11891-6.
- Cai J, Cheng A, Luo Y, Lu C, Mattson MP, Rao MS, et al. Membrane properties of rat embryonic multipotent neural stem cells. *J Neurochem.* 2004;88:212-26.
- Calderaro J, Nault JC, Bioulac-Sage P, Laurent A, Blanc JF, Decaens T, et al. ALDH3A1 is overexpressed in a subset of hepatocellular carcinoma characterised by activation of the Wnt/ss-catenin pathway. *Virchows Arch.* 2014;464:53-60.
- Cariati M, Naderi A, Brown JP, Smalley MJ, Pinder SE, Caldas C, et al. Alpha-6 integrin is necessary for the tumorigenicity of a stem cell-like subpopulation within the MCF7 breast cancer cell line. *Int J Cancer.* 2008;122:298-304.
- Chang H, Rha SY, Jeung HC, Im CK, Ahn JB, Kwon WS, et al. Association of the ABCB1 gene polymorphisms 2677G>T/A and 3435C>T with clinical outcomes of paclitaxel monotherapy in metastatic breast cancer patients. *Ann Oncol.* 2009;20:272-7.
- Chen Y, Thompson DC, Koppaka V, Jester JV, Vasiliou V. Ocular aldehyde dehydrogenases: protection against ultraviolet damage and maintenance of transparency for vision. *Prog Retin Eye Res.* 2013;33:28-39.
- Chen YW, Chen KH, Huang PI, Chen YC, Chiou GY, Lo WL, et al. Cucurbitacin I suppressed stem-like property and enhanced radiation-induced apoptosis in head and neck squamous carcinoma--derived CD44(+)/ALDH1(+) cells. *Mol Cancer Ther.* 2010;9:2879-92.
- Cheung AM, Wan TS, Leung JC, Chan LY, Huang H, Kwong YL, et al. Aldehyde dehydrogenase activity in leukemic blasts defines a subgroup of acute myeloid leukemia with adverse prognosis and superior NOD/SCID engrafting potential. *Leukemia.* 2007;21:1423-30.
- Chung J, Mercurio AM. Contributions of the alpha6 integrins to breast carcinoma survival and progression. *Mol Cells.* 2004;17:203-9.
- Cioce M, Valerio M, Casadei L, Pulito C, Sacconi A, Mori F, et al. Metformin-induced metabolic reprogramming of chemoresistant ALDHbright breast cancer cells. *Oncotarget.* 2014;5:4129-43.

566 Corti S, Locatelli F, Papadimitriou D, Donadoni C, Del Bo R, Crimi M, et al. Transplanted
567 ALDHhiSSClo neural stem cells generate motor neurons and delay disease progression of
568 nmd mice, an animal model of SMARD1. *Hum Mol Genet.* 2006a;15:167-87.

569 Corti S, Locatelli F, Papadimitriou D, Donadoni C, Salani S, Del Bo R, et al. Identification of
570 a primitive brain-derived neural stem cell population based on aldehyde dehydrogenase
571 activity. *Stem Cells.* 2006b;24:975-85.

572 Croker AK, Allan AL. Inhibition of aldehyde dehydrogenase (ALDH) activity reduces
573 chemotherapy and radiation resistance of stem-like ALDHhiCD44(+) human breast cancer
574 cells. *Breast Cancer Res Treat.* 2012;133:75-87.

575 Dalerba P, Cho RW, Clarke MF. Cancer stem cells: models and concepts. *Annu Rev Med.*
576 2007;58:267-84.

577 Deng S, Yang X, Lassus H, Liang S, Kaur S, Ye Q, et al. Distinct expression levels and
578 patterns of stem cell marker, aldehyde dehydrogenase isoform 1 (ALDH1), in human
579 epithelial cancers. *PLoS One.* 2010;5:e10277.

580 Deng Z, Wu Y, Ma W, Zhang S, Zhang YQ. Adoptive T-cell therapy of prostate cancer
581 targeting the cancer stem cell antigen EpCAM. *BMC Immunol.* 2015;16:1.

582 Dolle L, Boulter L, Leclercq IA, van Grunsven LA. Next generation of ALDH substrates and
583 their potential to study maturational lineage biology in stem and progenitor cells. *Am J*
584 *Physiol Gastrointest Liver Physiol.* 2015;308:G573-8.

585 Doyle L, Ross DD. Multidrug resistance mediated by the breast cancer resistance protein
586 BCRP (ABCG2). *Oncogene.* 2003;22:7340-58.

587 Dylla SJ, Beviglia L, Park IK, Chartier C, Raval J, Ngan L, et al. Colorectal cancer stem cells
588 are enriched in xenogeneic tumors following chemotherapy. *PLoS One.* 2008;3:e2428.

589 Emmink BL, Van Houdt WJ, Vries RG, Hoogwater FJ, Govaert KM, Verheem A, et al.
590 Differentiated human colorectal cancer cells protect tumor-initiating cells from irinotecan.
591 *Gastroenterology.* 2011;141:269-78.

592 Estey T, Chen Y, Carpenter JF, Vasiliou V. Structural and functional modifications of corneal
593 crystallin ALDH3A1 by UVB light. *PLoS One.* 2010;5:e15218.

594 Estey T, Piatigorsky J, Lassen N, Vasiliou V. ALDH3A1: a corneal crystallin with diverse
595 functions. *Exp Eye Res.* 2007;84:3-12.

596 Fallon P, Gentry T, Balber AE, Boulware D, Janssen WE, Smilee R, et al. Mobilized
597 peripheral blood SSCloALDHbr cells have the phenotypic and functional properties of
598 primitive haematopoietic cells and their number correlates with engraftment following
599 autologous transplantation. *Br J Haematol.* 2003;122:99-108.

600 Gasparetto M, Sekulovic S, Brocker C, Tang P, Zakaryan A, Xiang P, et al. Aldehyde
601 dehydrogenases are regulators of hematopoietic stem cell numbers and B-cell development.
602 *Exp Hematol.* 2012;40:318-29 e2.

603 Ghaffari S. Cancer, stem cells and cancer stem cells: old ideas, new developments. *F1000*
604 *Med Rep.* 2011;3:23.

605 Goel HL, Gritsko T, Pursell B, Chang C, Shultz LD, Greiner DL, et al. Regulated splicing of
606 the alpha6 integrin cytoplasmic domain determines the fate of breast cancer stem cells. *Cell*
607 *Rep.* 2014;7:747-61.

608 Gong C, Yao H, Liu Q, Chen J, Shi J, Su F, et al. Markers of tumor-initiating cells predict
609 chemoresistance in breast cancer. *PLoS One.* 2010;5:e15630.

610 Gottesman MM, Fojo T, Bates SE. Multidrug resistance in cancer: role of ATP-dependent
611 transporters. *Nat Rev Cancer.* 2002;2:48-58.

612 Guo C, Liu H, Zhang BH, Cadaneanu RM, Mayle AM, Garraway IP. Epcam, CD44, and
613 CD49f distinguish sphere-forming human prostate basal cells from a subpopulation with
614 predominant tubule initiation capability. *PLoS One.* 2012;7:e34219.

615 Guo Z, Li LQ, Jiang JH, Ou C, Zeng LX, Xiang BD. Cancer stem cell markers correlate with
616 early recurrence and survival in hepatocellular carcinoma. *World J Gastroenterol.*
617 2014;20:2098-106.

618 Hermann PC, Huber SL, Herrler T, Aicher A, Ellwart JW, Guba M, et al. Distinct populations
619 of cancer stem cells determine tumor growth and metastatic activity in human pancreatic
620 cancer. *Cell Stem Cell.* 2007;1:313-23.

621 Hess DA, Meyerrose TE, Wirthlin L, Craft TP, Herrbrich PE, Creer MH, et al. Functional
622 characterization of highly purified human hematopoietic repopulating cells isolated according
623 to aldehyde dehydrogenase activity. *Blood*. 2004;104:1648-55.

624 Horak CE, Pusztai L, Xing G, Trifan OC, Saura C, Tseng LM, et al. Biomarker analysis of
625 neoadjuvant doxorubicin/cyclophosphamide followed by ixabepilone or Paclitaxel in early-
626 stage breast cancer. *Clin Cancer Res*. 2013;19:1587-95.

627 Jang JH, Bruse S, Liu Y, Duffy V, Zhang C, Oyamada N, et al. Aldehyde dehydrogenase 3A1
628 protects airway epithelial cells from cigarette smoke-induced DNA damage and cytotoxicity.
629 *Free Radic Biol Med*. 2014;68:80-6.

630 Kaimala S, Bisana S, Kumar S. Mammary gland stem cells: more puzzles than explanations. *J*
631 *Biosci*. 2012;37:349-58.

632 Kastan MB, Schlaffer E, Russo JE, Colvin OM, Civin CI, Hilton J. Direct demonstration of
633 elevated aldehyde dehydrogenase in human hematopoietic progenitor cells. *Blood*.
634 1990;75:1947-50.

635 Kim SW, Lee J, Lee B, Rhim T. Proteomic analysis in pterygium; upregulated protein
636 expression of ALDH3A1, PDIA3, and PRDX2. *Mol Vis*. 2014;20:1192-202.

637 Lassen N, Bateman JB, Estey T, Kuszak JR, Nees DW, Piatigorsky J, et al. Multiple and
638 additive functions of ALDH3A1 and ALDH1A1: cataract phenotype and ocular oxidative
639 damage in *Aldh3a1(-)/Aldh1a1(-)* knock-out mice. *J Biol Chem*. 2007;282:25668-76.

640 Lassen N, Pappa A, Black WJ, Jester JV, Day BJ, Min E, et al. Antioxidant function of
641 corneal ALDH3A1 in cultured stromal fibroblasts. *Free Radic Biol Med*. 2006;41:1459-69.

642 Lee HE, Kim JH, Kim YJ, Choi SY, Kim SW, Kang E, et al. An increase in cancer stem cell
643 population after primary systemic therapy is a poor prognostic factor in breast cancer. *Br J*
644 *Cancer*. 2011;104:1730-8.

645 Lekakis L, Tryfonopoulos D, Pistamatzian N, Panopoulos C, Koumakis G, Demiri S, et al.
646 Salvage chemotherapy with cisplatin and 5-fluorouracil in metastatic breast cancer. Particular
647 activity against liver metastases. *Anticancer Res*. 2012;32:1833-7.

648 Liang D, Shi Y. Aldehyde dehydrogenase-1 is a specific marker for stem cells in human lung
649 adenocarcinoma. *Med Oncol*. 2012;29:633-9.

650 Lin KH, Brennan MD, Lindahl R. Expression of tumor-associated aldehyde dehydrogenase
651 gene in rat hepatoma cell lines. *Cancer Res*. 1988;48:7009-12.

652 Liu G, Yuan X, Zeng Z, Tunici P, Ng H, Abdulkadir IR, et al. Analysis of gene expression
653 and chemoresistance of CD133+ cancer stem cells in glioblastoma. *Mol Cancer*. 2006;5:67.

654 Liu Y, Jiang X, Zeng Y, Zhou H, Yang J, Cao R. Proliferating pancreatic beta-cells
655 upregulate ALDH. *Histochem Cell Biol*. 2014;142:685-91.

656 Livak KJ, Schmittgen TD. Analysis of relative gene expression data using real-time
657 quantitative PCR and the 2(-Delta Delta C(T)) Method. *Methods*. 2001;25:402-8.

658 Loi S, Sirtaine N, Piette F, Salgado R, Viale G, Van Eenoo F, et al. Prognostic and predictive
659 value of tumor-infiltrating lymphocytes in a phase III randomized adjuvant breast cancer trial
660 in node-positive breast cancer comparing the addition of docetaxel to doxorubicin with
661 doxorubicin-based chemotherapy: BIG 02-98. *J Clin Oncol*. 2013;31:860-7.

662 Luo XJ, Liu B, Ma QL, Peng J. Mitochondrial aldehyde dehydrogenase, a potential drug
663 target for protection of heart and brain from ischemia/reperfusion injury. *Curr Drug Targets*.
664 2014;15:948-55.

665 Ma I, Allan AL. The role of human aldehyde dehydrogenase in normal and cancer stem cells.
666 *Stem Cell Rev*. 2011;7:292-306.

667 Ma S, Lee TK, Zheng BJ, Chan KW, Guan XY. CD133+ HCC cancer stem cells confer
668 chemoresistance by preferential expression of the Akt/PKB survival pathway. *Oncogene*.
669 2008;27:1749-58.

670 Mack JT, Brown CB, Tew KD. ABCA2 as a therapeutic target in cancer and nervous system
671 disorders. *Expert Opin Ther Targets*. 2008;12:491-504.

672 Marcato P, Dean CA, Pan D, Araslanova R, Gillis M, Joshi M, et al. Aldehyde dehydrogenase
673 activity of breast cancer stem cells is primarily due to isoform ALDH1A3 and its expression
674 is predictive of metastasis. *Stem Cells*. 2011;29:32-45.

675 Moitra K, Im K, Limpert K, Borsa A, Sawitzke J, Robey R, et al. Differential gene and
676 microRNA expression between etoposide resistant and etoposide sensitive MCF7 breast
677 cancer cell lines. *PLoS One*. 2012;7:e45268.

678 Moreb JS, Baker HV, Chang LJ, Amaya M, Lopez MC, Ostmark B, Chou W. ALDH
679 isozymes downregulation affects cell growth, cell motility and gene expression in lung cancer
680 cells. *Mol Cancer*. 2008;7:87. doi: 10.1186/1476-4598-7-87.

681 Moreb JS, Mohuczy D, Ostmark B, Zucali JR. RNAi-mediated knockdown of aldehyde
682 dehydrogenase class-1A1 and class-3A1 is specific and reveals that each contributes equally
683 to the resistance against 4-hydroperoxycyclophosphamide. *Cancer Chemother Pharmacol*.
684 2007;59:127-36.

685 Moreb JS, Ukar D, Moreb JS, Ucar D, Han S, Amory JK, Goldstein AS, Ostmark B, Chang
686 LJ. The enzymatic activity of human aldehyde dehydrogenases 1A2 and 2 (ALDH1A2 and
687 ALDH2) is detected by Aldefluor, inhibited by diethylaminobenzaldehyde and has significant
688 effects on cell proliferation and drug resistance. *Chem Biol Interact*. 2012;195:52-60.

689 Munz M, Baeuerle PA, Gires O. The emerging role of EpCAM in cancer and stem cell
690 signaling. *Cancer Res*. 2009;69:5627-9.

691 Pappa A, Brown D, Koutalos Y, DeGregori J, White C, Vasiliou V. Human aldehyde
692 dehydrogenase 3A1 inhibits proliferation and promotes survival of human corneal epithelial
693 cells. *J Biol Chem*. 2005;280:27998-8006.

694 Pappa A, Chen C, Koutalos Y, Townsend AJ, Vasiliou V. Aldh3a1 protects human corneal
695 epithelial cells from ultraviolet- and 4-hydroxy-2-nonenal-induced oxidative damage. *Free
696 Radic Biol Med*. 2003a;34:1178-89.

697 Pappa A, Estey T, Manzer R, Brown D, Vasiliou V. Human aldehyde dehydrogenase 3A1
698 (ALDH3A1): biochemical characterization and immunohistochemical localization in the
699 cornea. *Biochem J*. 2003b;376:615-23.

700 Pappa A, Sophos NA, Vasiliou V. Corneal and stomach expression of aldehyde
701 dehydrogenases: from fish to mammals. *Chem Biol Interact*. 2001;130-132:181-91.

702 Parajuli B, Georgiadis TM, Fishel ML, Hurley TD. Development of selective inhibitors for
703 human aldehyde dehydrogenase 3A1 (ALDH3A1) for the enhancement of cyclophosphamide
704 cytotoxicity. *Chembiochem*. 2014;15:701-12.

705 Patel M, Lu L, Zander DS, Sreerama L, Coco D, Moreb JS. ALDH1A1 and ALDH3A1
706 expression in lung cancers: correlation with histologic type and potential precursors. *Lung
707 Cancer*. 2008;59:340-9.

708 Pearce DJ, Taussig D, Simpson C, Allen K, Rohatiner AZ, Lister TA, et al. Characterization
709 of cells with a high aldehyde dehydrogenase activity from cord blood and acute myeloid
710 leukemia samples. *Stem Cells*. 2005;23:752-60.

711 Reisdorph R, Lindahl R. Constitutive and 3-methylcholanthrene-induced rat ALDH3A1
712 expression is mediated by multiple xenobiotic response elements. *Drug Metab Dispos*.
713 2007;35:386-93.

714 Russo JE, Hilton J. Characterization of cytosolic aldehyde dehydrogenase from
715 cyclophosphamide resistant L1210 cells. *Cancer Res*. 1988;48:2963-8.

716 Shien K, Toyooka S, Ichimura K, Soh J, Furukawa M, Maki Y, et al. Prognostic impact of
717 cancer stem cell-related markers in non-small cell lung cancer patients treated with induction
718 chemoradiotherapy. *Lung Cancer*. 2012;77:162-7.

719 Stagos D, Chen Y, Cantore M, Jester JV, Vasiliou V. Corneal aldehyde dehydrogenases:
720 multiple functions and novel nuclear localization. *Brain Res Bull*. 2010;81:211-8.

721 Storms RW, Trujillo AP, Springer JB, Shah L, Colvin OM, Ludeman SM, et al. Isolation of
722 primitive human hematopoietic progenitors on the basis of aldehyde dehydrogenase activity.
723 *Proc Natl Acad Sci U S A*. 1999;96:9118-23.

724 Sullivan JP, Spinola M, Dodge M, Raso MG, Behrens C, Gao B, et al. Aldehyde
725 dehydrogenase activity selects for lung adenocarcinoma stem cells dependent on notch
726 signaling. *Cancer Res*. 2010;70:9937-48.

727 Tirino V, Desiderio V, Paino F, De Rosa A, Papaccio F, La Noce M, et al. Cancer stem cells
728 in solid tumors: an overview and new approaches for their isolation and characterization.
729 *FASEB J*. 2013;27:13-24.

730 Todaro M, Alea MP, Di Stefano AB, Cammareri P, Vermeulen L, Iovino F, et al. Colon
731 cancer stem cells dictate tumor growth and resist cell death by production of interleukin-4.
732 *Cell Stem Cell.* 2007;1:389-402.

733 Tsukamoto N, Chen J, Yoshida A. Enhanced expressions of glucose-6-phosphate
734 dehydrogenase and cytosolic aldehyde dehydrogenase and elevation of reduced glutathione
735 level in cyclophosphamide-resistant human leukemia cells. *Blood Cells Mol Dis.*
736 1998;24:231-8.

737 Vasiliou V, Pappa A, Estey T. Role of human aldehyde dehydrogenases in endobiotic and
738 xenobiotic metabolism. *Drug Metab Rev.* 2004;36:279-99.

739 Vasiliou V, Pappa A, Petersen DR. Role of aldehyde dehydrogenases in endogenous and
740 xenobiotic metabolism. *Chem Biol Interact.* 2000;129:1-19.

741 Vinogradov S, Wei X. Cancer stem cells and drug resistance: the potential of nanomedicine.
742 *Nanomedicine (Lond).* 2012;7:597-615.

743 Voulgaridou GP, Anestopoulos I, Franco R, Panayiotidis MI, Pappa A. DNA damage induced
744 by endogenous aldehydes: current state of knowledge. *Mutat Res.* 2011;711:13-27.

745 Voulgaridou GP, Mantso T, Chlichlia K, Panayiotidis MI, Pappa A. Efficient *E. coli*
746 expression strategies for production of soluble human crystallin ALDH3A1. *PLoS One.*
747 2013;8:e56582.

748 Wang Y, Shenouda S, Baranwal S, Rathinam R, Jain P, Bao L, et al. Integrin subunits alpha5
749 and alpha6 regulate cell cycle by modulating the chk1 and Rb/E2F pathways to affect breast
750 cancer metastasis. *Mol Cancer.* 2011;10:84.

751 Wang YC, Yo YT, Lee HY, Liao YP, Chao TK, Su PH, et al. ALDH1-bright epithelial
752 ovarian cancer cells are associated with CD44 expression, drug resistance, and poor clinical
753 outcome. *Am J Pathol.* 2012;180:1159-69.





754 Ween MP, Armstrong MA, Oehler MK, Ricciardelli C. The role of ABC transporters in
755 ovarian cancer progression and chemoresistance. *Crit Rev Oncol Hematol.* 2015.


756 Wulf GG, Wang RY, Kuehnle I, Weidner D, Marini F, Brenner MK, et al. A leukemic stem
757 cell with intrinsic drug efflux capacity in acute myeloid leukemia. *Blood.* 2001;98:1166-73.


758 Yan J, De Melo J, Cutz JC, Aziz T, Tang D. Aldehyde dehydrogenase 3A1 associates with
759 prostate tumorigenesis. *Br J Cancer.* 2014;110:2593-603.

760 Zhang L, Wang L, Liu X, Zheng D, Liu S, Liu C. ALDH expression characterizes G1-phase
761 proliferating beta cells during pregnancy. *PLoS One.* 2014;9:e96204.

762







Figure(s)
[Click here to download high resolution image](#)








Table 1. Primers used for the real-time PCR comparative quantification

GENE	FORWARD PRIMER	REVERSE PRIMER
<i>β-actin</i>	GCGCGGCTACAGCTTCA	CTTAATGTCACGCACGATTTCC
<i>ALDH3A1</i>	CAGCGGCATGGGATCCTA	GCGGCGGTGAGAGAAAGTC
<i>Cyclin A</i>	ACGGGTTGCACCCCTTAAG	CCAAGGAGGAACGGTGACA
<i>Cyclin B1</i>	GGCCTCTACCTTTGCACTTCT	GCTCGACATCAACCTCTCCAA
<i>Cyclin B2</i>	AAGCTTTTTCTGATGCCTTGCT	AGGGTTCTCCAATCTTCGTTAT
<i>Cyclin D</i>	AGACCTTCGTTGCCTCTTG	ATGGAGGGCGGATTGGAA
<i>Cyclin E</i>	GGCCTTGTATCATTCTCGTCAT	CGCACCCTGATACCCTGAA
<i>p53</i>	TCTGTCCCTTCCCAGAAAACC	CAAGAAGCCCAGACGGAAAC
<i>p21</i>	GGCGGGCTGCATCCA	AGTGGTGTCTCGGTGACAAAGTC
<i>ABCA2</i>	AGATGGACAAGATGATCGAG	GCTTGTACTTCAGGATGAGG
<i>ABCB1</i>	GAGGAAGACATGACCAGGTA	CTGTGCATTATAGCATGAA
<i>ABCG2</i>	ACCTGAAGGCATTTACTGAA	TCTTCCTTGCAGCTAAGAC
<i>CXCR4</i>	GGCCGACCTCCTTTTGTC	TTGCCACGGCATCAACTG
<i>Notch1</i>	GCACCTCAGCCTGCACAGT	CTGTGTTGCTGGAGCATCTTCT

<i>SOX2</i>	TGCGAGCGCTGCACAT	TCATGAGCGTCTTGGTTTTCC
<i>SOX4</i>	CTGCGCCTCAAGCACATG	TTCTTCCTGGGCCGGTACT
<i>Oct4</i>	CGACCATCTGCCGCTTTG	GCCGCAGCTTACACATGTTCT
<i>JAG1</i>	TGAAGTAGAAGAGGACGACATGGA	CGGCTGCTTGGCAAACC
<i>EpCAM</i>	TTATGATCCTGACTGCGATGAGA	GGTGCCGTTGCACTGCTT
<i>CD49F</i>	GATCCCGCCTGTGATTAATATT	CTGGCGGAGGTCAATTCTGT