

# Northumbria Research Link

Citation: Graham, Kirsty, Sparagano, Olivier and Finn, Robert (2016) Isolation of the monooxygenase complex from *Rhipicephalus (Boophilus) microplus* – clues to understanding acaricide resistance. *Ticks and Tick-borne Diseases*, 7 (4). pp. 614-623. ISSN 1877-959X

Published by: Elsevier

URL: <http://dx.doi.org/10.1016/j.ttbdis.2016.01.012>  
<<http://dx.doi.org/10.1016/j.ttbdis.2016.01.012>>

This version was downloaded from Northumbria Research Link:  
<http://nrl.northumbria.ac.uk/25926/>

Northumbria University has developed Northumbria Research Link (NRL) to enable users to access the University's research output. Copyright © and moral rights for items on NRL are retained by the individual author(s) and/or other copyright owners. Single copies of full items can be reproduced, displayed or performed, and given to third parties in any format or medium for personal research or study, educational, or not-for-profit purposes without prior permission or charge, provided the authors, title and full bibliographic details are given, as well as a hyperlink and/or URL to the original metadata page. The content must not be changed in any way. Full items must not be sold commercially in any format or medium without formal permission of the copyright holder. The full policy is available online: <http://nrl.northumbria.ac.uk/policies.html>

This document may differ from the final, published version of the research and has been made available online in accordance with publisher policies. To read and/or cite from the published version of the research, please visit the publisher's website (a subscription may be required.)

[www.northumbria.ac.uk/nrl](http://www.northumbria.ac.uk/nrl)



Isolation of the monooxygenase complex from *Rhipicephalus (Boophilus) microplus* – clues  
to understanding acaricide resistance

Kirsty M Graham<sup>a</sup>, Olivier AE Sparagano<sup>b</sup>, Robert D Finn<sup>a c</sup>

<sup>a</sup> Department of Applied Sciences, Faculty of Health and Life Sciences, Northumbria University Newcastle, Ellison Building, Ellison Place, Newcastle Upon Tyne, NE1 8ST, UK.

<sup>b</sup> Coventry University, Alan Berry Building, Coventry University, Priory Street, Coventry, CV1 5FB, UK.

[kirsty.graham@northumbria.ac.uk](mailto:kirsty.graham@northumbria.ac.uk)

[olivier.sparagano@coventry.ac.uk](mailto:olivier.sparagano@coventry.ac.uk)

[robert.finn@northumbria.ac.uk](mailto:robert.finn@northumbria.ac.uk)

<sup>c</sup> Corresponding author: Dr. Robert Finn, Department of Applied Sciences, Faculty of Health and Life Sciences, Northumbria University Newcastle, Ellison Building, Ellison Place, Newcastle Upon Tyne, NE1 8ST, UK. Tel: +44 191 2274554, email:

[robert.finn@northumbria.ac.uk](mailto:robert.finn@northumbria.ac.uk)

## Abstract

The monooxygenase complex is composed of three key proteins, a cytochrome P450 (CYP), the cytochrome P450 oxidoreductase (CPR) and cytochrome  $b_5$  and plays a key role in the metabolism and detoxification of xenobiotic substances, including pesticides. In addition, overexpression of these components has been linked to pesticide resistance in several important vectors of disease. Despite this, the monooxygenase complex has not been isolated from the Southern cattle tick *Rhipicephalus (Boophilus) microplus*, a major disease vector in livestock.

Using bioinformatics 115 transcriptomic sequences were analyzed to identify putative pesticide metabolizing CYPs. RACE-PCR was used to amplify the full length sequence of one CYP; CYP3006G8 which displays a high degree of homology to members of the CYP6 and 9 subfamilies, known to metabolize pyrethroids. mRNA expression levels of CYP3006G8 were investigated in 11 strains of *R. microplus* with differing resistance profiles by qPCR, the results of which indicated a correlation with pyrethroid metabolic resistance. In addition to this gene, the sequences for CPR and cytochrome  $b_5$  were also identified and subsequently isolated from *R. microplus* using PCR.

CYP3006G8 is only the third CYP gene isolated from *R. microplus* and the first to putatively metabolize pesticides. The initial results of expression analysis suggest that CYP3006G8 metabolizes pyrethroids but further biochemical characterization is required to confirm this. Differences in the kinetic parameters of human and mosquito CPR in terms of NADPH binding have been demonstrated and could potentially be used to design species specific pesticides. Similar differences in the tick CPR would confirm that this is a characteristic of hematophagous arthropods.

**Keywords:** Cytochrome P450 (CYP); cytochrome P450 oxidoreductase (CPR); cytochrome *b<sub>5</sub>*; *Rhipicephalus microplus*; pesticide resistance.

## Introduction

The detoxification of xenobiotics is an essential function for life and is carried out by three specific enzyme families: Glutathione S transferases, Esterases (or Carboxylesterases) and the Cytochrome P450s (CYPs). Cytochrome P450s are an ancient protein family found in all forms of life (Scott and Wen 2001; Sztal et al., 2012) and are known to be the major phase I metabolizing enzymes. In humans, CYPs metabolize approximately 65% of currently used clinical drugs (Zhou et al., 2005), emphasising the importance of this group of enzymes in xenobiotic metabolism. CYPs are characterized by the presence of a highly conserved heme-binding domain located towards the C terminus of the protein. This signature motif, with the sequence FxxGxxxCxG is found in all CYPs (Syed and Mashele 2014; Werck-Reichhart and Feyereisen 2000) and plays a vital role in the electron transport mechanism used by these enzymes to, in many cases, incorporate molecular oxygen into xenobiotics thus rendering them more water soluble and more readily excreted from the cell (Mamidala et al., 2011).

Xenobiotic metabolism is an important topic of investigation amongst arthropods due to this group of organisms containing numerous vectors of medical/veterinary diseases and a host of agricultural pests that are developing resistance to commonly used chemical control agents (Brogdon and McAllister 1998). Amongst the arthropods, there has been a huge amount of research into species such as *Musca domestica* (house fly) and *Drosophila melanogaster* (fruit fly) due to their use as model organisms and of course various mosquito

species due to the medical and economic importance (Drali et al., 2012; Kawada et al., 2011; Kushwah et al., 2015; Wondji et al., 2011).

Pesticide resistance is not a novel problem but one that is ever increasing due to its association with disease transmission to both animals and plants. Pesticide resistance has been described by the World Health Organisation (WHO) as “the ability of a parasite strain to survive and/or multiply despite the administration and absorption of a drug given in doses equal to or higher than those usually recommended but within the limits of tolerance of the subject” Abbas et al., (2014). It is this ability to survive despite the application of pesticides that is of interest as it provides a mechanism to continue the transmission of diseases to both plants and animals. One of the most studied of the arthropods with a view of disease transmission are the mosquito species and research has shown that CYPs play a key role in the development of pesticide resistance (Corbel et al., 2007; Gong et al., 2005; Hemingway et al., 2004; Marcombe et al., 2009). Cytochrome P450s are split into clades depending on their function and this allows key CYPs to be more easily identified. Four main clades (mitochondrial clade, CYP2, CYP3 and CYP4 clade) exist and in terms of pesticide metabolism the clade that is of interest is the CYP3 clade, a clade which includes the important mammalian CYP3A4 and insect homologues of this enzyme, namely the CYP6 and CYP9 families e.g. CYP6Z1 (Chiu et al., 2008) and CYP6P3 (Muller et al., 2008).

Despite the importance of ticks, as they are second to mosquitoes as vectors of human disease and the most important vector of disease to animals (Hill and Wikel, 2005), with the ability to transmit a variety of pathogens and transmitting the widest array of disease causing organisms out of all the hematophagous arthropods (Bissinger and Roe 2010), little research has been carried out into the cytochrome P450 mediated metabolism and their subsequent associated resistance in this group. The vast majority of pesticide resistance research in these arthropods has focused on the role of GSTs (Kwon et al., 2010) and

carboxylesterases (Baxter and Barker 1998; Cossio-Bayugar et al., 2002), with the involvement of CYPs only being inferred by synergist studies (Li et al., 2004; Rodriguez-Vivas et al., 2013). Furthermore, despite the ever increasing problem of pesticide resistance little of this research has involved deciphering the mechanisms underlying the development of resistance phenotypes. The identification of CYPs, and indeed other drug metabolizing enzymes, in acari is hindered by the lack of available annotated genome data. Some data is available on public databases such as NCBI for *Ixodes scapularis* and the *I. scapularis* genome has been published (Van Zee et al., 2007) however for species such as *R. microplus*, the data available is even more sparse (Guerrero et al., 2006). It is therefore important to develop these resources such that a bioinformatics approach can be initiated to identify members of the key CYP families and where possible, identify putative homologues of the key insect CYPs that have been isolated and characterized to ascertain what they metabolize.

In order to be fully functional cytochrome P450s require an oxidoreductase (CPR in this case) and additionally in some cases cytochrome *b*<sub>5</sub>. CPR is important in transferring electrons from NADPH through to the CYP allowing the incorporation of one molecule of oxygen, hence a monooxygenase reaction, to various substrates (Guengerich 2007). CPR is a highly conserved diflavin protein composed of two flavin cofactors, flavin mononucleotide (FMN) and flavin adenine dinucleotide (FAD), that transfer the electrons one at a time to a CYP (Iyanagi et al., 2012; Lian et al., 2011). Due to the nature of CYP research among mammals and insects, the CPR gene has been isolated from an array of species and been characterized in many of these (Chen and Zhang 2014; Liu et al., 2014; Zhu et al., 2012) however, CPR has not been isolated or characterized from a tick species to date, despite the importance of this protein to the functioning of the monooxygenase complex and subsequent detoxification of xenobiotics. Differences have been identified

between human CPR and *An. gambiae* CPR which is potentially paving the way for a new target and method of control (Lian et al., 2011), but further emphasises the importance of characterising CPR isoforms from other arthropods.

Cytochrome  $b_5$  is a small but highly conserved protein with a variety of roles in different reactions such as lipid and sterol biosynthesis (McLaughlin et al., 2010; Porter 2002; Zhang et al., 2005). The role cytochrome  $b_5$  plays within the monooxygenase complex and CYP mediated reactions has been described as controversial for many years (Porter 2005) as cytochrome  $b_5$  has been shown to have a stimulatory/modifying effect on CYP reactions, be it obligatory for some reactions and in other cases have an inhibitory effect (Im and Waskell 2011; Porter 2002; Schenkman and Jansson 2003). Despite this extensive amount of research into the role of cytochrome  $b_5$  in CYP-mediated drug metabolism in mammals, much less equivalent research exists in arthropods considering the fact cytochrome  $b_5$  may have a role in pesticide metabolism. Stevenson et al., (2012) found that by co-expressing numerous CYPs such as CYP9J2 from *Ae. aegypti* with cytochrome  $b_5$  from *An. gambiae* the catalytic activity of the CYP was enhanced. Conversely, Chandor-Proust (2013) found that co-expression of *Ae. aegypti* cytochrome  $b_5$  with *Ae. aegypti* CYP6Z8 did not cause a significant effect on the activity of the CYP (Chandor-Proust et al., 2013). This emphasises the need to co-express cytochrome  $b_5$  when investigating CYP activity. Thus it is reasonable to assume that further investigation into the role of cytochrome  $b_5$  in humans and some mosquito species will identify potential differences between species that similar to CPR, may additionally be exploited in the development of arthropod control measures.

In this paper we provide an update of our recent work towards filling the gap in the current knowledge of cytochrome P450s involved in pesticide metabolism in acari describing the isolation of the P450 monooxygenase complex from the Southern cattle tick, *R. microplus*.

Isolation of these individual components will allow detailed biochemical characterization and the ascertainment of their role in pesticide metabolism/resistance.

## **Materials & Methods**

### *Bioinformatic analysis*

115 putative cytochrome P450 sequences from *Rhipicephalus (Boophilus) microplus* provided by Dr Felix Guerrero and colleagues (Guerrero et al., 2006) were initially analyzed using the National Center for Biotechnology Information (NCBI) public database (<http://www.ncbi.nlm.nih.gov/>) to identify those having nucleotide sequence similarity to CYP3, CYP6 and/or CYP9. Of the 115 initial sequences 5 were identified as having the highest 'query coverage' and 'max ident' to known pesticide metabolizing CYPs from other species. Within these 115 were sequences were *R. microplus* homologues of cytochrome P450 oxidoreductase (CPR) and cytochrome *b*<sub>5</sub> (cytb<sub>5</sub>) genes were also identified.

### *Tick samples*

Tick larval samples from the species *R. microplus*, were provided from USDA, Texas (courtesy of Dr Felix Guerrero) preserved in RNA<sup>later</sup>-Ice® (Life Technologies, New York, USA), and stored at -80 °C to maintain RNA stability and integrity.

### *RNA Extraction and cDNA preparation*

RNA was extracted from all 11 strains of *R. microplus* shown in Table 1 (the resistance profile of each strain is also included). The ticks were snap frozen using liquid nitrogen and RNA was extracted using TRIzol® reagent followed by a PureLink RNA Mini Kit (Life Technologies, New York, USA) according the manufacturer's instructions. RNA quantity



and purity was measured using a Nanodrop ND-100 (Nanodrop Technologies, Wilmington, USA) and integrity was checked using agarose gel electrophoresis.

cDNA was generated from the 11 strains of *R. microplus* ticks using PrimerDesign Reverse Transcription Premix (PrimerDesign, Southampton UK).

### *RACE Reactions*

The Gonzales strain of *R. microplus* was used to generate RACE (Rapid Amplification of cDNA Ends) ready cDNA using a GeneRacer Kit (Life Technologies, New York, USA) and 1.44 µg of RNA was used to generate both 5' and 3' RACE ready cDNA. RACE PCRs were carried out using the RACE ready cDNA as follows:

5' RACE reactions contained, 300 µM dNTPs, 1 Unit VELOCITY DNA polymerase, 10 µl 5x amplification buffer, 900 nM GeneRacer 5' Primer (5' GGACTGACATGGACTGAAGGAGTA 3') and 300 nM GSP Reverse primer (5' GGGCGAAGCCAGACGACACC 3'), 1 mM MgCl<sub>2</sub>, 30µl sterile distilled water, 1 µl RACE ready cDNA template in a final reaction volume of 50 µl.

3' RACE reactions contained, 300 µM dNTPs, 1 Unit VELOCITY DNA polymerase, 10 µl 5x amplification buffer, 900 nM GeneRacer 3' Primer (5' GCTGTCAACGATACGCTACGTAACG 3') and 300 nM GSP Forward primer (10 µM) (5' TCCAGGTCCCCACCAAGTTT 3'), 1 mM MgCl<sub>2</sub>, 30µl sterile distilled water, 1 µl RACE ready cDNA template in a final reaction volume of 50 µl.

5' and 3' RACE fragments were generated using the following PCR cycling conditions: 1 cycle of 98 °C for 2 minutes, followed by 5 cycles of 98 °C for 30 seconds, 72 °C for 2 minutes, followed by 5 cycles of 98 °C for 30 seconds, 70 °C for 2 minutes, followed by 30 cycles of 98 °C for 30 seconds, 56 °C for 30 seconds, 72 °C for 2 minutes with a final

extension of 72 °C for 10 minutes. RACE products were double strand sequenced in order to facilitate the generation of gene-specific primers for amplification of the full length cDNA.

#### *PCR amplification of full length CYP3006G8*

PCR reactions were set up using VELOCITY DNA Polymerase (Bioline, London, UK) as follows; 0.5 µl of cDNA (per 12.5 µl reaction), 10 µl 5 x Hi-Fi Reaction Buffer, 400 µM dNTPs, 10 µM of each forward (5' CCGCACTCGCTCGAATCAT 3') and reverse (5' GCTGCTGTAGCAAACGATGTC 3') primers, 1 mM MgCl<sub>2</sub>, 2 Units of polymerase enzyme in a total of 50 µl. Reactions were subjected to the following PCR cycling conditions. 1 cycle of 98 °C followed by 35 cycles of 98 °C for 30 seconds, 52 °C for 30 seconds and 72 °C for 1 minute with a final step of 72 °C for 10 minutes. Following PCR, products were analyzed on a 1 % agarose gel. The full length cDNA was cloned into pCR Blunt (Life Technologies, Paisley, UK) and the sequence confirmed via double strand sequencing using the vector specific M13 forward and reverse primers.

#### *qPCR mRNA expression*

cDNA was generated from the 11 strains of *R. microplus* ticks. This was carried out using Reverse Transcription Premix (PrimerDesign, Southampton, UK).

The full length CYP3006G8 was used to design a primer/probe set to carry out quantitative expression analysis: forward 5' TTTCCTTACTGCTTGCTTTCC 3', reverse, 5' CGTCAAACACTTGTCTGCC 3', probe 5' FAM - TCATATCATTTCGGTTCACGACATAACAAGACG – TAM. Additionally a primer/probe set was designed to ELF1α which was the housekeeping gene used to normalise the data (Nijhof et al., 2009): forward, 5' CTTACCGACTTTCTCCCC 3', reverse 5' GCCTCCTTGGGTTTGACAGA 3', probe, 5' FAM - AACGGTCGCCGTCGGTGTC – TAM

3'. Following optimization of primer/probe sets, for CYP3006G8 200 nM forward primer, 100 nM reverse primer and 300 nM probe were used in each reaction and for ELF1 $\alpha$  100 nM forward, 200 nM reverse and 300 nM probe was used. All reactions were set up as follows, 6.25  $\mu$ l 2 x qPCR Master Mix for the iCycler (Primer Design, Southampton, UK), 2.5  $\mu$ l RNase free water, primer/probe at the appropriate concentration indicated above and 5 ng/ $\mu$ l of cDNA. Reactions were then put through the following cycling conditions, 95 °C for 10 minutes followed by 40 cycles of 95 °C for 15 seconds, 60 °C for one minute, data collection, using a BioRad CFX96 real-time PCR machine. The Ct values were converted into copy numbers ( $2^{-Ct}$ ) and normalized against the corresponding values for the reference gene. Normalized values were subsequently compared to the reference strain (Gonzales) to determine the fold change in expression level of CYP3006G8 for each strain. Data was analyzed using the Students T-test to determine levels of significance.

#### *Isolation of CPR and cytochrome b<sub>5</sub>*

Bioinformatic analysis allowed the 5' and 3' ends of the *R. microplus* CPR gene to be identified, allowing gene specific primers to be designed for use in PCR to amplify the full length cDNA. Reactions were set up as described above using the following primer set: forward, 5' CACGCGAGCCATGGAAGGGACG 3' and reverse, 5' GCAGCATCAACTCCAGACGTC 3'.

The full length cDNA of *R. microplus* cytochrome b<sub>5</sub> was identified via bioinformatics and primers were designed to amplify this gene. Reactions were set up as described above using the following primer set: forward, 5' CGTTCGCCAGTAGCTTTCCT 3' and reverse, 5' CCTGGTCTGTTTGCCCTCAT 3'. Both full length cDNAs were cloned into pCR Blunt and sequences confirmed via double strand sequencing using the vector specific M13 forward and reverse primers.

### *Peptide sequence analysis*

Each of the isolated nucleotide sequences was analyzed using ExPASy translation software (<http://web.expasy.org/translate/>). This allowed the putative start and stop codons to be identified and determination of the open reading frame of each protein. It also allowed the identification of key residues and motifs to further confirm the isolation of each gene. The molecular weight of each protein was estimated using the ExPASy Bioinformatics Resource Portal and the compute pI/MW page ([http://web.expasy.org/compute\\_pi/](http://web.expasy.org/compute_pi/)). Having used NCBI Blast (<http://blast.ncbi.nlm.nih.gov/Blast.cgi>) to investigate the homology of CYP3006G8 to other CYP proteins on the NCBI database and then using the same software to determine the homology of the putative CPR and cytochrome *b*<sub>5</sub> proteins to other similar proteins on the database, this allowed alignments to be carried out to show this homology. Alignments were carried out using Clustal omega software (<http://www.ebi.ac.uk/Tools/msa/clustalo/>).

## **Results**

### *Isolation of CYP3006G8*

Using known pesticide metabolizing CYPs as bait, bioinformatic analysis identified 5 potential homologues within the *R. microplus* transcriptome data set however during RACE-PCR only one set of primers designed to amplify sequence 60, successfully isolated both 5' and 3' products. Using these products gene specific primers were designed to amplify the full length cDNA of the encoded gene. The full length fragment was cloned, sequenced and submitted to Dr. David Nelson (University of Tennessee) for naming and has subsequently been recorded as CYP3006G8. The nucleotide and translated peptide sequence of

CYP3006G8 is shown in Figure 1, highlighting the key heme-binding domain essential for electron transfer and the membrane binding domain. CYP3006G8 has a calculated pI of 8.78 and a predicted molecular weight of 61.4 kDa.

Sequence homology analysis carried out on CYP3006G8 using NCBI Blastp confirmed homology to CYP6 and CYP9 sequences from several mosquito species (Table 2). Pairwise analysis found that CYP3006G8 shares maximum percent identities of 30.41 % to CYP6P3 (accession number AAL93295) from *Anopheles gambiae* and 30.14 % to CYP9J2 (accession number AAK17188) from *Aedes aegypti*. CYP6P3 has been shown to be involved in pyrethroid metabolism and resistance to pyrethroids (Muller *et al.*, 2008) and members of the CYP9J family have been shown to metabolize pyrethroids also (Stevenson *et al.*, 2012). Figure 2 shows an alignment of the protein sequences of these two mosquito CYPs along with CYP3006G8 and an additional NCBI top hit of CYP6I1 (accession number XP\_003744550) from a species of mite (*Metaseiulus occidentalis*).

#### *mRNA expression analysis*

In order to gain further support for the hypothesis that CYP3006G8 is involved in pesticide metabolism, its mRNA expression levels in 11 different strains of *R. microplus* with varying resistance profiles was determined. Figure 3 shows the results of this analysis expressed as log<sub>10</sub> fold change following normalization with ELF1 $\alpha$  and using the Gonzales strain as the reference strain. The results of this analysis revealed that 8 of the 11 strains showed increased levels of expression of CYP3006G8 with the strains Cz, Co and Pq showing increased expression levels that were highly significant (p = 0.0046, p = 0.0046 and p = 0.0056, respectively). The Tx strain was also found to have an increase in mRNA expression level that was significant (p = 0.027). The strain Ca was found to have a significant decrease in mRNA expression (p = 0.016).

These results are interesting as the strains Cz, Co and Tx have all been shown to demonstrate resistance to pesticides, with Cz and Co displaying metabolic resistance to pyrethroids and Tx both target site and metabolic resistance to organophosphates. In addition the strain Pq has been shown to have target site resistance to pyrethroids but it is believed that metabolic resistance may also play a role in this resistance to pyrethroids. Conversely, the reduced mRNA expression level of the strain Ca also correlates with a role for CYP3006G8 in pyrethroid metabolism as this strain is susceptible to pyrethroids but resistant to organophosphates.

However this correlation is not entirely clear cut as Figure 3 shows the expression level of CYP3006G8 is highly variable across the 11 strains, in particular in the De strain which has been shown to demonstrate susceptibility to all classes of pesticides. As such, further biochemical analysis is required.

#### *Isolation of CPR*

Bioinformatic analysis of the transcriptome data set also identified the 5' and 3' ends of *R. microplus* CPR allowing the generation of gene specific primers and the isolation of a full length cDNA. The full length nucleotide and predicted peptide sequences of *R. microplus* CPR are shown in Figure 4. The encoded peptide has a predicted pI of 5.62 and a molecular weight of 77.8 kDa, a sequence alignment with CPR homologues from other species identified the conserved binding motifs for FMN, FAD and NADPH (Figure 5). The results of the pairwise analysis following the alignment are shown in Table 3 and clearly show the high degree of sequence homology amongst cytochrome P450 oxidoreductase from a variety of species but also shows that variation exists.

### *Isolation of Cytochrome b<sub>5</sub>*

A full length cDNA for *R. microplus* cytochrome *b<sub>5</sub>* was identified following bioinformatic analysis of the 115 transcriptomic sequences allowing this gene to be successfully amplified. The full length nucleotide and translated protein sequence are shown in Figure 6. The encoded protein has a predicted pI of 5.3 and a molecular weight of 15.1 kDa and displays a high degree of homology to other species as shown by the alignment in Figure 7, including the highly conserved histidine residues which are important for heme binding and residues important for creating the “*b<sub>5</sub> fold*”. This high degree of homology is further illustrated by values obtained for pairwise analysis presented in Table 4. Figure 6 and Figure 7 illustrate the high number of acidic residues which generates the acidic nature of this protein.

### **Discussion**

Pesticide resistance amongst the arthropods is known to be a major problem in both medicine and veterinary. There are a huge number of arthropods that act as vectors for disease and others that cause massive destruction of farm crops. Despite this there is still very little knowledge and understanding of how arthropods are able to develop resistance via metabolism of commonly used classes of pesticides such as pyrethroids and organophosphates. Cytochrome P450s through the monooxygenase complex have been shown to be a key mechanism of metabolic resistance to pesticides in numerous arthropods yet this area of research has not been well investigated in the Acari, especially the *Ixodidea* which includes the Southern cattle tick and the black-legged tick both of which transmit disease that affect humans and animals.

CYP3006G8 becomes only the third, cytochrome P450 to be isolated from the Southern cattle tick, *Rhipicephalus (Boophilus) microplus*, with the previous two being isolated over a decade ago. Crampton et al., (1999a) isolated a gene that was named Cyp4w1, however the authors used only sequence analysis and phylogenetics to investigate the gene isolated and did not carry out any biochemical characterization. Additionally, CYP4W1 was found not to be associated with pesticide resistance. He et al., (2002) isolated the gene Cyp319a1 from *R. microplus* and carried out semi-quantitative PCR and found no variation in its expression in six different strains with varying resistance profiles. Similar to CYP4W1, no heterologous expression or biochemical characterization has been carried out for CYP319A1.

Bioinformatic analysis suggests that CYP3006G8 displays some homology to CYP6 and CYP9 members although the level of homology to these key pesticide metabolizing CYPs is not greater than the 40% homology required to be placed in these families. Despite this, the results of the alignments as shown in Figure 2 and the mRNA expression analysis of CYP3006G8 in 11 *R. microplus* strains does provide support of a role of this novel CYP in pesticide metabolism.

The strains Cz, Co and Pq all of which have been profiled to display metabolic resistance to pyrethroids were found to have significant increases in the mRNA expression level of CYP3006G8. Additionally, the Tx strain which has been shown to have target site resistance and metabolic resistance to organophosphates showed an increase in expression of CYP3006G8 which despite being resistance to a different class still supports the hypothesis that this enzyme plays a role in metabolism/pesticide resistance. The other strain that had a significant difference in mRNA expression levels of CYP3006G8 was the Ca which displays target site and some metabolic resistance to organophosphates however is susceptible to all other classes including pyrethroids. This is of interest as this strain



displayed down-regulation of CYP3006G8 further supporting the hypothesis that CYP3006G8 is involved in pyrethroid pesticide metabolism. In view the data published for the previous two CYPs from this species, CYP4W1 (Crampton et al., 1999a) and CYP319A1 (He et al., 2002), CYP3006G8 is the first cytochrome P450 isolated from *R. microplus* that putatively metabolizes pyrethroids based on sequence homology and the correlation of mRNA expression with resistance profiling.

In addition to isolating a potential pyrethroid metabolizing CYP, the other components of the monooxygenase complex have also been isolated, the first of which is cytochrome P450 oxidoreductase (Figure 4) the sole electron transfer protein to CYPs. Isolation of this component is essential in determining not only the ability of CYP3006G8 to metabolize pyrethroids but also the activity of all future CYPs isolated from *R. microplus*. Recent work has been investigated the co-factor binding traits of CPR from different species, in particular between human and insects such as *Anopheles gambiae*. This is of interest as the comparison between the human and mosquito isoforms has identified differences in traits which may have far reaching implications (Lian et al., 2011). The results of this comparison found the concentration of the bound co-factors FMN and FAD per mol of enzyme were lower in *An. gambiae* than in the human CPR and that *An. gambiae* CPR was found to bind more weakly to 2',5'-ADP (an analogue to NADP) compared to human CPR. This latter finding is important as 2',5'-ADP can exert conformational changes which can affect the rate of electron transfer and thus affect the P450 catalysis. It is believed these subtle differences could be exploited to design new species specific pesticides. Isolating CPR for *R. microplus* will now allow similar studies to be carried out to identify if the same binding characteristics exists within all hematophagous organisms.

The final component of the monooxygenase complex that was isolated was cytochrome *b<sub>5</sub>*. Again, similar to CPR, this is the first reported isolation of this gene from *R. microplus*,

indeed from any acari species and completes the isolation of the monooxygenase complex from this species. Cytochrome  $b_5$  is known to stimulate, inhibit or have no effect on catalysis by cytochrome P450s with much research carried out in mammalian CYPs (Finn et al., 2011; McLaughlin et al., 2010; Zhang et al., 2005), therefore it is important to be able to co-express  $cytb_5$  to accurately determine the activity of CYPs with different substrates. The isolation of *R. microplus*  $cytb_5$  will now allow co-expression with CYP3006G8 and any future CYPs that are isolated and expressed from *R. microplus* to determine the effect  $cytb_5$  on catalysis and the role it may have in pesticide metabolism and the development of resistance. As no biochemical characterization has been carried out on CYPs from this species, there is no research showing what effect  $cytb_5$  has and therefore this is an important aspect that requires future studies.

By employing the techniques used here and amplifying additional tick CYP genes attempts can continue to increase our knowledge and understanding of the role of cytochrome P450 genes in pesticide metabolism and ultimately the role they have in pesticide resistance. Identifying those CYPs involved provides the potential to investigate novel compounds (or even synergists) either natural such as plant derived products or synthetic that can be used to control tick populations in this case *R. microplus*.

### *Conclusions*

Ultimately, this paper highlights the lack of research currently being undertaken in identifying cytochrome P450s in acari. This is despite the huge problem of pesticide resistance and the transmission of diseases to both humans and wildlife via ticks and mites acting as vectors. However using what data is available, we have been able to start identifying, isolating and characterising novel pesticide metabolizing CYPs and use the isolated monooxygenase components as an important tool in this area of research.

Characterising arthropod CYPs is an important process as it allows a greater understanding as to which pesticides may still have an effect due to susceptibility and which are unlikely to have any substantial effect due to the establishment of resistance in the field and additionally facilitates a more targeted based approach to the development of new pesticides.

### **Acknowledgements**

We thank Dr Felix Guerrero and Dr Adalberto A. Pérez de León for the provision of the transcriptomic sequence data and for the provision of ticks from the 11 strains used in this work. Funding for this study was provided by Northumbria University at Newcastle.

## References

- Abbas, R.Z., Zaman, M.A., Colwell, D.D., Gilleard, J., Iqbal, Z. 2014 Acaricide resistance in cattle ticks and approaches to its management: The state of play. *Vet. Parasitol.* 203, 6-20.
- Baxter, G.D., Barker, S.C. 1998. Acetylcholinesterase cDNA of the cattle tick, *Boophilus microplus*: characterization and role in organophosphate resistance. *Insect Biochem. Mol. Biol.* 28, 581-589.
- Bissinger, B.W., Roe, R.M. 2010. Tick repellents: Past, present, and future. *Pestic. Biochem. Physiol.* 96, 63-79.
- Brogdon, W.G., McAllister, J.C. 1998. Insecticide resistance and vector control. *Emerg. Infect. Dis.* 4, 605-613.
- Chandor-Proust, A., Bibby, J., Regent-Kloeckner, M., Roux, J., Guittard-Crilat, E., Poupardin, R., Riaz, M.A., Paine, M., Dauphin-Villemant, C., Reynaud, S., David, J.P. 2013. The central role of mosquito cytochrome P450 CYP6Zs in insecticide detoxification revealed by functional expression and structural modelling. *Biochem. J.* 455, 75-85.
- Chen, X., Zhang, Y. 2014. Identification and characterization of NADPH-dependent cytochrome P450 reductase gene and cytochrome *b*<sub>5</sub> gene from *Plutella xylostella*: Possible involvement in resistance to beta-cypermethrin. *Gene* 558, 208-14.
- Chiu, T.L., Wen, Z.M., Rupasinghe, S.G., Schuler, M.A. 2008. Comparative molecular modeling of *Anopheles gambiae* CYP6Z1, a mosquito P450 capable of metabolizing DDT. *Proc. Natl. Acad. Sci. U. S. A.* 105, 8855-8860.
- Corbel, V., N'Guessan, R., Brengues, C., Chandre, F., Djogbenou, L., Martin, T., Akogbeto, A., Hougard, J.M., Rowland, M. 2007. Multiple insecticide resistance mechanisms in

*Anopheles gambiae* and *Culex quinquefasciatus* from Benin, West Africa. *Acta Tropica* 101, 207-216.

Cossio-Bayugar, R., Barhoumi, R., Burghardt, R.C., Wagner, G.G., Holman, P.J. 2002. Basal cellular alterations of esterase, glutathione, glutathione S-transferase, intracellular calcium, and membrane potentials in coumaphos-resistant *Boophilus microplus* (Acari : Ixodidae) cell lines. *Pestic. Biochem. Physiol.* 72, 1-9.

Crampton, A.L., Baxter, C.D., Barker, S.C. 1999a. A new family of cytochrome P450 genes (CYP41) from the cattle tick, *Boophilus microplus*. *Insect Biochem. Mol. Biol.* 29, 829-834.

Crampton, A.L., Baxter, G.D., Barker, S.C. 1999b. Identification and characterization of a cytochrome P450 gene and processed pseudogene from an arachnid: the cattle tick, *Boophilus microplus*. *Insect Biochem. Mol. Biol.* 29, 377-384.

Drali, R., Benkouiten, S., Badiaga, S., Bitam, I., Rolain, J.M., Brouqui, P. 2012. Detection of a knockdown resistance mutation associated with permethrin resistance in the body louse *Pediculus humanus corporis* by use of melting curve analysis genotyping. *J. Clin. Microbiol.* 50, 2229-2233.

Finn, R.D., McLaughlin, L.A., Hughes, C., Song, C.L., Henderson, C.J., Wolf, C.R. 2011. Cytochrome *b*<sub>5</sub> null mouse: a new model for studying inherited skin disorders and the role of unsaturated fatty acids in normal homeostasis. *Transgenic Res.* 20, 491-502.

Gong, M.Q., Gu, Y., Hu, X.B., Sun, Y., Ma, L., Li, X.L., Sun, L.X., Sun, J., Qian, J., Zhu, C.L. 2005. Cloning and overexpression of CYP6F1, a cytochrome P450 gene, from deltamethrin-resistant *Culex pipiens pallens*. *Acta Biochim. Biophys. Sin.* 37, 317-326.

Guengerich, F.P. 2007. Mechanisms of cytochrome P450 substrate oxidation: MiniReview. *J. Biochem. Mol. Toxicol.* 21, 163-168.

Guerrero, F.D., Nene, V.M., George, J.E., Barker, S.C., Willadsen, P. 2006. Sequencing a new target genome: The *Boophilus microplus* (Acari : Ixodidae) genome project. J. Med. Entomol. 43, 9-16.

He, H., Chen, A.C., Davey, R.B., Ivie, G.W. 2002. Molecular cloning and nucleotide sequence of a new P450 gene, CYP319A1, from the cattle tick, *Boophilus microplus*. Insect Biochem. Mol. Biol. 32, 303-309.

Hemingway, J., Hawkes, N.J., McCarroll, L., Ranson, H. 2004. The molecular basis of insecticide resistance in mosquitoes. Insect Biochem. Mol. Biol. 34, 653-665.

Hill, C.A., Wikel, S.K. 2005. The *Ixodes scapularis* Genome Project: an opportunity for advancing tick research. Trends Parasitol. 21, 151-153.

Im, S.C., Waskell, L. 2011. The interaction of microsomal cytochrome P450 2B4 with its redox partners, cytochrome P450 reductase and cytochrome *b*<sub>5</sub>. Arch. Biochem. Biophys. 507, 144-53.

Iyanagi, T., Xia, C.W., Kim, J.J.P. 2012. NADPH-cytochrome P450 oxidoreductase: Prototypic member of the diflavin reductase family. Arch. Biochem. Biophys. 528, 72-89.

Kawada, H., Futami, K., Komagata, O., Kasai, S., Tomita, T., Sonye, G., Mwatele, C., Njenga, S.M., Mwandawiro, C., Minakawa, N., Takagi, M. 2011. Distribution of a knockdown resistance mutation (L1014S) in *Anopheles gambiae* s.s. and *Anopheles arabiensis* in Western and Southern Kenya. Plos One 6, e24323.

Kushwah, R.B., Dykes, C.L., Kapoor, N., Adak, T., Singh, O.P. 2015. Pyrethroid-resistance and presence of two knockdown resistance (kdr) mutations, F1534C and a novel mutation T1520I, in Indian *Aedes aegypti*. PLoS Negl. Trop. Dis. 9, e3332.

- Kwon, D.H., Clark, J.M., Lee, S.H. 2010. Extensive gene duplication of acetylcholinesterase associated with organophosphate resistance in the two-spotted spider mite. *Insect Mol. Biol.* 19, 195-204.
- Li, A.Y., Davey, R.B., Miller, R.J., George, J.E. 2004. Detection and characterization of amitraz resistance in the southern cattle tick, *Boophilus microplus* (Acari : Ixodidae). *J. Med. Entomol.* 41, 193-200.
- Lian, L.Y., Widdowson, P., McLaughlin, L.A., Paine, M.J.I. 2011. Biochemical comparison of *Anopheles gambiae* and Human NADPH P450 Reductases reveals different 2'-5'-ADP and FMN binding traits. *Plos One* 6, e20574.
- Liu, D., Zhou, X.J., Li, M., Zhu, S.Y., Qiu, X.H. 2014. Characterization of NADPH-cytochrome P450 reductase gene from the cotton bollworm, *Helicoverpa armigera*. *Gene* 545, 262-270.
- Mamidala P., Jones S.C., Mittapalli, O. 2011. Metabolic resistance in Bed Bugs. *Insects* 2, 36-48.
- Marcombe, S., Poupardin, R., Darriet, F., Reynaud, S., Bonnet, J., Strode, C., Brengues, C., Yebakima, A., Ranson, H., Corbel, V., David, J.P. 2009. Exploring the molecular basis of insecticide resistance in the dengue vector *Aedes aegypti*: a case study in Martinique Island (French West Indies). *BMC Genomics* 10, 494.
- McLaughlin, L.A., Ronseaux, S., Finn, R.D., Henderson, C.J., Wolf, C.R. 2010. Deletion of microsomal cytochrome *b*<sub>5</sub> profoundly affects hepatic and extrahepatic drug metabolism. *Mol. Pharmacol.* 78, 269-278.
- Muller, P., Warr, E., Stevenson, B.J., Pignatelli, P.M., Morgan, J.C., Steven, A., Yawson, A.E., Mitchell, S.N., Ranson, H., Hemingway, J., Paine, M.J.I., Donnelly, M.J. 2008. Field-

caught permethrin-resistant *Anopheles gambiae* overexpress CYP6P3, a P450 that metabolizes pyrethroids. Plos Genet. 4, e1000286.

Nijhof, A.M., Balk, J.A., Postigo, M., Jongejan, F. 2009. Selection of reference genes for quantitative RT-PCR studies in *Rhipicephalus (Boophilus) microplus* and *Rhipicephalus appendiculatus* ticks and determination of the expression profile of Bm86. BMC Mol. Biol. 10, 112.

Porter, T.D. 2002. The roles of cytochrome *b*<sub>5</sub> in cytochrome P450 reactions. J. Biochem. Mol. Toxicol. 16, 311-6.

Rodriguez-Vivas, R.I., Alonso-Diaz, M.A., Rodriguez-Arevalo, F., Fragoso-Sanchez, H., Santamaria, V., Rosario-Cruz, R. 2006. Prevalence and potential risk factors for organophosphate and pyrethroid resistance in *Boophilus microplus* ticks on cattle ranches from the State of Yucatan, Mexico. Vet. Parasitol. 136, 335-342.

Scott, J.G., Wen, Z.M. 2001. Cytochromes P450 of insects: the tip of the iceberg. Pest Manag. Sci. 57, 958-967.

Schenkman, J.B., Jansson, I. 2003. The many roles of cytochrome *b*<sub>5</sub>. Pharmacol. Ther. 97, 139-52.

Stevenson, B.J., Pignatelli, P., Nikou, D., Paine, M.J.I. 2012. Pinpointing P450s associated with pyrethroid metabolism in the Dengue Vector, *Aedes aegypti*: Developing new tools to combat insecticide resistance. Plos Negl. Trop. Dis. 6, e1595.

Syed, K., Mashele, S.S. 2014. Comparative analysis of P450 signature motifs EXXR and CXG in the large and diverse kingdom of fungi: Identification of evolutionarily conserved amino acid patterns characteristic of P450 family. Plos One 9, e95616.



Sztaf, T., Chung, H., Berger, S., Currie, P.D., Batterham, P., Daborn, P.J. 2012. A Cytochrome P450 conserved in insects is involved in cuticle formation. *Plos One* 7, e36544.

Van Zee, J.P., Geraci, N.S., Guerrero, F.D., Wikel, S.K., Stuart, J.J., Nene, V.M., Hill, C.A. 2007. Tick genomics: The Ixodes genome project and beyond. *Int. J. Parasitol.* 37, 1297-1305.

Werck-Reichhart, D., Feyereisen, R. 2000. Cytochromes P450: a success story. *Genome Biol.* 1, REVIEWS3003.

Wondji, C.S., Dabire, R.K., Tukur, Z., Irving, H., Djouaka, R., Morgan, J.C. 2011. Identification and distribution of a GABA receptor mutation conferring dieldrin resistance in the malaria vector *Anopheles funestus* in Africa. *Insect Biochem. Mol. Biol.* 41, 484-491.

Zhang, H.M., Myshkin, E., Waskell, L. 2005. Role of cytochrome *b*<sub>5</sub> in catalysis by cytochrome P4502B4. *Biochem. Biophys. Res. Comm.* 338, 499-506.

Zhou, S., Chan, E., Li, X., Huang, M. 2005. Clinical outcomes and management of mechanism-based inhibition of cytochrome P450 3A4. *Ther. Clin. Risk Manag.* 1, 3-13.

Zhu, F., Sams, S., Mural, T., Haynes, K.F., Potter, M.F., Palli, S.R. 2012. RNA interference of NADPH-Cytochrome P450 Reductase results in reduced insecticide resistance in the bed bug, *Cimex lectularius*. *Plos One* 7, e31037.

**Table 1:** The 11 strains of *R. microplus* along with the resistance profile of each strain used during expression analysis.

<b><i>R. microplus</i> Strain</b>	<b>Pesticide resistance or susceptibility</b>
<b>Gonzalez (Gz)</b>	Susceptible to all, very low level of metabolic resistance
<b>Coatzoacoalcos (Cz)</b>	Metabolic resistance to pyrethroids, target site pyrethroid resistance is absent, susceptible to other classes through low level of metabolic resistance to Organophosphates (OPs).
<b>San Roman (SR)</b>	Target site and metabolic resistance to OPs, susceptible to other classes.
<b>Caporal (Ca)</b>	Target site and metabolic resistance to OPs, susceptible to other classes.
<b>San Alphonso (San Alfonso) (SA)</b>	Metabolic resistance to amitraz probably. Target site resistance to pyrethroids.
<b>Santa Luiza (SL)</b>	Amitraz and pyrethroid resistance though mechanism unknown. Expect metabolic resistance is important
<b>Corrales (Co)</b>	Target site resistance to pyrethroids, some metabolic resistance to both pyrethroids and OPs also present.
<b>Pesqueria (Pq)</b>	Target site resistant to pyrethroids, probably some metabolic resistance to pyrethroids also present.
<b>Deutsch (De)</b>	Susceptible to OPs and pyrethroids.
<b>San Felipe (SF)</b>	Target site and metabolic resistance to pyrethroids. Probably low level of metabolic resistance to OPs.
<b>Tuxpan (Tx)</b>	Target site and metabolic resistance to OPs

**Table 2: Pairwise analysis results.** Results of pairwise analysis following alignment of CYP3006G8 with other arthropod CYPs shown in Figure 2

	Rm_CYP3006G8	M.occidentalis_CYP6I1	An.gambiae_CYP6P3	Ae.aegypti_CYP9J2
Rm_CYP3006G8	100.00	33.52	30.41	30.14
M.occidentalis_CYP6I1	33.52	100.00	25.10	23.96
An.gambiae_CYP6P3	30.41	25.10	100.00	31.09
Ae.aegypti_CYP9J2	30.14	23.96	31.09	100.00

**Table 3: Pairwise analysis results.** Results of pairwise analysis following alignment of *R. microplus* CPR with other CPR sequences

	H.sapiens	R.microplus	Ae.aegypti	D.melanogaster
H.sapiens	100.00	56.95	56.12	56.87
R.microplus	56.95	100.00	58.01	60.00
Ae.aegypti	56.12	58.01	100.00	78.85
D.melanogaster	56.87	60.00	78.85	100.00

**Table 4: Pairwise analysis results.** Results of pairwise analysis following alignment of *R. microplus* cytochrome *b*<sub>5</sub> with other cytochrome *b*<sub>5</sub> sequences

	H.sapiens	R.microplus	D.melanogaster	An.gambiae	Cu.quinquefasciatus
H.sapiens	100.00	51.16	46.15	44.44	46.46
R.microplus	51.16	100.00	50.00	48.82	53.12
D.melanogaster	46.15	50.00	100.00	60.94	67.44
An.gambiae	44.44	48.82	60.94	100.00	85.94
Cu.quinquefasciatus	46.46	53.12	67.44	85.94	100.00

**Figure 1: Full Length sequence of CYP3006G8.** Nucleotide and translated peptide sequence of the full length CYP3006G8 with the start and stop codons in bold/ underlined and characteristic heme binding domain (FxxGxxxCxG) bold/shaded. Membrane binding domain dashed underline.

**Figure 2: Amino acid alignment of CYP3006G8 with other arthropod CYPs.** CYP3006G8 aligned using Clustal omega with CYP6I1, CYP6P3 and CYP9J2 returned as top hits following Blastp searches. Heme-binding domain highlighted grey, ExxR motif boxed with solid line and PERF motif boxed with dashed line.

**Figure 3: CYP3006G8 mRNA expression level as fold change compared to Gonzales.** Fold change data transformed ( $\text{Log}_{10}$ ). Strains Cz (P = 0.0046), Co (P = 0.0046), and Pq (P = 0.0056) showed highly significant difference (\*\*) and strains Tx (P = 0.0027) and Ca (P = 0.016) showed significant difference (\*). Strains labelled as shown in Tables 1.

**Figure 4: Nucleotide and translated peptide sequence of *R. microplus* Cytochrome P450 Oxidoreductase.** Data obtained following successful PCR amplification of CPR and subsequent sequencing. Start and stop codons in bold and underlined.

**Figure 5: Protein alignment of *R. microplus* CPR with key regions highlighted.** Alignment showing high degree of homology of CPR between different species. Membrane binding region dashed underlined. Key co-factor-binding regions are highlighted, FMN domains shaded grey, FAD domains boxed with a dashed line and NADP(H) domains boxed with a solid line.

**Figure 6: Nucleotide and translated protein sequence of *R. microplus* cytochrome *b<sub>5</sub>*.** Data obtained following successful PCR amplification. Start and stop codons are in bold and underlined.

**Figure 7: Peptide sequence alignment of *R. microplus* cytochrome *b*<sub>5</sub> and other arthropod and mammalian Cyt*b*<sub>5</sub> sequences.** An alignment showing the high degree of homology between species for the protein, cytochrome *b*<sub>5</sub>. Conserved heme-coordinating histidine residues in the heme binding domain boxed. Residues highlighted grey indicates amino acids in the “*b*<sub>5</sub> fold”. Membrane-binding domain dashed underlined.

Figure 1

atgctcgaagtgtttctcttgagcttgatcgctcgtcttcgcaacatggttcttcatacaacgaaggagg  
**M** \_ **L** \_ **E** \_ **V** \_ **F** \_ **L** \_ **L** \_ **S** \_ **L** \_ **I** \_ **V** \_ **V** \_ **F** \_ **A** \_ **T** \_ **W** \_ **F** \_ **F** \_ **I** \_ **Q** \_ **R** \_ **R** \_ **R**  
cggttttccttctttaagacctcggaattccagggtccccacccaagttttctttccggaaactgtgcc  
R F S F F K D L G I P G P P P S F L S G N L S  
gaacttatacaaaaaggaacactggagaaatacaaggaatggctggacaaatattggtgacattgttgg  
E L I Q K G T L E K Y K E W L D K Y G D I V G  
ttttacaacggcgctcaccattttcttatcgtaaaagacccggagctaatcaagaagatccaaataaag  
F Y N G A H P F L I V K D P E L I K K I Q I K  
gatttccacaattttcatggccgaggggtgtcatctggcttcgcgagaactcatccgatcaacaaagaa  
D F H N F H G R G V S S G F A R T H P I N K E  
agtatgataaacgctcaaggagagcgctggaagaagatgcgagctcttctgacgcccgtttcagcagc  
S M I N A Q G E R W K K M R S L L T P A F T T  
agcaacatgaagaagatggcaagcctaagcctaattggacgacagctccaacgagttttcttcaagtcatc  
S N M K K M A S L S L M D D S S N E F L Q V I  
gaatccctgaggaagaaggacgaagctctcgagttccgcgatcttttccagagactcaccgcagacgctc  
E S L R K K D E A L E F R D L F Q R L T A D V  
atcattcgatcggcggtttgcggtttggctctcaaatccgacctgcagcaaaaggaccgattgaagagcaca  
I I R S A F A F G L K S D L Q Q K D R L K S T  
acagagtcgctgtttcgggagacggttgatagcttgcaacaattttcgccgggctggataaatttcctt  
T E S L F R E T L D S L Q Q F R R A W I N F L  
actgcttgctttccggagtttaatoctttgtggaggctgatcatatcattcggttcacgacataacaag  
T A C F P E F N P L W R L I I S F G S R H N K  
acggcggcagacaagtgttttgacgagatcacttccatcatacagtttcgctcgtgaaaaccgtgagaga  
T A A D K C F D E I T S I I Q F R R E N R E R  
gataggtgcatcttctgcagctgatgctcaatgctgaagtggaggacgccacccttgtgaacgtgcac  
D R C D L L Q L M L N A E V E D A T L V N V H  
tcgctcacggcatccgggatgctgacagcgcatctgaagggaaccagcctgaaaaggtaaaagcagggc  
S L T A S G D A D S A S E G N Q P E K V K A G  
ggagcaggcggaaaaaacgtgtgttctcacgaatacagaaatcttggccaatggcttcagtttcttctg  
G A G G K T C V L T N T E I L A N G F S F F V  
gcaggggttcgagacgacaggctcatcgatggcattttctgtcatatctcctcgcgaaacaccaggacatt  
A G F E T T G S S M A F L S Y L L A K H Q D I  
caggaccgactcagagaagacgctccttgccgctcctcaatagagacgggtgccttcacgtacgacaacgctc  
Q D R L R E D V L A V L N R D G A F T Y D N V  
tttggcataaaaatacctggaccaggctatatcagaatctctgcgcttttatttcgccagctcgtagggttc  
F G I K Y L D Q A I S E S L R F Y S P V V G F  
actacaagaagatgctcgtgaatacgtgcacaagggtatgaaaatacccgtggaaccagcatcgtc  
T T R R C A R E Y V H K G M K I P A G T S I V  
gtcatccccaaccaccactgagtcatgaccgaatttctgggagcaaccggaagtcttctgatccagaa  
V I P N H H L S H D P N F W E Q P E V F D P E  
agattcagcccacaaaataaaggccttggatcccgtgggtgtaccagcctt**ttcggtcagggccccgc**  
R F S P Q N K G L V D P V V Y Q P **F G Q G P R**  
**aattgogtggg**atgagattcgcagctggagatgaaactaacatggcgaaattattggcaaaatac  
**N C V G** M R F A Q L E M K L T M A K L L A K Y  
aagctttttctggatgaacgtcatattaaggagaaaaacttggagctggaatccactttcatattcgcg  
K L F L D E R H I K E K N L E L E S T F I F A  
atgcctaaggatggcatctggctcaagatcgaaaaagttatc**taa**  
M P K D G I W L K I E K V I **STOP**




## Figure 2

```

Rm_CYP3006G8      MLEVFLLSLIVV----FATWFFIQRRRRFSFFKDLGIPGP-PPSFLSGNLSELIQKGTLE
M.occidentalis_CYP611 MLFAVLSTLLVA----YFVYCLRWRLKVLISIFRRLNIPGP-SPRFLWGNMYEYIKHGHVLV
An.gambiae_CYP6P3    ---MELINAVLAAFI FAVSIVYLFIRNKHNYWKDNGFPYAPNPHFLFGHAKGQAQTRHGA
Ae.aegypti_CYP9J2    MVEVNLFSALAV--GAVILLLYHYIAKKYHYFLTKPIPCI-KPTFLLGIFDMVVLKRVEL
                    * . : . : : * * * *
                    -----
Rm_CYP3006G8      K-----YKEWLDKYGDIVGFYNGAHPFLIVKDPELIKKIQIKDFHNFHGRGVSSGFA-RT
M.occidentalis_CYP611 M-----QKKWHEQYGPVVGfyfGLFPCLLVSDTELLKRIILKDFHNFADRSHIISG--EN
An.gambiae_CYP6P3    DIHQELYRYFKQRGERYGGISQFIVPSVLVIDPELAKTILVKDFNVFHDHGVFTN---AK
Ae.aegypti_CYP9J2    VFGS-KLLYNSYPDAKIIGYYELTKPTYMVRDPEMIKKAIAIKDFDSFTDRTPVFGDAVPA
                    * * : * * * : * * : * * * * *
Rm_CYP3006G8      HPINKESMINAQGERWKKMRSLLTPAFTTSMKMMASLSLMDSSNEFLQVI--ESLRKK
M.occidentalis_CYP611 GSIIDDSLLTLKGQRWKEVRSTLTPSFTSKKLLKVTPE--VSSAADEFMKNV--DRHFAE
An.gambiae_CYP6P3    DDPLTGHLFALEGQPWRLMRQKLTPTFTSGRMKQMFGTI--RDVGLELEKCM---EQSYN
Ae.aegypti_CYP9J2    DSLFFNSLFLSLRGQKWRDMRSTLSPAFTGSRMRMYAELV--VKCATSMTDFIHSEAKAGR
                    :: . * : * : * : * * * * * : : : : . . . . .
Rm_CYP3006G8      DEALEFRDLFQRLTADVIRSAFAFGLK--SDLQQKDRLKSTTESLFRETLDLSLQQFRRA
M.occidentalis_CYP611 DAECDFYELFQALTLDTICTKGMGVDFEIQSNIDNSAILR-----QVKMMLAYQID
An.gambiae_CYP6P3    QPEVEMKDILGRFTTVDVIGTCAFGIECNTLKTTDSEFRKYG-NKAFELNTMIM-----
Ae.aegypti_CYP9J2    RLEFNMKDTFSRFVCDAIASVAFGIEVDSFRDPENEFYKKG-NETQKIHTFKS-----
                    :: : : : . * . * . . . . . : . . . . .
Rm_CYP3006G8      WINFLTACFPFENPLWRLI-ISFGSRHNKTAAD---KCFDEITSIIQFRR-ENRERDRCD
M.occidentalis_CYP611 LILLIIVSFPSSLQLAGKLLTALAQRAKNSGRDPVIVKLKQCGDVVKLRRSDTEKSERSD
An.gambiae_CYP6P3    MKTFLASSYPTL-VRNLHMKIT-----YNDVERFFLDIVKETVDYRE--ANNVCRND
Ae.aegypti_CYP9J2    LATFVTLRFVFPFLQKVFNFDFV-----DANVAGYFKKLI SDNMDQRK--KQGIVRND
                    :: : : : : : : : . . . * . . * *
Rm_CYP3006G8      LLQLMLNAEVEDATLVNVHSLTASGDADSASEGNQPEKV-----KAGGAGGKTCVLTN
M.occidentalis_CYP611 LLQLMMDAQNNTVAHATDLKSMIAGDDAEAVEA-EASLKI AKKQSI PSECFAGKPKGLTD
An.gambiae_CYP6P3    FMNLMLQIKNKGLDD-----SDD-----GSLGKG--EVGMTQ
Ae.aegypti_CYP9J2    LVNMLMETKKGALKYE-----EPDLQVSEGYATVE-----ESHVGKSTHSRIWTD
                    ::::: : : . * :
Rm_CYP3006G8      TEILANGFSFFVAGFETTGSMAFLSYLLAKHQDIQDRLEDVLAVL--NRDGAFTYDNV
M.occidentalis_CYP611 SEVIDNALLLLLAGYETTSSMAFMKLLVRFPEVQERMREELLDAT--GGGKVFDFERL
An.gambiae_CYP6P3    NELAAQAFVFFLAGFETSSTTQSFCLYELAKNPDIQERLREEINRAI-AENGGEVTYDVV
Ae.aegypti_CYP9J2    NELISQCFFFFFAAFDNVSSILAFSLYELTVNQDIQRRLYEEIAATESTLNQOPITYEAL
                    . * : : : : * . . : * * * : * : . . : :
Rm_CYP3006G8      FGIKYLDQAISESLRFYSPVVGFTTRRCAREYVHK-----GMKIPAGTSIVVIPNHHLSH
M.occidentalis_CYP611 QRCQYTEAVLQESLRMYPPPIYFLTRVAAAEIKYG-----SLTIPRGMNIFSSV-NQLHY
An.gambiae_CYP6P3    MNIKYLDNVIDETLRKYPPVES-LTRVPSVDYLIP---GTKHVIPKRTLVIQI-PAYAIQR
Ae.aegypti_CYP9J2    QKMAYLDMVVSSEALRKYPTATL-TDRYANKDYVFDDEEGLRFVIEKGTIWI-SMLALHH
                    * : . . * : * * * * : : :
Rm_CYP3006G8      DPNFWEQPEVFDPERFSPQNKGLVDPVVYQPFQGGPRNCVGMRFAQLEMKLTMAKLLAKY
M.occidentalis_CYP611 DEDVFPLPYEFQPERFLPENKTPAMAWYQPFQGGPRNCIGMRFAQMEIKLTMAKLLTKY
An.gambiae_CYP6P3    DPDHYDPERFNPDRFLPEEVKRRHPFTFIPFGEGPRICIGLRFGLMQTKVGLITLLRKF
Ae.aegypti_CYP9J2    DPKYFPEPERFDPERFSEDNRSKIVPGTYLPGAGPRSCIGPRLALLEVKMALYHLVKDF
                    * . : * * * * * : : : * * * * * * * * : : * : : * : .
                    -----
Rm_CYP3006G8      KLFLDERHIKEKNLELESTFIFAMPKDGIIWLKIEKVI--
M.occidentalis_CYP611 RMTSESEPEHEAYIETLVMPVLQKIKDPVKCKLHLL---
An.gambiae_CYP6P3    RFSPSARTPE--RVEYDPKMITIAPKAGNYLKVEKL---
Ae.aegypti_CYP9J2    NLQPSEKTQI--PLRLSKSAFTMQAENGWVLELKARPKA
                    . : . . : . : : : :

```

Figure 3



**Figure 4**

**atg**gaagggacgagccaagatgttccgctgagtgtcgacagcagcccgctgggaagtcagaatgaggaggcccc  
**M** E G T S Q D V P L S V D S S P L G S Q N E E A P  
ctgtttggcgtgttgacgtgctcatccttctggcattgctgggctttgccttctactggctcttctgaggcgc  
L F G V L D V L I L L A L L G F A F Y W L F L R R  
aagaagggccccaccttcgatccggccgcatcaaacggtttccattgaaacaagtatacagaaggctgacaac  
K K A P T F D P A A I K T F S I E T S I Q K A D N  
accagctttattggaaagatgaaatcaacgggttaggaatattgtcatattctatggatcacagacgggcacagcg  
T S F I G K M K S T G R N I V I F Y G S Q T G T A  
gaagagtttctgctcgtcttgcaaaagaagccaacagatttgggtctaaaagcaatggttgctgacccccgaggag  
E E F P A R L A K E A N R F G L K A M V A D P E E  
tgtgaaatggaggacctgaccaagctgccagagatcagcaattcgatggccatcttttgatggcaacgtacggt  
C E M E D L T K L P E I S N S M A I F C M A T Y G  
gaagtgacccccacagacaatgcgcaagactttaccagtggttcaggatggcagcgttgacctgctggagtc  
E G D P T D N A Q D F Y Q W L Q D G S V D L P G V  
aattatgcagtggtttgcattgggaaacaagacgtacgagcacttcaatgcgatgggcaagtagctcgacaagcgc  
N Y A V F A L G N K T Y E H F N A M G K Y V D K R  
atggaggaactgggagccaccctgtctttgaactgggctcggagatgatgatgctaacattgaggaagacttc  
M E E L G A T R V F E L G L G D D D A N I E E D F  
gtgacctggaaggagcgttctggaatgccgtttgtgaaaacttcgaaaaggtcttccacggagagatctcgcgc  
V T W K E R F W N A V C E N F H L E I S G E D I N  
ctccacctggagatctctggcgaggatatacaacctccgpcagtagcattgtgacactgatcttccatct  
L R Q Y Q L I V H T D L P S E K V F H G E I S R L  
aactcataccaccagaaaatgccctttgatgcaaaagaatccattcctggctcctggttcgagtgacagaa  
N S Y T T Q K M P F D A K N P F L A P V R V H K E  
ttatacaaaaggctctcgtcgtcatgcatgcacattgagattagcattgccggctccaaaatgaggtatgatgctggt  
L Y K G S R S C M H I E I S I A G S K M R Y D A G  
gaccacgtggccgtctatccgatgaatgacgtcgccatagtgaggaaacctaggacaaaatgctcaaggtggactg  
D H V A V Y P M N D V A I V E N L G Q M L K V D L  
gacacagtcatcacactgaaaaatcttgatgaggacagctccaagaagcatccatttccctgcccattgtagttat  
D T V I T L K N L D E D S S K K H P F P C P C S Y  
agaaccgcactgctgtactactcgacatcaccactccccctaggacacacagctcttgaaggaatctccgagat  
R T A L L Y V D I T T P R T H V L K E I S E Y  
gccaccaacgaggaggagaaaaaadgtgaagttgatgagctcttcttcagatgaaggaagcctgtacaag  
A T N E E E K K M L K L M S S S S D E G K S L Y K  
cagtggttactgaatgactgcccagtgctgctgacacattctggaggacctgccatcggcagctcctcctttggac  
Q W V L N D C R S V V H I L E D L P S A R P P L D  
catttgctggagcttatgcccaggctgcaggctcgatactactctatatcatcgctcgccaaaggtgcatccagac  
H L L E L M P R L Q A R Y Y S I S S S P K V H P D  
agcatccacatgacggctgtaaaagtggagtacgagactccaaccaagcgcataaaccacgggtgtggccacaggc  
S I H M T A V K V E Y E T P T K R I N H G V A T G  
tggcttgcaactcaagcgaccgacaacggcactcagcctacgctgctgtgtatgtgcgggcgtgcagttcaag  
W L A L K R P D N G T Q P T L P V Y V R R S Q F K  
ctgccctcacggccgagattcccatcgtgatgggtggggccgggcacgggcttggcgccattccgcggttcatc  
L P S R P Q I P I V M V G P G T G L A P F R G F I  
caagaaaggatttcccttcgaaagaaagcaagcccgtgggtgaagctgtactctactttggctgcccgaagaag  
Q E R D F L R K E S K P V G E A V L Y F G C R K K  
gctgaagactacctgtaccaggaggagctggaggagtacctggcaaatggtacactgaccaagttgtacctcgcc  
A E D Y L Y Q E E L E E Y L A N G T L T K L Y L A  
ttctcgcgagaccagccacaaaagtgtatgtgacacatctcctgcccagaacaaggatgaggtctgggacctc  
F S R D Q P H K V Y V T H L L R Q N K D E V W D L  
attggcaagaagaatggccacttctacatttgggtgatgcacgtaacatggcaagagacgtgcatgagatccta  
I G K K N G H F Y I C G G D A R N M A R D V H E I L  
ctagagatcttccgagagaatggcaacatgtctgaagacgaggcagtgctcctacctcaagcgcagtgagtcgag  
L E I F R E N G N M S E D E A V S Y L K R M E S Q  
cggcgctactcggctgacgtctggagt**tga**  
R R Y S A D V W S STOP

**Figure 5**

H.sapiens	MINMG-DSHVDTSSSTVSEAVAAEEVSLFSMTDMILFSLIVGLLTYWFLFRKKKEEVPEFTK
R.microplus	MEGTSQDVPLSVDSSPLGSQNEEAPLFGVLDVLLALLLGFAYWFLFRKKKAPTDFPAA
Ae.aegypti	---M--D--AQTEPEVPPAPISDEPFLGPLDIILLAVLIGGAAYFLKSKKKTQT--SQ
D.melanogaster	MASE--Q--TIDGAAAIPSGGGDEPFLGLLDVALLAVLIGGAAYFLRSRKKEEE----P
H.sapiens	IQ-----TLTSSVRESSFVEKMKKTGRNIIIVFYGSQTGTAEFFANRLSKDAHRYGMR
R.microplus	IKTFSIET---SIQKADNTSFIGKMKSTGRNIIIVFYGSQTGTAEFFPARLAKEANRFLK
Ae.aegypti	FKYSYIQPTTVNTMTMAENSFIKKLKSSGRRLVVFYGSQTGTAEFFAGRLAKEGLRYQMK
D.melanogaster	TRSYSIQPTTVCTTSASDNSFIKKLKASGRSLVVFYGSQTGTGEEFAGRLAKEGIRYRLK
H.sapiens	GMSADPEEYDLADLSSLPEIDNALVFCMATYGEEDPTDNAQDFYDWLQETDVDLSGVKF
R.microplus	AMVADPEECEMEDLTKLPEISNSMAIFCMATYGEEDPTDNAQDFYQWLQDGSVDLPGVNY
Ae.aegypti	GMVADPEECDMEELLSLKDIDKSLAVFCLATYGEEDPTDNCMEFYDWIQNDVDFSGLNY
D.melanogaster	GMVADPEECDMEELLQLKIDNSLAVFCLATYGEEDPTDNAMEFYEWITSGDVLISGLNY
H.sapiens	AVFGLGNKTYEHFNAMGKYVDKRLLEQLGAQRIFELGLGDDDNLEEDFITWREQFWPAVC
R.microplus	AVFALGNKTYEHFNAMGKYVDKRMEELGATRVFELGLGDDANIEEDFVTWKERFWNAVC
Ae.aegypti	AVFGLGNKTYEHYKNGVIYVDKRLLEELGANRVFELGLGDDANIEEDFITWKDKFWPAVC
D.melanogaster	AVFGLGNKTYEHYKVAIYVDKRLLEELGANRVFELGLGDDANIEEDFITWKDRFWPAVC
H.sapiens	EHFGVEATGEESSIRQYELVVHTDIDAAKVYMGEMGRKLSYENQKPPFDAKNPFLAAVTT
R.microplus	ENFHLEISGEDINLRQYQLIVHTDLPSEKVFHGEISRLNSYTTQKMPFDAKNPFLAPVRV
Ae.aegypti	DHFGIESSGEEVLMRQYRLLEQPETPTERLYTGEVARLHSLQTQRPPFDAKNPFLAPIKV
D.melanogaster	DHFGIEGGGEEVLIQRYLLEQPDPVQPDRIYTGELIARLHSIQNRPPFDAKNPFLAPIKV
H.sapiens	NRKLNQGTERHLMHLELDISDSKIRYESGDHVAVYPANDSALVNQLGKILGADLDVVMSSL
R.microplus	HKELYK-GSRSCMHIEISIASGSKMRYDAGDHVAVYPMNDVAIVENLQMLKVDLDTVITL
Ae.aegypti	NRELHKAGGRSCMHIEFDIEGSKMRYEAGDHLAMYPVNDQDLVLRGLKLCNADLDTIFSL
D.melanogaster	NRELHKGGGRSCMHIELSIEGSKMRYDAGDHVAMFPVNDKSLVEKLGQLCNADLDTVFSL
H.sapiens	NNLDEESNKKHPFPCPTSRYRTALTYLDITNPPRTNVLYELAQYASEPSEQELLRKMASS
R.microplus	KNLDEDSKKHPFPCPCSYRTALLYVDITPPRTHVLKEISEYATNEEEKMLKLMSSS
Ae.aegypti	INTDTDSSKKHPFPCPTTYRTALTHYLEITAIPTRTHILKELAEYCSEEDKFEFLRMCST
D.melanogaster	INTDTDSSKKHPFPCPTTYRTALTHYLEITAIPTRTHILKELAEYCTDEKEKELLRSMASI
H.sapiens	SGEGKELYLSWVVEARRHILAILQDCPSLRPPIIDHLCCELLPRLQARYYSIASSSKVHPNSI
R.microplus	SDEGKSLYKQWVLNDCRSVWHILEDLPSARPLDHLLELMPRLQARYYSISSSPKVHPDSI
Ae.aegypti	NPEGKAKYQEWVQDCRNIVHVLEDLPSCRPPIIDHICELLPRLQPRYYSISSSSKLYPTTI
D.melanogaster	SPEGKEYQSWIQDACRNIVHILEDIKSCRPPIIDHVCELLPRLQPRYYSISSSAKLHPTTI
H.sapiens	VHICAVVVEYETKAGRINKGVATNWLRAKEPAGENGRALVPMFVRKSQLRFPFKATTEV
R.microplus	IHMATAVKVEYETPTKRINHGVATGWLALKRPDNGT--QPTLPVYVRRSQFKLPSRQIEPI
Ae.aegypti	VHVTAVLVKYETKTGRVNHGVATTFLSQKHPDGE-PLPRVPIFIRKSQLRPAKTETEIV
D.melanogaster	VHVTAVLVVEYKTPTGRINKGVATTYLNKQPOGSE--EVKVPVIFIRKSQLRPLTKPETEIV
H.sapiens	IMVGPGTGVAPFIFGFIQERAWLRQQGKEVGETLLLYGCRRSDEDYLYREELAQFHRDGL
R.microplus	VMVGPGTGLAPFRGFIQERDFLRKESKPVGEAVLYFGCRKKAEDYLYQEELEEYLANGTL
Ae.aegypti	IMVGPGTGLAPFRGFIQERDFNKKDGKEVGQTIYLYFGCRKRSEDIYIEELEEYVQRGIM
D.melanogaster	IMVGPGTGLAPFRGFIQERQFLRDEGKTVGESILYFGCRKRSEDIYIEELEEYVKKGTL
H.sapiens	TQLNVAFSRECSHKVYVQHLLKQDREHLWKLIE-GGAHIYVCGDARNMAHDVQNTFYDIV
R.microplus	TKLYLAFSRDQPHKVYVTHLLRQNKDEVDLIGKKNHFYICGDARNMAHDVHEILLEIF
Ae.aegypti	-KLRTAFSRDQAHKVYVTHLLEEDMDLLWNVIGENKGFYICGDAKNMATDVRNILLKVL
D.melanogaster	-NLKAAFSRDQGKKVYVQHLLQDADLIWNVIGENKGFYICGDAKNMATDVRNILLKIL
H.sapiens	AELGAMEHAQAVDYIKKLMTKGRYSLDVWS
R.microplus	RENGNMSEDEAVSYLKRMEQGRYSADVWS
Ae.aegypti	QTKGSMSESEAIQYIKKMEAQGRYSADVWS
D.melanogaster	STKGNMSEADAVQYIKKMEAQGRYSADVWS

**Figure 6**

atggccacgccgacgaaaacctacacgctagacgaaatcgagaagcacaacgaaaagtattctgcatgg  
**M** A T P T K T Y T L D E I E K H N E K Y S A W  
ttactaatccacaacgcagtgtagcagctgacgaaatgatggaagagcaccaggcggtgaagaagtt  
L L I H N A V Y D V T K F M E E H P G G E E V  
cttttgagcaggctggaagcatgcaactgaagcatttgaagatggttgacattccacagatgccaga  
L L E Q A G K H A T E A F E D V G H S T D A R  
gagttgatgaaacagtacaagattggtgatctttgtgaggaggaccagaagaaaatcggtcaggttgct  
E L M K Q Y K I G D L C E E D Q K K I G Q V A  
aagaaaactcagtgggcagctaccacctccaacgaaagctcctggatgagctggctgattcctggttga  
K K T Q W A A T T S N E S S W M S W L I P V G  
gtggcagctgctgctccattttgtaccgactcttctgtcctatggcgctcatcagtga  
V A A A A S I L Y R L F L S Y G A H Q -

**Figure 7**

H.sapiens	MAEQSDEAVKYITLLEI QKHNSKSTW L I L H H K V Y D L T K F L E E H P G G E E V L R E Q A G G D A T
R.microplus	----MATPTKTYTLDEIEKHNEKYSAWLLIHNNAVYDVTKFMEHHPGGEEVLLEQAGKHAT
D.melanogaster	---MSSEETKTFTRAEVAKHNTNKDTWLLIHNNIYDVTAFLNEHPGGEEVLIEQAGKDAT
An.gambiae	-----MSEVKTYSLADVKSHNTNKSTWIVIHNDIYDVTEFLNEHPGGEEVLLEQAGREAT
Cu.quinquefasciatus	----MSSEVKTYSLAEIKAHNTNKSSWIVIHNNIYDVTEFLNEHPGGEEVLLEQAGKEAT
	* :: : : ** . . : * : : : * . : * : * * : : * * * * * * * * * * * . * *

H.sapiens	ENFEDVGHSTDAREMSKTFIIGELHPDDRPKLNKPPE--TLITTIDSSSSWWTNHWVIPA
R.microplus	EAFEDVGHSTDARELMKQYKIGDLCEEDQKKIGQVAKKTQWAATTS-NESSWMSWLPVPG
D.melanogaster	ENFEDVGHSTNDARDMMKKYKIGELVESERTSVAQ-KSEPTWSTEQQTEESSVKSWLVPV
An.gambiae	EAFEDVGHSSDAREMMKFKVGE LIEAERKQIPV-KKEPDWKMDQQD-DNQLKQWIVPLI
Cu.quinquefasciatus	EAFEDVGHSTDAREMMKFKVGE LIESERKQVPV-KKEPDWSTEQKD-ENSLKSWIVPLI
	* * * * * * * . * * * : : * : : * * : : . . . . . * : : * * .

H.sapiens	SAVAVALMYRLYMAED---
R.microplus	VAAAASILYRLFLSYGAHQ
D.melanogaster	LCLVATLFYKFFFGALQ-
An.gambiae	LGLLATILYRFYFTQ----
Cu.quinquefasciatus	LGLLATIIRFYFTQ----
	. : : * : : : :
	-----