

Northumbria Research Link

Citation: Pound, Matthew and Salzmann, Ulrich (2016) Global vegetation distribution and terrestrial climate evolution at the Eocene-Oligocene transition. In: EGU General Assembly 2016, 17th - 22nd April 2016, Vienna, Austria.

URL: <http://meetingorganizer.copernicus.org/EGU2016/EGU...>
<<http://meetingorganizer.copernicus.org/EGU2016/EGU2016-2724.pdf>>

This version was downloaded from Northumbria Research Link:
<http://nrl.northumbria.ac.uk/27425/>

Northumbria University has developed Northumbria Research Link (NRL) to enable users to access the University's research output. Copyright © and moral rights for items on NRL are retained by the individual author(s) and/or other copyright owners. Single copies of full items can be reproduced, displayed or performed, and given to third parties in any format or medium for personal research or study, educational, or not-for-profit purposes without prior permission or charge, provided the authors, title and full bibliographic details are given, as well as a hyperlink and/or URL to the original metadata page. The content must not be changed in any way. Full items must not be sold commercially in any format or medium without formal permission of the copyright holder. The full policy is available online: <http://nrl.northumbria.ac.uk/policies.html>

This document may differ from the final, published version of the research and has been made available online in accordance with publisher policies. To read and/or cite from the published version of the research, please visit the publisher's website (a subscription may be required.)

www.northumbria.ac.uk/nrl





Global vegetation distribution and terrestrial climate evolution at the Eocene-Oligocene transition

Matthew Pound and Ulrich Salzmann

Northumbria University, Faculty of Engineering and Environment, Geography, Newcastle upon Tyne, United Kingdom
(ulrich.salzmann@northumbria.ac.uk)

The Eocene – Oligocene transition (EOT; ca. 34–33.5 Ma) is widely considered to be the biggest step in Cenozoic climate evolution. Geochemical marine records show both surface and bottom water cooling, associated with the expansion of Antarctic glaciers and a reduction in the atmospheric CO₂ concentration. However, the global response of the terrestrial biosphere to the EOT is less well understood and not uniform when comparing different regions. We present new global vegetation and terrestrial climate reconstructions of the Priabonian (late Eocene; 38–33.9 Ma) and Rupelian (early Oligocene; 33.9–28.45 Ma) by synthesising 215 pollen and spore localities. Using presence/absence data of pollen and spores with multivariate statistics has allowed the reconstruction of palaeo-biomes without relying on modern analogues. The reconstructed palaeo-biomes do not show the equator-ward shift at the EOT, which would be expected from a global cooling. Reconstructions of mean annual temperature, cold month mean temperature and warm month mean temperature do not show a global cooling of terrestrial climate across the EOT. Our new reconstructions differ from previous global syntheses by being based on an internally consistent statistically defined classification of palaeo-biomes and our terrestrial based climate reconstructions are in stark contrast to some marine based climate estimates. Our results raise new questions on the nature and extent of terrestrial global climate change at the EOT.