

Northumbria Research Link

Citation: Akram, Umair, Ellis, Jason, Myachykov, Andriy and Barclay, Nicola (2014) The mediating role of anxiety and depression on the association between perfectionism, sleep-disturbances, and insomnia severity. *Journal Of Sleep Research*, 23 (S1). p. 130. ISSN 1365-2869

Published by: Blackwell Publishing

URL: <http://dx.doi.org/10.1111/jsr.12213> <<http://dx.doi.org/10.1111/jsr.12213>>

This version was downloaded from Northumbria Research Link:
<http://nrl.northumbria.ac.uk/23597/>

Northumbria University has developed Northumbria Research Link (NRL) to enable users to access the University's research output. Copyright © and moral rights for items on NRL are retained by the individual author(s) and/or other copyright owners. Single copies of full items can be reproduced, displayed or performed, and given to third parties in any format or medium for personal research or study, educational, or not-for-profit purposes without prior permission or charge, provided the authors, title and full bibliographic details are given, as well as a hyperlink and/or URL to the original metadata page. The content must not be changed in any way. Full items must not be sold commercially in any format or medium without formal permission of the copyright holder. The full policy is available online: <http://nrl.northumbria.ac.uk/policies.html>

This document may differ from the final, published version of the research and has been made available online in accordance with publisher policies. To read and/or cite from the published version of the research, please visit the publisher's website (a subscription may be required.)

www.northumbria.ac.uk/nrl



See discussions, stats, and author profiles for this publication at: <http://www.researchgate.net/publication/265397357>

The mediating role of anxiety and depression on the association between perfectionism, sleep-disturbances, and insomnia severity

ARTICLE *in* JOURNAL OF SLEEP RESEARCH · SEPTEMBER 2014

Impact Factor: 2.95

4 AUTHORS:



[Umair Akram](#)

Northumbria University

6 PUBLICATIONS 3 CITATIONS

[SEE PROFILE](#)



[Jason Gordon Ellis](#)

Northumbria University

53 PUBLICATIONS 394 CITATIONS

[SEE PROFILE](#)



[Andriy Myachykov](#)

Northumbria University

28 PUBLICATIONS 72 CITATIONS

[SEE PROFILE](#)



[Nicola L Barclay](#)

Northumbria University

33 PUBLICATIONS 237 CITATIONS

[SEE PROFILE](#)

Sleep disorders – Insomnia 1

P440

Preliminary evidence for quality of life improvements after acceptance and commitment therapy in primary insomnia

E. Hertenstein, N. Thiel, M. Lueking, A. K. Kuelz, E. Schramm, C. Baglioni, K. Spiegelhalter, D. Riemann and C. Nissen
University Medical Center Freiburg, Freiburg, Germany

Background: Cognitive behavior therapy for insomnia (CBT-I) improves subjective and objective parameters of sleep in patients with primary insomnia (PI). However, the perception of poor sleep and an impaired quality of Life (QoL) often persists despite treatment. This study aimed to provide first evidence that Acceptance and Commitment Therapy (ACT), an intervention including mindfulness and value-based changes of behavior, can improve subjective sleep quality and QoL in patients with chronic PI.

Methods: Eleven patients with chronic PI who were non- or partial responders to CBT-I were included. Data were collected 6 weeks prior to the intervention (T_{-1}), directly before (T_0) and after the intervention (T_1), and at 3-months follow-up (T_2). The intervention consisted of six ACT sessions in an outpatient group setting. Primary outcomes were sleep-related QoL (Glasgow Sleep Impact Index), global QoL (World Health Organization Quality of Life scale) and subjective sleep quality (measured by sleep diaries).

Results: Ten patients completed the study, one dropped out due to scheduling problems. All measures remained stable between T_{-1} and T_0 . Significant improvements after the intervention were observed for sleep-related QoL, global QoL, and subjective sleep quality (ANOVA with factor Time, post-hoc contrasts T_1 and T_2 vs. T_0 , all P s < 0.05, large effect sizes). Subjective total sleep time, sleep onset latency and wake time after sleep onset did not significantly change across the study.

Conclusions: The findings provide preliminary evidence that ACT might improve subjective sleep quality and the QoL in patients with PI.

Disclosure: CN has received speaker honoraria from Servier. DR has received a consulting fee from Abbvie Germany. The other authors indicate no financial conflicts of interest.

P441

The mediating role of anxiety and depression on the association between perfectionism, sleep-disturbances, and insomnia severity

U. Akram, J. G. Ellis, A. Myachykov and N. L. Barclay
Northumbria Centre for Sleep Research, Northumbria University,
Newcastle upon Tyne, United Kingdom

Individuals with sleep-disturbances, specifically insomnia, often exhibit aspects of perfectionism, and symptoms of anxiety and depression, however the investigation of these factors together has been diminutive. The present study examined the mediating role of anxiety and depression on the associations between sleep-disturbances, insomnia-severity and aspects of perfectionism.

Eighty six participants (Age: $M = 25$, $SD = 10.02$; 80% female) were assessed using the Pittsburgh Sleep Quality Index (PSQI) and Insomnia Severity Index (ISI). Participants also completed two Multidimensional Perfectionism Scales (F-MPS; HF-MPS), and the Hospital Anxiety and Depression Scale (HADS).

Poor-sleepers demonstrated significantly higher scores on the F-MPS subscale *doubts-about-action* compared to normal-sleepers ($t(84) = 2.54$, $P = 0.01$). Additionally, global PSQI scores were signif-

icantly associated with *doubts-about-action* ($r = 0.22$, $P = 0.04$). Insomnia-severity was significantly associated with the F-MPS subscales of *doubts-about-action* ($r = 0.32$, $P = 0.01$), *concern-over-mistakes* ($r = 0.26$, $P = 0.02$), and *parental-criticism* ($r = 0.30$, $P = 0.01$). Hierarchical multiple-regression analyses, separately examining the global PSQI and insomnia-severity scores, demonstrated that when anxiety and depression were controlled for, none of the previous associations remained significant.

The results indicate that:

1. both increased sleep-disturbance and insomnia severity appear to be associated with increased *doubts-about-action*;
2. increased insomnia-severity also appears to be associated with increased *concern-over-mistakes*, and *parental criticism*; and
3. these associations are mediated by anxiety and depression.

These results could suggest that treatments for sleep-disturbance and insomnia should address anxiety and depressive symptoms, with the prospect of alleviating aspects of perfectionism, specifically *doubts-about-action*, *concern-over-mistakes*, and *parental criticism*, that may perpetuate pre-sleep arousal, consequently disturbing sleep.

Disclosure: Nothing to disclose.

P442

Sublingual flumazenil for the residual effects of hypnotics: zolpidem and brotizolam

N. Katz¹, G. Pillar², E. Peled² and N. Peled³

¹NICU, Wolfson Medical Center, Holon, ²Technion Faculty of Medicine, Haifa, ³Tel Aviv University Faculty of Medicine, Hod Hasharon, Israel

Background: Residual morning drowsiness is a frequent side effect of hypnotics which limits optimal treatment in Insomnia.

Purpose: To study the safety and efficacy of sublingual Flumazenil (GABA-A antidote) in reversing the residual hypnotic effect.

Methods: A double blind, placebo controlled, randomized cross over study with twenty healthy subjects who slept for 1.5 h following sleep induction (Zolpidem $n = 10$; Brotizolam $n = 10$). Upon awakening, they underwent neurocognitive tests including immediate word recall test (iWRT), Digit Symbol Substitution Test (DSST), and mood/performance questionnaires. They were then treated by Flumazenil or placebo ($n = 20;20$) and were re-evaluated after 20 and 60 min. A week later, the same procedures were performed (Placebo/Flumazenil accordingly).

Results: All 20 volunteers completed the study. Flumazenil was superior to placebo by 59–93% ($P < 0.05$ – 0.001) in improving performance in the various neurocognitive tasks. Subjects reported a significant improvement in vigilance with Flumazenil, both at 20 min (3.0 ± 0.6 cm vs. 1.3 ± 0.6 cm, $P < 0.02$) and 60 min (4.7 ± 0.6 cm vs. 2.8 ± 0.7 cm, $P < 0.03$). iWRT improved with Flumazenil vs. placebo at 20 min (4.2 ± 0.8 vs. 1.3 ± 0.9 words, $P < 0.005$) and 60 min (5.4 ± 1.1 vs. 1.2 ± 1.2 words, $P < 0.02$).

Conclusions: Sublingual administration of Flumazenil is safe and effective in reversing the hypnotic effects of Zolpidem and Brotizolam. This may offer a comprehensive and safer treatment for insomniac patients.