

# Northumbria Research Link

Citation: Gibbon, Karl, Debuse, Dorothee and Caplan, Nick (2013) Low impact weight-bearing exercise in an upright posture achieves greater lumbopelvic stability than overground walking. *Journal of Bodywork and Movement Therapies*, 17 (4). pp. 462-468. ISSN 1360-8592

Published by: Elsevier

URL: <http://dx.doi.org/10.1016/j.jbmt.2013.02.004>  
<<http://dx.doi.org/10.1016/j.jbmt.2013.02.004>>

This version was downloaded from Northumbria Research Link:  
<http://nrl.northumbria.ac.uk/11150/>

Northumbria University has developed Northumbria Research Link (NRL) to enable users to access the University's research output. Copyright © and moral rights for items on NRL are retained by the individual author(s) and/or other copyright owners. Single copies of full items can be reproduced, displayed or performed, and given to third parties in any format or medium for personal research or study, educational, or not-for-profit purposes without prior permission or charge, provided the authors, title and full bibliographic details are given, as well as a hyperlink and/or URL to the original metadata page. The content must not be changed in any way. Full items must not be sold commercially in any format or medium without formal permission of the copyright holder. The full policy is available online: <http://nrl.northumbria.ac.uk/policies.html>

This document may differ from the final, published version of the research and has been made available online in accordance with publisher policies. To read and/or cite from the published version of the research, please visit the publisher's website (a subscription may be required.)

[www.northumbria.ac.uk/nrl](http://www.northumbria.ac.uk/nrl)



1 **LOW IMPACT WEIGHT-BEARING EXERCISE IN AN UPRIGHT POSTURE**  
2 **ACHIEVES GREATER LUMBOPELVIC STABILITY THAN OVERGROUND**  
3 **WALKING**  
4

5  
6  
7 <sup>1</sup>Gibbon, K.C., <sup>1</sup>Debuse, D. & <sup>1</sup>Caplan, N.  
8

9  
10  
11 1 Faculty of Health & Life Sciences, Northumbria University, Newcastle upon  
12 Tyne, United Kingdom  
13

14  
15 Corresponding author:

16 Dr Nick Caplan

17 Faculty of Health and Life Sciences

18 Northumbria University

19 Northumberland Building

20 Newcastle upon Tyne

21 NE1 8ST

22 United Kingdom  
23

24 Tel: +44 191 243 7382

25 Email: [nick.caplan@northumbria.ac.uk](mailto:nick.caplan@northumbria.ac.uk)  
26  
27  
28

29

## **ABSTRACT**

30

31 The aim of this study was to determine the kinematic differences between  
32 movements on a new exercise device (EX) that promotes a stable trunk over a  
33 moving, unstable base of support, and overground walking (OW). Sixteen male  
34 participants performed EX and OW trials while their movements were tracked using  
35 a 3D motion capture system. Trunk and pelvis range of motion (ROM) were similar  
36 between EX and OW in the sagittal and frontal planes, and reduced for EX in the  
37 transverse plane. The pelvis was tilted anteriorly, on average, by about 16 degrees  
38 in EX compared to OW. Hip and knee ROM were reduced in EX compared to OW.  
39 The exercise device appears to promote similar or reduced lumbopelvic motion,  
40 compared to walking, which could contribute to more tonic activity of the local  
41 lumbopelvic musculature.

42

43 **Keywords:** kinematics, walking, lumbopelvic stability, exercise

## INTRODUCTION

44

45

46 *In vitro* studies have shown the thoracolumbar and lumbar spine, devoid of any  
47 musculature, will experience structural failure under compressive loadings as small  
48 as 20 and 90 N in magnitude, respectively (Crisco et al 1992). Considering spinal  
49 loadings experienced *in vivo* can range from 6 kN during selected everyday tasks  
50 (McGill & Norman 1986) to in excess of 36 kN during competitive powerlifting  
51 (Cholewicki et al 1991) the human vertebral column is intrinsically incapable of  
52 meeting the physiological demands placed upon it without additional stabilisation at a  
53 segmental level (Panjabi et al 1989).

54

55 The role of the lumbopelvic trunk musculature in providing the required  
56 supplementary stability at a segmental level is well documented (Bergmark 1989;  
57 Panjabi 1992; Cholewicki & McGill 1996; Vera-Garcia et al 2007). In particular, due  
58 to their anatomical positioning, morphology and function, the deeper fibres of the  
59 lumbar multifidus (LM) and the transversus abdominis (TrA) are considered crucial  
60 for local stability of the lumbar spine (Hodges & Richardson 1996; Hodges 1999; Kim  
61 et al 2007).

62

63 A growing body of evidence links structural and functional changes of local  
64 stabilising trunk muscles with low back pain (LBP) (Hides et al 1994; Hides et al  
65 1996; Hodges & Richardson 1996; Danneels et al 2000; Oddsson & De Luca 2003;  
66 Hides et al 2008; Hides et al 2008; MacDonald et al 2009; Teyhen et al 2009;  
67 Wallwork et al 2009). In people with LBP, muscle fibre atrophy and fatty infiltrations

68 of the LM have been observed (Kader et al 2000), as well as a dysfunction of the  
69 anticipatory activity of the LM and TrA (Hodges & Richardson 1998).

70

71 Corrective/restorative treatment strategies for such dysfunction of the local  
72 lumbopelvic musculature have included specific motor control exercises (Hides et al  
73 2008), 'core stability' training, muscular strength and endurance training (Danneels  
74 et al 2001), aerobic exercise (Frost et al 1995) and the use of an unstable base of  
75 support (BOS) (Marshall & Murphy 2006), often in a tailored combination (Demoulin  
76 et al 2010). The majority of these approaches tend to show only modest  
77 effectiveness (Keller et al 2007; van Middelkoop et al 2010), possibly due to a lack of  
78 carry-over to functional day-to-day activities (Richardson & Hides 2004; Hodges &  
79 Cholewicki 2007).

80

81 Recently a new method promoting activation of LM and TrA has been proposed as  
82 an alternative to the current approaches for addressing local lumbopelvic muscle  
83 dysfunction (Debusse et al 2013). The users of the exercise device move their feet in  
84 a quasi-elliptical path in anti-phase against virtually no external resistance. The  
85 absence of external resistance creates the need for much greater motor control of  
86 the legs and pelvis, to control leg movement, whilst maintaining an upright trunk  
87 posture, than in conventional exercise devices. The exercise device was found to  
88 recruit LM and TrA to a greater extent than a range of control activities, including  
89 standing on the ground or on an unstable base of support and voluntary muscle  
90 contractions. The authors postulated that the method promotes a relatively stable  
91 lumbopelvic area during a functional lower limb movement and results in an  
92 automatic recruitment/activity of TrA and LM (Debusse et al 2013). Richardson and

93 Jull in their seminal paper of 1995 proposed that local muscles work tonically, as  
94 opposed to global muscles which tend to work phasically. This is widely accepted by  
95 other authors working in this field (for example Sahrman 2002; Hides 2004; Hides  
96 et al 2004; Hodges & Cholewicki 2007). Debusse et al (2013) imply that tonic muscle  
97 activity is likely to be responsible for the stable lumbopelvic region when using the  
98 exercise device. However, no information was provided on the lumbopelvic and  
99 lower limb kinematics of the user while exercising to identify how the exercise device  
100 promoted lumbopelvic stability and, thus, tonic muscle activity.

101

102 The aim of the current study was to compare lower limb, pelvic and trunk kinematics  
103 during the use of a newly developed exercise device (EX) and overground walking  
104 (OW), with a particular focus on the level of lumbopelvic stability in both activities.

105

106

107

108

## **METHOD**

109

### **Participants**

111 Sixteen healthy adult male volunteers (mean  $\pm$  SD age: 26.5  $\pm$  3.38 years, body  
112 mass: 82.158  $\pm$  7.21 kg, height: 1.78  $\pm$  0.05 m, and body mass index: 25.89  $\pm$  2.16  
113 kg·m<sup>-2</sup>) with no recent history of LBP, gait impairments, or other conditions affecting  
114 their ability to walk or exercise, agreed to participate in this study. Participants gave  
115 their fully informed written consent to take part. The study had received ethical  
116 approval from the Institutional Review Board prior to data collection.

117

118 Three-dimensional Motion Capture

119 Three-dimensional trajectories of 39 retro-reflective markers ( $\text{\O} = 14\text{mm}$ ) were  
120 captured at a sampling frequency of 200 Hz using a 12 camera near-infrared motion  
121 capture facility (MX T20, Vicon Motion Systems, Oxford, UK). Markers were placed  
122 in accordance with a standard full-body model (Plug-in-Gait, Vicon Motion Systems,  
123 Oxford, UK), which consists of a 15 segment rigid-linked model of the head, thorax,  
124 pelvis, and bilateral upper arms, forearms, hands, thighs, lower legs and feet. Only  
125 the segmental orientations of the thorax, pelvis, thighs and lower legs were  
126 subsequently used for analysis.

127

128 The motion capture system was calibrated before all testing sessions using a  
129 standard dynamic protocol, with a 5 marker calibration wand (Vicon Motion Systems,  
130 Oxford, UK). System calibration was accepted when the image error of all 12  
131 cameras was less than 0.2 mm.

132

133 Body mass, height and anthropometric measurements, including leg length (anterior  
134 superior iliac spine to medial malleolus), ankle widths and knee widths, necessary for  
135 the correct operation of the model used were taken in triplicate and the mean value  
136 used thereafter.

137

138 Experimental protocol

139 Participants completed an overground walking (OW) condition and a condition using  
140 the exercise device (EX – Figure 1) in a counterbalanced random order within a  
141 single session. In the OW condition participants were asked to walk along a level 7.5  
142 m walkway, instrumented with embedded force plates (OR6-7, AMTI, Watertown,

143 Massachusetts, USA), at a self-selected comfortable speed. Starting positions were  
144 adjusted individually to ensure that 'clean' foot contacts with the force plates could  
145 be achieved without direct targeting by the participant. A minimum of 10 trials were  
146 completed, before six trials - without evidence of targeting - were selected for  
147 subsequent analysis.

148

149 In the EX condition participants were given an initial five minute period to familiarise  
150 themselves with the exercise device. Following this, 30 seconds of trajectory data  
151 were captured during exercise in standing. Subsequently, six cycles were chosen at  
152 random for analysis. All participants were given standardised instructions on the  
153 correct use of the device emphasising the need for a 'slow controlled movement'  
154 whilst maintaining 'an upright posture' during each cycle.

155

#### 156 Data processing and reduction

157 Marker trajectories collected during OW and EX trials were reconstructed and  
158 processed within Vicon Nexus (1.7, Vicon Motion Systems, Oxford, UK). Lost or  
159 obscured trajectory segments were interpolated using a quintic-spline function for  
160 gaps less than or equal to 10 frames (0.05 s) or a pattern fill function for gaps less  
161 than 10 frames, which utilises the trajectory of a marker with a similar predicted  
162 displacement trajectory. Marker trajectories were then low pass filtered at 5 Hz using  
163 a fourth-order zero lag Butterworth filter (Saunders et al 2005).

164

165 Key "gait cycle" phases (stance and swing) were demarcated for both the OW and  
166 EX conditions using discrete gait cycle events. Heel strikes and toe offs during OW  
167 were detected using the vertical component of the ground reaction force obtained



168 from the force plates embedded flush with the walkway surface at the centre of the  
169 calibrated capture volume. When using the new exercise device, the feet remain in  
170 contact with the foot plates at all times during both stance and swing phase.  
171 Therefore, data collected during EX were divided into a stance and swing phase  
172 based on the trajectory of a marker placed on the front corner of the foot plate:  
173 stance was defined as the most anterior to the most posterior foot plate position, and  
174 swing was from the most posterior to most anterior foot plate position.

175

176 Three-dimensional angular displacements for the trunk (thorax with respect to [wrt]  
177 pelvis), pelvis (wrt the room, rather than a relative position between body segments),  
178 hip (pelvis wrt thigh) and knee (thigh wrt lower leg) were time normalised to cycle  
179 duration in 2% increments (51 data points from 0-100%) for the right sided cycles of  
180 both OW and EX conditions. Angular range of motion (ROM) was calculated as the  
181 maximum minus the minimum joint angle achieved within one cycle. This was done  
182 for each of the six trials and averaged within each participant, and then between all  
183 participants in both conditions. The mean angular position of each segment or joint  
184 was determined as the average of each angle throughout the gait cycle for OW and  
185 EX. The difference in mean angular positions, or offset, between OW and EX was  
186 calculated. Data for each variable were checked for normality of distribution using Q-  
187 Q and box plots. For variables that were normally distributed, paired samples t-tests  
188 were used to compare ROM and mean angular position between conditions with  
189 significance set at  $p < 0.05$ . For variables that were not normally distributed,  
190 Wilcoxon signed rank tests were instead used. Confidence intervals (95%) were  
191 also calculated for each pairwise comparison. All statistical analyses were  
192 performed using SPSS (version 19).

193

194

195

## **RESULTS**

196

### **Spatiotemporal characteristics**

198 All spatiotemporal data were normally distributed. Statistically significant differences  
199 were observed in all six spatiotemporal parameters (Table 1). The EX condition was  
200 characterised by reduction in cadence ( $t=21.220$ ,  $df=15$ ,  $p<0.001$ ), stride length  
201 ( $t=14.041$ ,  $df=15$ ,  $p<0.001$ ), stride duration ( $t=26.380$ ,  $df=15$ ,  $p<0.001$ ), speed  
202 ( $t=20.506$ ,  $df=15$ ,  $p<0.001$ ), and effective stance phase ( $t=15.354$ ,  $df=15$ ,  $p<0.001$ )  
203 compared to those observed during OW. Step width was significantly greater in the  
204 EX condition compared to OW ( $t=2.662$ ,  $df=15$ ,  $p<0.05$ ).

205

206

### **Kinematics**

208 All angular ROM data were normally distributed with the exception of the hip in the  
209 transverse plane. Angular ROM was found to be similar between EX and OW  
210 conditions for the trunk in the sagittal ( $t=1.622$ ,  $df=15$ ,  $p=0.126$ ) and frontal ( $t=1.203$ ,  
211  $df=15$ ,  $p=0.248$ ) planes, and was similar for the pelvis in the sagittal ( $t=1.607$ ,  $df=15$ ,  
212  $p=0.129$ ) and frontal ( $t=0.213$ ,  $df=15$ ,  $p=0.834$ ) planes. In the transverse plane,  
213 ROM was significantly reduced for the trunk ( $t=8.513$ ,  $df=15$ ,  $p<0.001$ ) and the  
214 difference approached significance in the pelvis ( $t=1.854$ ,  $df=15$ ,  $p=0.083$ ) between  
215 EX and OW (Table 2).

216

217 All mean angular position data were normally distributed with the exception of the  
218 pelvis and hip in the transverse plane. The pelvis was significantly tilted anteriorly  
219 for the EX condition compared to OW with an offset of  $6.49^\circ$  ( $t=4.697$ ,  $df=15$   
220  $p<0.001$ ) (Table 3). Hip ROM was significantly reduced in the EX condition  
221 compared to OW in the sagittal ( $t=7.359$ ,  $df=15$ ,  $p<0.001$ ), frontal ( $t=2.572$ ,  $df=15$ ,  
222  $p=0.021$ ) and transverse ( $Z=3.516$ ,  $p<0.001$ ) planes (Table 2). Knee ROM was also  
223 reduced in EX in the sagittal ( $t=8.463$ ,  $df=15$ ,  $p<0.001$ ), frontal ( $t=7.041$ ,  $df=15$ ,  
224  $p<0.001$ ) and transverse ( $t=7.120$ ,  $df=15$ ,  $p<0.001$ ) planes. The hip ( $t=13.297$ ,  
225  $df=15$ ,  $p<0.001$ ) and knee ( $t=19.878$ ,  $df=15$ ,  $p<0.001$ ) were both more flexed  
226 throughout the gait cycle in the EX condition than in OW, with offsets of  $22.31^\circ$  and  
227  $24.11^\circ$ , respectively, which were significant (Table 3). Despite the reduced ROM,  
228 peak knee and hip angles occurred at a similar point in the gait cycle for OW and EX  
229 (Figure 2).

230

231

232

## **DISCUSSION**

233

234 The aim of this investigation was to compare the kinematics of lower limb and trunk  
235 motion during the use of a newly developed exercise device (EX), and overground  
236 walking (OW). The key findings of this study were that the lumbopelvic region was at  
237 least as stable whilst exercising on the new exercise device as overground walking.  
238 In the transverse plane, reduced ROM was observed during EX compared to OW.  
239 This stable lumbopelvic region was achieved over a dynamically moving base of  
240 support, where the ROM of the knees and hips was lower in EX than in OW. All

241 spatiotemporal variables were significantly reduced in EX compared to OW,  
242 suggesting a slower, more controlled motion.

243

244 Trunk motion in the sagittal and frontal planes demonstrated similar ranges for both  
245 EX and OW. In the transverse plane, a reduced ROM was observed for EX  
246 suggesting increased lumbopelvic stability. Similar observations were made for the  
247 pelvis in terms of ROM, although in the transverse plane, a smaller reduction in  
248 range of motion was found for EX, with this reduction approaching statistical  
249 significance.

250

251 As a fundamental human activity, walking has previously been investigated as an  
252 intervention strategy in the treatment of LBP (Torstensen et al 1998; Joffe et al 2002;  
253 Taylor et al 2003; Mirovsky et al 2006). However, heterogeneity of study design and  
254 methodological quality have contributed to inconsistent findings (Hendrick et al  
255 2010). Of these studies only Torstensen et al. (1998) and Taylor et al. (2003) used  
256 walking independently, while Joffe et al. (2002) and Mirovsky et al. (2006) combined  
257 walking with bodyweight support and traction, respectively. Notwithstanding the lack  
258 of evidence supporting walking as an effective intervention strategy for low back  
259 pain, the movement itself, involving control of trunk and pelvis motion during lower  
260 limb movements, is known to contribute to recruitment of the TrA and LM (Saunders  
261 et al 2004; Saunders et al 2005). Importantly, walking tends to be advocated by  
262 health care professionals in line with recommendations that ordinary physical  
263 activities should be continued as much as possible in order to aid recovery from LBP  
264 and prevent long-term disability (van Tulder et al 2000).

265

266 Similarities observed in both trunk and pelvic ROM between EX and OW in the  
267 sagittal and frontal planes suggest that the exercise device may be similar to  
268 walking, in terms of enabling tonic recruitment of the local lumbopelvic muscles such  
269 as TrA and LM. Previously Saunders et al. (2004; 2005) reported tonic TrA but  
270 phasic LM activity at walking speeds comparable to those reported here. However,  
271 no data were presented describing changes in activity amplitude, if any, within each  
272 gait cycle. The phasic activity of LM previously reported during walking (Saunders et  
273 al 2004) could be a factor leading to the questionable effectiveness of walking as a  
274 successful intervention for LBP (Hendrick et al 2010). The reduced transverse ROM,  
275 and thus the inherently more tonic muscle actions, in EX compared with OW seen in  
276 the current study could further indicate facilitation of greater tonic activity of the local  
277 lumbopelvic muscles (Richardson & Jull 1995) when using the new exercise device  
278 than in overground walking. If this reduced axial rotation results in more tonic  
279 recruitment of LM at a segmental level, then this could lead to the exercise device  
280 being a more successful intervention for LBP than walking. Current research within  
281 our group is exploring differences in lumbopelvic muscle recruitment between the  
282 exercise device and walking using ultrasound imaging and electromyography. Future  
283 studies in symptomatic populations are required to examine the clinical effectiveness  
284 of the exercise device.

285

286 No angular offsets were found between EX and OW for the trunk or pelvic position in  
287 all three planes, with the exception of a greater degree of anterior tilt of the pelvis in  
288 the EX condition. Influences of anterior pelvic tilt (O'Sullivan et al 2006) and  
289 accompanying lordotic spinal posture (Claus et al 2009), similar in magnitude to that  
290 observed within this investigation, have previously been shown to recruit both the

291 superficial and deep fibres of the LM to approximately 30-40% of maximal voluntary  
292 isometric contraction capabilities, which is known to facilitate stabiliser muscle  
293 recruitment (McArdle et al 1991). Thus, this angular offset could be beneficial for the  
294 recruitment of the LM, provided care is taken to avoid over-recruitment of the  
295 superficial fibres of LM.

296

297 Hip and knee joints were more flexed throughout the gait cycle in EX than during  
298 OW. The increase in hip flexion was partly due to the angular definition being relative  
299 to a perpendicular axis of the pelvis. Therefore, the observed increase in anterior tilt  
300 creates a greater degree of flexion at the hip. The increased flexion of the knee  
301 throughout the gait cycle during EX are linked to the reduced stride length that was  
302 caused by the mechanical constraints of the device. By reducing stride length, the  
303 knee was unable to reach full extension during the stance phase of the gait cycle, as  
304 is normally seen during OW. What was apparent for knee and hip motion in the  
305 sagittal plane, was that the change in angle throughout the gait cycle showed a more  
306 sinusoidal pattern in EX compared to OW. This, more regular, movement pattern  
307 could contribute, to some extent, to more continuous/tonic muscle recruitment, a key  
308 training requirement of the local stabilising musculature (Richardson & Jull 1995).

309

310 There has been a drive, in recent years, for training interventions for the local  
311 muscles of the lumbopelvic region to be made more functional (Hodges 2011). A  
312 number of studies have brought into question the transferability of any training  
313 effects seen following less functional activities such as gym ball training where the  
314 base of support is simply unstable (Drake et al 2006). Debuse et al. (2013)  
315 demonstrated that the local lumbopelvic muscles were recruited to a greater extent

316 with lower limb movement and an unstable base of support than with standing still on  
317 an unstable base of support (i.e. no voluntary lower limb movement). While  
318 overground walking involves lower limb movement, it does not usually involve an  
319 unstable base of support. During exercising on the new device, the requirement to  
320 control the descent of the “front” leg by gradually unloading the “back” leg within  
321 each gait cycle may result in greater recruitment of the local lumbopelvic muscles  
322 than overground walking. Our ultrasound imaging studies will examine this aspect in  
323 greater detail.

324

325 This study has a number of limitations. It examined relative motion between the  
326 pelvis and trunk. In order to gain a better understanding of how the exercise device  
327 influences the kinematics of the lumbopelvic region, a more detailed model of the  
328 thoracic and lumbar spine is needed. This would enable vertebral motion to be  
329 evaluated at a segmental level. Participants were asked to walk at their preferred  
330 walking speed. Due to the nature of the exercise device, movements were slower  
331 compared to walking. Saunders et al (2005) reported reduced axial rotation of the  
332 spine when walking slower. Thus, slow walking could lead to similar kinematics that  
333 were observed for the exercise device, and this should be explored further.  
334 However, walking slower does not involve an unstable base of support, or the  
335 complex motor control associated with using the exercise device, both of which could  
336 be contributing to increased local lumbopelvic muscle recruitment.

337

338

339

340

## **CONCLUSION**

341  
342  
343  
344  
345  
346  
347  
348  
349  
350  
351  
352

Key differences between exercising on the device and overground walking included reduced transverse plane range of trunk motion with respect to the pelvis (i.e. increased lumbopelvic stability), a more anteriorly tilted pelvis, and reduced stride length, knee and hip range of motion in the sagittal plane. The greater anterior tilt of the pelvis potentially moved the pelvis into a more advantageous position for the recruitment of TrA and LM. However, the unstable base of support afforded by the new exercise device would seem to add a challenge to movement control that may result in greater TrA and LM activity than overground walking. Future investigations should examine TrA and LM activity during walking and exercising on the new device using ultrasound imaging.



## REFERENCES

353

354

355 Bergmark A 1989 Stability of the lumbar spine. A study in mechanical engineering.

356 Acta Orthopaedica Scandinavica. Supplementum 230: 1-54.

357 Cholewicki J, McGill SM 1996 Mechanical stability of the in vivo lumbar spine:

358 implications for injury and chronic low back pain. Clinical Biomechanics 11: 1-

359 15.

360 Cholewicki J, McGill SM, Norman RW 1991 Lumbar spine loads during the lifting of

361 extremely heavy weights. Medicine and Science in Sports and Exercise 23:

362 1179-86.

363 Claus AP, Hides JA, Moseley GL, Hodges PW 2009 Different ways to balance the

364 spine: subtle changes in sagittal spinal curves affect regional muscle activity.

365 Spine 34: E208-14.

366 Crisco JJ, Panjabi MM, Yamamoto I, Oxland TR 1992 Euler Stability of the Human

367 Ligamentous Lumbar Spine Part II: Experiment. Clinical Biomechanics 7: 27-

368 32.

369 Danneels LA, Cools AM, Vanderstraeten GG, Cambier DC, Witvrouw EE, Bourgois

370 J, de Cuyper HJ 2001 The effects of three different training modalities on the

371 cross-sectional area of the paravertebral muscles. Scandinavian Journal of

372 Medicine and Science in Sports 11: 335-41.

373 Danneels LA, Vanderstraeten GG, Cambier DC, Witvrouw EE, De Cuyper HJ 2000

374 CT imaging of trunk muscles in chronic low back pain patients and healthy

375 control subjects. European Spine Journal 9: 266-72.

376 Debusse D, Birch O, St Clair Gibson A, Caplan N 2013 Low impact weight-bearing

377 exercise in an upright posture increases the activation of two key local

378 muscles of the pumbo-pelvic region. *Physiotherapy Theory and Practice* 29:  
379 51-60.

380 Demoulin C, Grosdent S, Capron L, Tomasella M, Somville PR, Crielaard JM,  
381 Vanderthommen M 2010 Effectiveness of a semi-intensive multidisciplinary  
382 outpatient rehabilitation program in chronic low back pain. *Joint, Bone, Spine :  
383 Revue du Rhumatisme* 77: 58-63.

384 Drake JDM, Fischer SL, Brown SHM, Callaghan JP 2006 Do Exercise Balls Provide  
385 a Training Advantage for Trunk Extensor Exercises? A Biomechanical  
386 Evaluation. *Journal of Manipulative and Physiological Therapeutics* 29: 354-  
387 362.

388 Frost H, Klaber Moffett JA, Moser JS, Fairbank JC 1995 Randomised controlled trial  
389 for evaluation of fitness programme for patients with chronic low back pain.  
390 *British Medical Journal* 310: 151-4.

391 Hendrick P, Te Wake A, Tikkisetty A, Wulff L, Yap C, Milosavljevic S 2010 The  
392 effectiveness of walking as an intervention for low back pain: a systematic  
393 review. *European Spine Journal* 19: 1613-1620.

394 Hides J 2004. Paraspinal mechanism in low back pain. In (Richardson C, Hodges P,  
395 Hides J). *Therapeutic exercise for lumbopelvic stabilization*. Churchill  
396 Livingstone, Edinburgh. 149-162.

397 Hides J, Gilmore C, Stanton W, Bohlscheid E 2008 Multifidus size and symmetry  
398 among chronic LBP and healthy asymptomatic subjects. *Manual Therapy* 13:  
399 43-9.

400 Hides J, Richardson C, Hodges P 2004. Local segmental control. In (Richardson C,  
401 Hodges P, Hides J). *Therapeutic Exercise for Lumbopelvic Stabilization*.  
402 Churchill Livingstone, Edinburgh. 185-220.

403 Hides J, Stanton W, Freke M, Wilson S, McMahon S, Richardson C 2008 MRI study  
404 of the size, symmetry and function of the trunk muscles among elite cricketers  
405 with and without low back pain. *British Journal of Sports Medicine* 42: 809-13.

406 Hides JA, Richardson CA, Jull GA 1996 Multifidus muscle recovery is not automatic  
407 after resolution of acute, first-episode low back pain. *Spine* 21: 2763-9.

408 Hides JA, Stanton WR, McMahon S, Sims K, Richardson CA 2008 Effect of  
409 stabilization training on multifidus muscle cross-sectional area among young  
410 elite cricketers with low back pain. *Journal of Orthopaedic and Sports Physical  
411 Therapy* 38: 101-8.

412 Hides JA, Stokes MJ, Saide M, Jull GA, Cooper DH 1994 Evidence of lumbar  
413 multifidus muscle wasting ipsilateral to symptoms in patients with  
414 acute/subacute low back pain. *Spine* 19: 165-72.

415 Hodges P, Cholewicki J 2007. Functional control of the spine. In (Vleeming A,  
416 Mooney V, Stoeckart R). *Movement, Stability and Lumbopelvic Pain*. Churchill  
417 Livingstone, Endinburgh.

418 Hodges PW 1999 Is there a role for transversus abdominis in lumbo-pelvic stability?  
419 *Manual Therapy* 4: 74-86.

420 Hodges PW 2011 Pain and motor control: From the laboratory to rehabilitation.  
421 *Journal of Electromyography and Kinesiology* 21: 220-8.

422 Hodges PW, Richardson CA 1996 Inefficient muscular stabilization of the lumbar  
423 spine associated with low back pain. A motor control evaluation of transversus  
424 abdominis. *Spine* 21: 2640-50.

425 Hodges PW, Richardson CA 1998 Delayed postural contraction of transversus  
426 abdominis in low back pain associated with movement of the lower limb.  
427 *Journal of Spinal Disorders* 11: 46-56.

428 Joffe D, Watkins M, Steiner L, Pfeifer BA 2002 Treadmill ambulation with partial body  
429 weight support for the treatment of low back and leg pain. *Journal of*  
430 *Orthopaedic and Sports Physical Therapy* 32: 202-13.

431 Kader DF, Wardlaw D, Smith FW 2000 Correlation between the MRI changes in the  
432 lumbar multifidus muscles and leg pain. *Clinical Radiology* 55: 145-9.

433 Keller A, Hayden J, Bombardier C, van Tulder M 2007 Effect sizes of non-surgical  
434 treatments of non-specific low-back pain. *European Spine Journal* 16: 1776-  
435 88.

436 Kim C, Gottschalk L, Eng C, Ward S, Lieber R 2007 The multifidus muscle is the  
437 strongest stabilizer of the lumbar spine. *The Spine Journal* 7: 76S.

438 MacDonald D, Moseley GL, Hodges PW 2009 Why do some patients keep hurting  
439 their back? Evidence of ongoing back muscle dysfunction during remission  
440 from recurrent back pain. *Pain* 142: 183-8.

441 Marshall PW, Murphy BA 2006 Evaluation of functional and neuromuscular changes  
442 after exercise rehabilitation for low back pain using a Swiss ball: a pilot study.  
443 *Journal of Manipulative and Physiological Therapeutics* 29: 550-60.

444 McArdle WD, Katch FI, Katch VL 1991. *Exercise physiology : energy, nutrition, and*  
445 *human performance.* Lea & Febiger, Philadelphia.

446 McGill SM, Norman RW 1986 Partitioning of the L4-L5 dynamic moment into disc,  
447 ligamentous, and muscular components during lifting. *Spine* 11: 666-78.

448 Mirovsky Y, Grober A, Blankstein A, Stabholz L 2006 The effect of ambulatory  
449 lumbar traction combined with treadmill on patients with chronic low back  
450 pain. *Journal of Back and Musculoskeletal Rehabilitation* 19: 73-78.

451 O'Sullivan PB, Dankaerts W, Burnett AF, Farrell GT, Jefford E, Naylor CS, O'Sullivan  
452 KJ 2006 Effect of different upright sitting postures on spinal-pelvic curvature  
453 and trunk muscle activation in a pain-free population. *Spine* 31: E707-12.

454 Oddsson LI, De Luca CJ 2003 Activation imbalances in lumbar spine muscles in the  
455 presence of chronic low back pain. *Journal of Applied Physiology* 94: 1410-  
456 20.

457 Panjabi M, Abumi K, Duranceau J, Oxland T 1989 Spinal stability and intersegmental  
458 muscle forces. A biomechanical model. *Spine* 14: 194-200.

459 Panjabi MM 1992 The stabilizing system of the spine. Part I. Function, dysfunction,  
460 adaptation, and enhancement. *Journal of Spinal Disorders* 5: 383-9.

461 Richardson C, Hides J 2004. Closed chain segmental control In (Richardson C,  
462 Hodges P, Hides J). *Therapeutic Exercise for Lumbopelvic Stabilization*.  
463 Churchill Livingstone, Edinburgh.

464 Richardson CA, Jull GA 1995 Muscle control-pain control. What exercises would you  
465 prescribe? *Manual Therapy* 1: 2-10.

466 Sahrmann S 2002. *Diagnosis and Treatment of Movement Impairment Syndromes*.  
467 Mosby, St Louis.

468 Saunders SW, Rath D, Hodges PW 2004 Postural and respiratory activation of the  
469 trunk muscles changes with mode and speed of locomotion. *Gait and Posture*  
470 20: 280-90.

471 Saunders SW, Schache A, Rath D, Hodges PW 2005 Changes in three dimensional  
472 lumbo-pelvic kinematics and trunk muscle activity with speed and mode of  
473 locomotion. *Clinical Biomechanics* 20: 784-93.

474 Taylor NF, Evans OM, Goldie PA 2003 The effect of walking faster on people with  
475 acute low back pain. *European Spine Journal* 12: 166-72.

476 Teyhen DS, Williamson JN, Carlson NH, Suttles ST, O'Laughlin SJ, Whittaker JL,  
477 Goffar SL, Childs JD 2009 Ultrasound characteristics of the deep abdominal  
478 muscles during the active straight leg raise test. Archives of Physical Medicine  
479 and Rehabilitation 90: 761-7.

480 Torstensen TA, Ljunggren AE, Meen HD, Odland E, Mowinckel P, Geijerstam S  
481 1998 Efficiency and costs of medical exercise therapy, conventional  
482 physiotherapy, and self-exercise in patients with chronic low back pain. A  
483 pragmatic, randomized, single-blinded, controlled trial with 1-year follow-up.  
484 Spine 23: 2616-24.

485 van Middelkoop M, Rubinstein SM, Verhagen AP, Ostelo RW, Koes BW, van Tulder  
486 MW 2010 Exercise therapy for chronic nonspecific low-back pain. Best  
487 Practice and Research. Clinical Rheumatology 24: 193-204.

488 van Tulder M, Malmivaara A, Esmail R, Koes B 2000 Exercise therapy for low back  
489 pain: a systematic review within the framework of the cochrane collaboration  
490 back review group. Spine 25: 2784-96.

491 Vera-Garcia FJ, Elvira JL, Brown SH, McGill SM 2007 Effects of abdominal  
492 stabilization maneuvers on the control of spine motion and stability against  
493 sudden trunk perturbations. Journal of Electromyography Kinesiology 17: 556-  
494 67.

495 Wallwork TL, Stanton WR, Freke M, Hides JA 2009 The effect of chronic low back  
496 pain on size and contraction of the lumbar multifidus muscle. Manual Therapy  
497 14: 496-500.

498

499

500 Table 1. Spatiotemporal characteristics of overground walking and exercise in the  
 501 standing position on the device. (SD = standard deviation, CI = confidence interval).

502

Gait Parameter	Overground Walking		Exercise Device		Mean Difference	
	Mean	±1SD	Mean	±1SD	(95% CI)	<i>P</i> value
<b>Cadence</b>						
(steps·min <sup>-1</sup> )	110.7	7.2	71.3	2.7	-39.4 (-43.4 to -35.45)	<0.001
Stride Length (m)	1.41	0.09	1.10	0.00	-0.31 (-0.35 to -0.26)	<0.001
Stride Duration (s)	1.09	0.07	1.69	0.06	0.60 (0.55 to 0.65)	<0.001
Speed (m·s <sup>-1</sup> )	1.30	0.13	0.65	0.03	-0.65 (-0.71 to -0.58)	<0.001
Step Width (m)	0.20	0.03	0.23	0.05	0.03 (0.01 to 0.06)	0.018
Stance Phase (%)	59.54	1.66	49.45	2.26	-10.09 (-11.49 to -8.69)	<0.001

503 Table 2. Angular range of motion of the trunk, pelvis, hip, and knee in all three  
 504 planes during overground walking and using the exercise device, also including the  
 505 mean difference between the two conditions. (SD = standard deviation, CI =  
 506 confidence interval).  
 507

Gait Parameter	Walking		Exercise Device		Mean Difference (95% CI)	P value
	Mean	±1SD	Mean	±1SD		
<b>Sagittal Plane</b>						
Trunk	3.93	1.80	3.01	1.67	-0.92 (-0.29 to 2.14)	0.126
Pelvis	2.89	0.78	3.69	1.91	0.8 (-1.86 to 0.26)	0.129
Hip	42.54	3.96	33.38	2.28	-9.16 (6.50 to 11.81)	<0.001
Knee	59.88	4.03	45.22	6.02	-14.66 (10.97 to 18.36)	<0.001
<b>Frontal Plane</b>						
Trunk	12.59	3.26	11.21	4.42	-1.39 (-1.07 to 3.84)	0.248
Pelvis	8.29	3.33	8.09	2.70	-0.20 (-1.85 to 2.26)	0.834
Hip	12.67	3.44	8.77	4.64	-3.90 (0.67 to 7.14)	0.021
Knee	16.50	5.91	9.42	5.22	-7.08 (4.93 to 9.22)	<0.001
<b>Transverse Plane</b>						
Trunk	12.55	3.85	3.92	1.14	-8.63 (6.47 to 10.79)	<0.001
Pelvis	12.00	3.28	9.25	4.18	-2.75 (-0.41 to 5.92)	0.083
Hip	16.93	7.34	8.87	2.73	-8.06 (4.95 to 11.17)	<0.001 <sup>a</sup>
Knee	20.66	5.37	10.59	3.96	10.07 (7.06 to 13.09)	<0.001

508 <sup>a</sup> indicates that these data were not normally distributed

509



510 Table 3. Mean angular position of the trunk, pelvis, hip and knee in all three planes  
 511 during overground walking and exercise in the standing position on the device. (SD  
 512 = standard deviation, CI = confidence interval).

513

Gait Parameter	Walking		Exercise Device		Mean Difference (±95% CI)	P value
	Mean	±1SD	Mean	±1SD		
Sagittal Plane						
Trunk	-5.37	6.15	-5.43	6.66	0.06 (-3.44 to 3.56)	0.970
Pelvis	9.06	4.06	15.55	6.18	-6.49 (-9.43 to -3.54)	<0.001
Hip	18.30	5.56	40.61	6.62	-22.31(-25.88 to -18.73)	<0.001
Knee	26.28	4.62	50.39	6.69	-24.11 (-26.69 to -21.52)	<0.001
Frontal Plane						
Trunk	-0.41	1.68	0.53	2.05	-0.94 (-2.27 to 0.38)	0.150
Pelvis	-0.25	1.23	-0.33	1.83	0.08 (-0.69 to 0.85)	0.827
Hip	-0.14	2.02	-0.91	2.33	0.77 (-0.13 to 1.67)	0.088
Knee	2.99	3.92	0.44	7.53	2.55 (-0.37 to 5.48)	0.082
Transverse Plane						
Trunk	-2.11	1.86	-1.70	2.03	-0.41 (-1.25 to 0.42)	0.311
Pelvis	-0.65	2.26	-1.63	2.93	0.98 (-0.05 to 2.01)	0.056 <sup>a</sup>
Hip	8.77	8.35	2.55	5.59	6.22 (1.77 to 10.66)	0.010 <sup>a</sup>
Knee	-8.77	9.14	1.16	8.58	-9.94 (-12.45 to -7.42)	<0.001

514 <sup>a</sup> indicates that these data were not normally distributed

515

516

517

## **FIGURE CAPTIONS**

518

519 Figure 1. The exercise device during use.


520

521 Figure 2. Hip (A) and knee (B) angles are presented for overground walking (- - -)

522 and exercise (—) conditions. The shaded region represents the standard deviation

523 for the exercise device data series.

524 **Figure 1**  
525  
526



527  
528  
529

