

Improving Tracheostomy Care in the United Kingdom: results of a guided quality improvement program in 20 diverse hospitals.

Brendan A McGrath^{1,2,*}, Sarah Wallace², James Lynch², Barbara Bonvento², Barry Coe², Anna Owen², Mike Firn³, Michael J Brenner⁴, Elizabeth Edwards⁵, Tracy L Finch⁶, Tanis Cameron⁷, Antony Narula⁸, David W Roberson⁹.

1. Manchester Academic Critical Care, Division of Infection, Immunity and Respiratory Medicine, School of Biological Sciences, Faculty of Biology, Medicine and Health, the University of Manchester, Manchester Academic Health Science Centre, UK.
2. Manchester University NHS Foundation Trust, Manchester, UK.
3. South West London and St George's Mental Health NHS Trust, Tooting, London.
4. University of Michigan, Ann Arbor, USA.
5. Patient Champion, Manchester, UK,
6. Department of Nursing, Midwifery and Health, The University of Northumberland, Newcastle-upon-Tyne
7. Tracheostomy Review and Management Service (TRAMS), Austin Health, Melbourne, Australia.
8. Imperial Healthcare, London
9. Bayhealth Medical Group, Milford, Delaware, US

*Corresponding author Brendan.mcgrath@manchester.ac.uk @BrendanAMcGrath

Presented in part at:

- BJA Research Forum, London, June 2019; abstracts:
 - T Liney, R Dawson, R Seth, J Lynch, S Wallace, B Bonvento, FK Ng, BA McGrath. Anxiety levels amongst patients with tracheostomies. *BJA* 2019;123(4):e504-e505
 - T Liney, R Dawson, R Seth, J Lynch, S Wallace, B Bonvento, FK Ng, BA McGrath. Patient satisfaction with tracheostomy care—a snapshot of UK practice. *BJA* 2019;123(4):e504-e505
 - FK Ng, BA McGrath, R Seth, J Lynch, S Wallace, B Bonvento, M Firn, T Finch. Measuring multidisciplinary staff engagement in a national tracheostomy quality improvement project using the NoMAD instrument. *BJA* 2019;123(4):e506
- World Congress of Intensive Care Medicine, Melbourne, October 2019
- World Airway Management Meeting, Amsterdam, November 2019
- NIAA/HSRC/ARS Anaesthesia Research 2019, York, December 2019

Running title: Improving Tracheostomy Care UK

Abstract

Background

Inconsistent and poorly coordinated systems of tracheostomy care commonly result in frustrations, delays, and harm. Quality improvement strategies described by exemplar hospitals of the Global Tracheostomy Collaborative have potential to mitigate such problems. This three-year guided implementation program investigated interventions designed to improve quality and safety of tracheostomy care.

Methods

The program management team guided implementation of 18 interventions over three phases (baseline/implementation/evaluation). Mixed methods interviews, focus groups, and **Hospital Anxiety and Depression** questionnaires defined outcome measures, with patient-level databases tracking and benchmarking **process metrics**. Appreciative Inquiry, interviews and **Normalisation Measure Development** questionnaires explored **change** barriers and enablers.

Results

All sites implemented at least 16/18 interventions, with the magnitude of some improvements linked to staff engagement (1536 questionnaires from 1019 staff). 2405 admissions (1868 ICU/HDU, 7.3% children) were prospectively captured. Median stay was 50 hospital days, 23 ICU days, and 28 tracheostomy days. **Incident severity score reduced significantly (n=606, p<0.01).** There were significant reductions in ICU (-0.25 days.month⁻¹), ventilator (-0.11 days.month⁻¹), tracheostomy (-0.35 days.month⁻¹) and hospital (-0.78 days.month⁻¹) **days** (all p<0.01). Time to first vocalisation and first oral intake both decreased by 7 days (n=733, p<0.01). Anxiety decreased by 44% (from 35.9% to 20.0%), and depression decreased by 55% (from 38.7% to 18.3%) (n=385, both p<0.01). Independent economic analysis demonstrated **£33 251** savings per patient, with projected annual **UK National Health Service** savings of £275 million.

Conclusions

This guided improvement program for tracheostomy patients significantly improved the quality and safety of care, contributing rich qualitative improvement data. Patient-centred outcomes were improved along with significant efficiency and cost savings across diverse UK hospitals.

Trial registry numbers IRAS-ID-206955, REC-Ref-16/LO/1196, NIHR Portfolio CPMS ID 31544

Keywords

Airway, Multidisciplinary, Quality Improvement, Safety, Tracheostomy

Introduction

Tracheostomies act as artificial airways for around 15 000 patients in England and Wales annually.¹⁻⁴ Patients often **have significant comorbidities, with medical needs that** cross traditional **specialty** working boundaries and locations. **These patients** are dependent on competent, knowledgeable **staff** to keep them safe. Landmark studies consistently highlight failings in **tracheostomy care provision in hospital, demonstrating how** inadequate staff training, **deficient** equipment provision and lack of **necessary** infrastructure lead to avoidable patient harm, morbidity and mortality.^{2 5-7} **Patients who undergo tracheostomy are often critically ill and have** in-hospital mortality reported from 25-60%, **with most of this mortality attributed to** underlying illness.^{8 9} However, up to 30% of tracheostomy patients experience an untoward incident **during their hospital stay**. Measurable harm occurs in 60-70% of such incidents, **including** hospital or Intensive Care Unit (ICU) (re)admission, prolonged in-patient stays, hypoxic brain injury **or** death.^{6 7} Delays in care are common due to the variety **and complexity** of services accessed by tracheostomised patients.¹⁰

Single hospitals or teams have previously reported success in improving outcomes, demonstrating that many problems in tracheostomy patients are amenable to prospective quality improvement (QI) strategies.¹¹⁻¹⁵ **In order to coordinate resources and strategies for such solutions at scale,** groups such as the UK National Tracheostomy Safety Project (www.tracheostomy.org.uk) and the Global Tracheostomy Collaborative (GTC, www.globaltrach.org) **have emerged,** providing **approaches** to improve care.

The GTC is a global community of **healthcare institutions, teams and individuals** focussed on collaborating to implement or expand upon best practices **that can improve the quality and/or safety of care.**¹⁶ **Multidisciplinary teams include members from the diverse specialities involved in tracheostomy care, and emphasise the central roles for patients, families and/or carers in decision-making and iterative improvement processes.**¹⁷ The GTC key drivers for improvement are described elsewhere,^{11 18} but briefly, comprise:

- **Multi-disciplinary care:** An institution-level multidisciplinary committee, and a multidisciplinary 'tracheostomy team' that meets and sees patients regularly.
- **Standardisation of care:** Planned protocols or care pathways.
- **Broad staff education**
- **Patient and family involvement**
- **Patient-level data:** To track changes, benchmark and drive improvements.

To date, only small-scale evaluations of adopting the GTC drivers for improvement and associated interventions have been reported, from individual sites or clusters of sites. Whilst a four-site UK implementation program positively impacted care, it remained unclear whether these interventions

could have a similar impact on patient outcomes at scale.¹² These patient outcomes include several widely used quality improvement metrics, such as mortality, adverse events, length of stay, and cost, as well as patient-centred measures most relevant to tracheostomy patients, such as time to first vocalisation, time to first oral intake, and measures of anxiety and depression. The aims of this study were; to conduct a large-scale demonstration program in geographically, demographically and politically diverse hospitals in the UK's public National Health Service (NHS); to refine existing interventions and evaluate their impact on safety and variation in care; and to understand the contextual implementation challenges for delivering reliable and sustainable change in patient outcomes. This study is also intended to share methods the readers can adapt to their own hospital and clinical practice. The *Improving Tracheostomy Care* program's key objectives were; to partner 20 UK hospitals, identifying leaders and champions from healthcare staff and patients; to rapidly implement GTC/National Tracheostomy Safety Project resources by creating a change culture; create a national collaborative environment for tracheostomy QI; and to describe and evaluate the experiences of patients and staff.

Methods

Study oversight

This investigator-initiated multicentre unblinded observational study was (competitively) funded by the Heath Foundation, in partnership with the Royal College of Anaesthetists, the NTSP and the GTC. **As part of the grant award independent** improvement consultancy was provided by Springfield Consultancy and independent economic evaluation by the University of East Anglia Health Economics Consulting. The study was designed by the authors and overseen by a representative steering committee.

Ethical considerations

The GTC has sought extensive advice in complying with country-specific Ethics Committee (Institutional Review Board) guidelines to fulfil its purpose as a QI Collaborative. There were clear additional aims for the **Improving Tracheostomy Care** program beyond the QI Collaborative, with detailed questioning, interviews and qualitative data collection from NHS patients and staff. National Research Ethics Committee approval was granted on 11th July 2016 (IRAS Project ID 206955, REC Ref 16/LO/1196), subsequently adopted onto the National Institute of Healthcare Research (NIHR) Portfolio (CPMS ID 31544).

Site selection

We identified **and contacted the** 44 potential UK hospitals from those with **prior active engagement** with either the **National Tracheostomy Safety Project** or GTC. **The first 20 sites who indicated a positive interest and multidisciplinary commitment to participating in the program, along with appropriate research capability and capacity, were included.** These 20 self-nominating sites represented the diverse nature of NHS hospitals, geographically, structurally and organisationally. Specifically, the hospitals spanned England, Wales and Scotland, included adult and paediatric district general and tertiary services with a range of tracheostomy services.

Interventions

Participating sites were grouped geographically, with study setup staggered over three months. An initial site visit by the study team profiled existing tracheostomy services and infrastructure. High-level executive engagement and support was secured, and local tracheostomy multi-disciplinary teams and leaders were identified, supported and/or developed. Interventions to improve care were identified from existing local practices, other participating sites, the wider GTC community, or newly developed to meet specific needs. Interventions were selected and prioritised by consensus processes previously described.¹⁹ Eighteen interventions were selected and grouped into themes addressing patient safety, patient-focussed quality of care, and organisational efficiency (**Table 1**). **There was no funding**

available for sites to develop or implement new interventions or services, although many developed internal business cases for new staff roles during the course of the program, supported by data generated by the project. GTC membership for all hospitals was paid for by the program along with funding for tracheostomy train-the-trainer and provider courses via the Advanced Life Support Group (ALSG, www.alsg.org) based on NTSP guidelines.^{17 18 20 21} Sites participated in 6-monthly themed national meetings and workshops with invited tracheostomy and QI expertise offering guidance on the content (tracheostomy care) and implementation (QI) elements of the program. Additional GTC webinars, meetings and forums were provided, along with peer support and guidance from fellow participants and the management team in order to promote a learning community around best practices. For example; if a particular site did not have an existing tracheostomy policy, competency standards or educational program, other sites were asked to provide not only their resources, but an explanation of how these resources had been developed and implemented. All meetings included strong patient and family representation. The number of interventions considered as 'fully implemented' (site **representatives'** opinion) was captured 6-monthly, so constructing aggregate implementation scores by site, intervention and time.

Data collection: patient-level

Patient-level data were entered by local staff into the GTC-specific REDCap database.^{19 22} Under Memoranda of Understanding and Data Sharing Agreements anonymous exports were provided for pooled analysis, with additional linked patient data made available from locally submitted critical care minimum datasets (CCMDS) and **local patient safety incident reporting**. Sites were provided with **data entry templates and examples**, and **regular** feedback was provided to encourage comprehensive data capture.

Recorded patient safety incidents were anonymised for site and date, then classified independently by three authors (AO, BC and BAM). A previously described harm score was applied,¹² summarised as: 0 No/minor physiological change (green); 1 Temporary harm (yellow); 2 Temporary harm with increased length of critical care or hospital stay (orange); 3 Permanent harm (red); 4 Intervention needed to sustain life (**dark red**); Reaction may have caused or contributed to death (black).

Each **adult** site was asked to recruit 10-20 patients or their families/carers, capturing experiences of tracheostomy care at three distinct phases; baseline (months 0-10), implementation (11-22) and evaluation (23-30) using HADS (Hospital Anxiety & Depression Scale) questionnaires. HADS consists of fourteen questions scored 0-3, with seven questions each focussing separately on anxiety and depression.^{20 23} Total scores of ≤ 7 are considered 'no case' (for depression/anxiety) in each category; 8-10 'borderline'; ≥ 11 'cases'. A free text field was included. Unstructured interviews conducted by local staff **guided by templates were offered to patient participants**, both providing more qualitative

narrative accounts. HADS is a standard questionnaire, validated for use by patients and families for assessing anxiety and depression.

Data collection: staff

Additionally, 10-20 frontline staff and site leads from all sites per phase completed Normalisation Measure Development (NoMAD) 'engagement' questionnaires, Appreciative Inquiry forms and semi-structured interviews. NoMAD is based on Normalization Process Theory and proposes four constructs (coherence, cognitive participation, collective action, and reflexive monitoring) addressing different aspects of implementing new practices.^{21 24} Answers are scored on a 5-point Likert scale bounded by 'Strongly agree' (score 5) through to 'Strongly disagree' (score 1). Staff could repeat surveys, but different staff were encouraged to participate, representing 'snapshots' of opinions. Appreciative Inquiry takes an action research approach that offers insight into positive and negative aspects of past, current and future practices and staff barriers and enablers. Appreciative Inquiry forms and interview questions are detailed elsewhere.¹⁹

Analysis

Data were pooled anonymously into Microsoft Excel, grouped by site, admission month (hence phase) and other discriminators. Simple descriptive statistics with means(SD) or median(IQR) values are reported as appropriate. Non-parametric linear regression investigated relationships between outcome and predictor variables, with confidence intervals for slopes based upon Kendall's τ constructed using StatsDirect 3.1.22 (StatsDirect Ltd). Cuzik's test identified trends in duration of care metrics.

After month 30, an additional 60 days of data collection occurred for length of stay (LoS). Patients not completing their hospital stay at this point had their LoS truncated. Sensitivity analyses investigated the impact; firstly, removing patients admitted during the first and last months; and secondly, removing patients with truncated LoS. ANOVA and Kruskal Wallis H tests were used to examine differences between groups using SPSS 22.0 (IBM Corp, New York, US). Fisher's exact test was used where data could be summarised into contingency tables. Cronbach's alpha (SPSS) evaluated reliability and consistency of the HADS and NoMAD tools in this setting, with an alpha of >0.80 representing good reliability.

For Qualitative Interviews and Appreciative Inquiry, thematic analysis was performed to identify, investigate, and report themes from the transcripts.^{22 25} Narratives were initially read line by line and coded into categories, without formal validation or double coding. Evaluation of large volumes of text was supported by NVivo 11 (QSR International); a qualitative data analysis software tool for coding and analysis of unstructured text. Codes were merged to develop themes representing participant experiences and perceptions.

Economic evaluation

Independent health economic evaluation was conducted to examine the cost-minimization associated with the implementation of the improvement strategies. The model specifically considered bed days and days of specific ICU organ support (CCMDS). Resource use was valued using the 2017/2018 NHS national schedule of reference cost.²⁶ Costs were calculated for Neonatal Intensive Care (NICU) days for infants, Paediatric Intensive Care (PICU) days for children, adult ICU days, and relevant ward days. The cost of care was calculated for each period (baseline, implementation, evaluation) and the incremental costs reported.

Results

Interventions

Sites had different baseline profiles with different interventions in place (**Figures 1-3**). Most sites took 12 months to start implementing substantial numbers of interventions. All sites made significant changes, with a median of 9 new interventions per site (range 4-13). Variation between sites in the number of implemented interventions reduced from a maximum difference of 9 to 2 items over the program. Sites had most difficulty implementing hospital-wide tracheostomy co-ordinators (eight sites unsuccessful) and ward-level tracheostomy link nurses (contact points between ward and hospital-wide specialist services; three sites unsuccessful). Patient champions (Q1) and patient-level data collection (Q7) were least likely to be implemented at baseline (two sites). Safety interventions appeared easiest to implement (group mean implementation score 89), followed by organisational (72.5) then quality interventions (71). A total of 371 staff attended national train-the-trainer days over the program, supported by 4000 local tracheostomy half-day training places.

Patient-level data

Hospital admissions were recorded from 1st August 2016 (month zero) to 31st January 2018 (month 30). Patient-level data were submitted from all sites with 2405 discrete patient admissions captured in the final combined database. A total of 1868 patients (77.7%) were admitted to ICU or HDU during their hospital stay, with detailed CCMDs data available for 1080. A total of 584 patients (24.3%) were admitted with existing tracheostomies and 177 patients (7.3%) were <16 years old (**Table 2 Supplemental**).

A total of 727 patient safety incidents were reported, with 26 considered non-clinical, leaving 701 incidents in 657 patients (27.3% of all patients). Fifty-eight patients experienced multiple incidents. **Table 3 Supplemental** describes incident categories, most commonly; accidental decannulation (18.4%), tube obstruction (10.4%), skin breakdown (7.8%) and bleeding (7.3%). There was a significant reduction over time in the severity score assigned to incidents by the blinded assessors in the 606 incidents reported with sufficient detail to assign a harm score (linear regression slope -0.044 (95% CI -0.034 to -0.055, ANOVA $p < 0.001$, **Figure 3**). Essentially, incidents occurred throughout, **but significantly fewer severe incidents occurred later in the program, with significantly less harm as a result**. Significantly more incidents occurred in paediatric admissions (47.5%) versus adult admissions (24.9%, $p < 0.001$). More patients experienced more than one incident in the paediatric group (17.9% vs 7.2%, $p < 0.001$).

The primary driver of cost reduction was the significant reduction in ICU and hospital length of stay associated with the guided improvement program. Considering all patients, there were significant reductions in tracheostomy time (equating to 0.35 days.month⁻¹ of program, $p < 0.001$), total hospital

LoS (0.8 days, $p<0.001$), ICU LoS (0.25 days, $p<0.001$) and ICU ventilator days (0.1 days, $p=0.002$) (**Figure 4**). These significant trends remained when sensitivity analyses were performed; firstly, removing 10 and 85 patients from the first and last months respectively; and secondly, removing the 33 patients in whom the final LoS was truncated.

The biggest LoS reductions were seen between the intervention and evaluation phases, mirroring the uptake of interventions and reflecting the time taken to establish new systems and treatment pathways. For the whole dataset, incremental costs between the baseline and evaluation periods translate into a cost saving per admission of £33,251 (£20,305 from ICU; £12,946 from wards). These cost savings do not account for GTC membership (£5,000 GBP per year) or the costs of new services, equipment, staff time for the program or staff posts; which varied considerably between sites.

A total of 385 consenting patients completed a HADS questionnaire following an in-patient admission (baseline $n=142$, implementation $n=128$, evaluation $n=114$). There was a 44.3% reduction in anxiety prevalence (decreasing from 54.2% to 37.4%, $p=0.008$) and a 52.7% reduction in depression prevalence (decreasing from 38.7% to 18.3%, $p<0.001$, **Table 4**). For the anxiety construct, 373 complete cases were analysed, producing a Cronbach's alpha of 0.86. For depression, 363 complete cases produced an alpha of 0.83. This represents good reliability of the HADS questionnaire in our setting.

Patients highlighted communication and oral nutritional intake as key areas of their care journey during baseline data collection; areas that the program's interventions could be expected to influence.^{26 27} Communication and nutritional metrics were available for 733 patient admissions (REDCap and additional local data). Time to cuff deflation decreased significantly over the three phases from a median of 17 to 10 days ($n=477$, $p<0.001$). Time to first use of a speaking valve with a ventilator decreased significantly from a median of 14 to 7 days ($n=199$, $p=0.037$), with clinically meaningful (but not statistically significant) reductions in time to speaking valve use with spontaneous ventilation (19 to 12 days, $n=204$, $p=0.77$). Time to first oral intake decreased significantly over the course of the program (26 to 9 days, $n=168$, $p<0.001$).

Staff data

At baseline, 204 Appreciative Inquiry forms (36 from leads, 168 from frontline staff) described quality concerns themed around harm, variation in practices, adequacy of training and safe staffing levels.¹⁹ Themes evolved during implementation (122 forms, 17 leads/105 frontline) and evaluation phases (125 forms, 18 leads/107 frontline) describing; positive improvements in education and training attendance; new collaborations resulting in better co-ordination of care; standardising or introducing new equipment or processes; the utility and more effective use of tracheostomy-specific data; a

perception of fewer patient safety incidents; the delivery of patient-centred care; and involvement of patients and families. Data collection burdens remained a prominent theme throughout.

Data was the dominant theme arising from 37 baseline site lead interviews, emphasising both opportunity and collection burdens. Concerns around staff training, resources and the challenges of multi-disciplinary relationships were also prominent with excitement around engaging in QI and raising the profile of tracheostomy care. Themes later evolved (from 22 lead interviews) demonstrating continued motivation supported by early local achievements, the enabling effect of the collaborative programme and rich testimony for the value of the programme in sharing strategies and driving improvements.

A total of 1019 unique participants (61.8% frontline) completed 1536 NoMAD forms, with over half declaring 3-10-years' experience caring for patients with tracheostomies. **Figure 6 Supplemental** shows overall mean construct scores increased significantly over the program by a mean difference of 0.26 ($p=0.02$). Stratifying sites into quintiles of aggregate NoMAD scores demonstrated significant differences in the rates of change in incident severity scores over the first 12 months of the program: slope coefficient -0.08 (a monthly reduction in incident severity) for the highest scoring (most engaged) quintile vs 0.02 (no reduction) for the lowest. There was good reliability for the four constructs (question groups) of the NoMAD questionnaire (coherence $\alpha=0.81$, cognitive participation $\alpha=0.85$, collective action $\alpha=0.80$, reflexive monitoring $\alpha=0.77$, general questions $\alpha=0.81$). When analysing all questions pooled together $\alpha=0.92$. This represents good to excellent reliability for NoMAD in this setting.^{27 28}

Discussion

This comprehensive program demonstrated that it is possible to significantly improve the quality and safety of tracheostomy care in a politically, geographically and operationally diverse group of UK NHS hospitals participating in a dedicated, guided quality improvement program. The views of patients and their families were actively sought and acted upon, designing, adopting and delivering innovative resources.^{17 26 27} Whilst difficulties were captured, meaningful change and improvements occurred in all sites (at different rates) reducing psychological distress associated with poor or less patient-focussed care. As expected, quality improvements led to organisational efficiencies, with motivated multidisciplinary teams acting proactively, decannulating patients appropriately and earlier, reducing tracheostomy days, and ICU and hospital lengths of stay. Qualitative data suggesting that the lower ICU admission rates observed towards the end of the program was primarily related to up-skilling of non-critical care locations and increasing staff confidence admitting or discharging to these locations.^{12 29 30} Our mixed methods research has captured a rich knowledge base for enabling change in this complex field which will be invaluable for future research and quality improvements.

Amongst interventions that appear most difficult to implement were a dedicated tracheostomy coordinator, link nurses on relevant wards, Speech & Language Therapists (SLTs) that can perform Fibreoptic Endoscopic Evaluation of Swallow (FEES) and patient champions. These organisational interventions had been shown by hospitals outside of the Improving Tracheostomy Care group to be effective methods of coordinating care, delivering efficient and effective proactive management.^{13–15 31–37} However, dedicated posts typically take 6-12 months to arrange and recruit to, perhaps longer to fund, and qualified and equipped SLTs take time to train.^{37 38} Commencing new services and embedding into practice can take years. Finding a suitable patient champion can also take time and the inclusion of a relevant patient in the team can be an unfamiliar experience to some, leading to barriers.^{11 39} Patient champions engaged in a number of core activities, including education, advocacy, strategy, and review of local material such as policies, information leaflets, care plans and care bundles, to ensure that all interventions remained as relevant and patient focussed as possible. Realising this value, all sites embedded patient champions by the end of the program, many of whom attended site lead meetings and actively participated in the group.

This program addressed a breadth of improvements aimed primarily at improving patient safety and the patient experience, which realised the anticipated associated improvements in organisational performance and therefore costs. This study was designed to build on the successes of smaller studies of tracheostomy QI in single sites and hospital clusters, but scaling up such initiatives does not guarantee success. The recent EPOCH study implemented a complex QI program for emergency laparotomy care in 93 UK hospitals,⁴⁰ comprising 37 component interventions, building on an evidence base of arguably weak, small-scale, before-and-after study designs; similar to the methodologies of

many of the studies underpinning this tracheostomy program. Whilst similarities exist between the findings of this study and EPOCH (high perceived data burden, limited staff time and limited resources dedicated to change management, in a complex patient population with around 20% 90-day mortality), our study was able to influence care, reduce variation and positively impact upon safety, quality and process measures. The smaller scale of the tracheostomy QI program and the self-selecting motivated cohort of hospital sites likely contributed to our positive outcomes. The national picture for tracheostomy care at baseline is also complex, fragmented, with unacceptably high rates of preventable harm and a lack of patient focus,² meaning that there may be more scope for positive change than the more evolved pathways of laparotomy care.

Whilst the majority of the site multidisciplinary teams expected to have knowledge of all tracheostomy in-patients, it is highly likely that not all admissions were recorded, and comprehensive outcome data was not collected for all. However, our pre-planned sensitivity analyses did not affect the observed reductions in duration of care metrics. The data burden for staff was perceived as high. In order to commend this program to the wider NHS and beyond, we recommend that contemporaneous electronic data capture systems that integrate with existing NHS systems are explored to ease burdens on staff.

Whilst this program has answered many questions, it has also identified many potential areas for future research. Further investigations should evaluate the impact of combinations of interventions on key outcomes, develop balanced score cards (providing a 'dashboard' of progress), develop ease impact matrices (guiding sites in balancing the difficulty of implementing a particular intervention with its potential impact), and continually develop resources for patients, families and staff wishing to embark on tracheostomy QI. The sites participating in this program were motivated, engaged and interested in tracheostomy care. This may not be the case in future sites, although this in itself may offer greater potential for improvement. We propose that future sites who are initially less engaged with improvement efforts may benefit from a tailored program starting with easy to implement interventions (bedhead signs for example) before building towards more difficult/complex interventions (such as multidisciplinary ward rounds). Our study did not measure the sustainability of change and impact beyond the program, but continued membership of the GTC will provide quarterly feedback and benchmarking to participating sites. This data tracking and feedback may help drive and sustain improvements.

This program is the first to demonstrate significant improvements in outcome measures developed in partnership with adult and paediatric tracheostomy patients at this scale. Improvements were seen in all sites in the domains of quality, safety and resultant organisational efficiency, translating into significant potential cost savings of around £275 million per year for the wider NHS. Importantly, we have learned what to do, how and when, contributing rich and deep new knowledge around making

changes in the NHS. We believe that these results will have a meaningful impact in the NHS and beyond.

Authors' contributions

Concept, design: BAM, SW, JL, BB, MF, TF, TC, AN DWR

Analysis and interpretation: BAM, SW, JL, BB, BC, AO, MF, TF

Drafting and revisions: All authors

Acknowledgements

The authors thank Prof Paul Dark, Dr Andrew Bentley, Dr Tim Felton, Dr Peter Alexander and Dr Malachy Columb for assistance in planning and statistical advice.

The authors are grateful to the Manchester Clinical Research Facility (Wythenshawe Hospital), NIHR and the Royal College of Anaesthetists for logistical support.

The authors acknowledge the support of the funder, The Health Foundation; an independent charity committed to better health & health care in the UK.

Study data were collected and managed using REDCap electronic data capture tools hosted at Vanderbilt University. REDCap (Research Electronic Data Capture) is a secure, web-based application designed to support data capture for research studies.

The authors wish to thank the staff who led and delivered the project at the individual sites, and to the wider support network consisting of the healthcare professionals, patients, families, committees and support staff of the Global Tracheostomy Collaborative.

In memory of our patient co-author Betty Edwards who sadly died during the preparation of this manuscript. As a critical care nurse and former patient with a tracheostomy, Betty had a unique insight into the problems and solutions highlighted in this program. She will be sadly missed.

Conflicts of interests

The authors declare that they have no conflicts of interest. For transparency: The following authors serve in varying capacities within the Global Tracheostomy Collaborative; David Roberson is the Founder and President; Erin Ward, Michael Brenner, Antony Narula, and Tanis Cameron serve as the Board of Directors; and Brendan McGrath and James Lynch serve as committee members. Brendan McGrath is also the Chair of the UK NTSP, European Lead of the GTC and National Clinical Advisor for Tracheostomy, NHS England.

Table 1. Interventions undertaken by sites, grouped into themes

- **Organisational Efficiency (6 items)**
 - O1 Implement a hospital steering group
 - O2 Ensure mandatory training for staff caring for tracheostomised patients
 - O3 Institute a hospital-wide tracheostomy policy
 - O4 Designated tracheostomy cohort wards
 - O5 Dedicated tracheostomy coordinator
 - O6 Tracheostomy link nurses in relevant wards

- **Patient-centred Quality of care interventions (7 items)**
 - Q1 Include patient champions
 - Q2 Implement Multidisciplinary Tracheostomy Team that sees patients
 - Q3 Integrate Speech & Language Therapist (SLTs) in ICU care
 - Q4 Involve SLT on Head & Neck wards
 - Q5 Involve SLT on general wards
 - Q6 Train SLTs to be Fiberoptic Endoscopic Evaluation of Swallow (FEES) proficient
 - Q7 Capture patient-level data (REDCap database)

- **Safety interventions (5)**
 - S1 Establish competency standards for staff caring for patients with tracheostomy
 - S2 Formalise MDT reviews of adverse incidents with learning
 - S3 Standardise bedside and ward area tracheostomy equipment
 - S4 Routinely place tracheostomy bedhead signs
 - S5 Use standardised tracheostomy care bundles

Supplemental Table 2. Patient descriptive statistics across the three phases. N/A: data not available.

Descriptive Statistics		Baseline	Intervention	Evaluation
N	2405 participants	656 participants	1178 participants	571 participants
Ethnicity	Caucasian	76.7%	78.9%	76.7
	African-American	4.0%	3.7%	4.0
	Asian	7.3%	7.3%	9.5
	Other	1.2%	1.3%	1.6
	N/A	10.8%	8.8%	8.2
Age (in years)	Mean	50.1	51.1	50.8
	Minimum	0.0	0.0	0.0
	Maximum	93.0	93.0	99.0
Sex	Male	60.2%	63.02%	61.8%
	Female	34.6%	33.82%	35.6%
	N/A	5.2%	3.22%	2.6%
Patients Category	Adult	91.0%	93.1%	92.5%
	Paediatric	8.7%	6.6%	7.2%
	N/A	0.3%	0.3%	0.3%
Tracheostomy planned prior to admission	Planned	9.0%	11.5%	10.9%
	Not Planned	8.2%	9.0%	10.7%
	N/A	82.8%	79.5%	78.5%
Existing tracheostomy present at admission	Yes	25.9%	22.6%	25.9%
	No	74.1%	77.4%	74.1%
	N/A	0.0%	0.0%	0.0%
Patient admitted to ICU	Yes	78.8%	79.1%	73.4%
	No	21.2%	20.5%	25.9%
	N/A	0%	0.3%	0.7%
Patient survive to hospital discharge	Yes	81.6%	81.8%	71.8%
	No	16.3%	14.2%	14.2%
	N/A	2.1%	4.0%	14.0%

Supplemental Table 3. Frequency of incidents occurring in patient groups. Individual patients may experience multiple incidents.

	Adult (n=2228, 93%)	Paediatric (n=177, 7%)	Total (n=2405, 100%)	% of all incidents
All incidents reported (incident count)	625	102	727	100%
Clinical incident reported (incident count)	619	82	701	96.4%
At least 1 incident reported (patient count)	554 / 2228 (24.9%)	84 / 177 (47.5%)	638 / 2405 (26.5%)	
				% of all clinical incidents
Accidental decannulation	105/619 (17.0%)	24/82 (29.3%)	129/701	18.4%
Tracheostomy tube obstruction	36/619 (5.8%)	37/82 (45.1%)	73/701	10.4%
Skin breakdown at tracheostomy site	47/619 (7.6%)	8/82 (9.8%)	55/701	7.8%
Significant bleeding from tracheostomy (>10mls fresh red blood)	50/619 (8.1%)	1/82 (1.2%)	51/701	7.3%
Failed decannulation (within 72hrs)	42/619 (6.8%)	4/82 (4.9%)	46/701	6.6%
Local skin or stoma infection/inflammation	45/619 (7.3%)	1/82 (1.2%)	46/701	6.6%
Air leak	41/619 (6.6%)	2/82 (2.4%)	43/701	6.1%
Laryngectomy patient - inadequate identification/provision	36/619 (5.8%)	0	36/701	5.1%
Communication between HCPs	30/619 (4.8%)	1/82 (1.2%)	31/701	4.4%
Delay in care	28/619 (4.5%)	0	28/701	4.0%
Tracheal injury (at insertion or later)	18/619 (2.9%)	0	18/701	2.6%
Infrastructure - No suitable bed	7/619 (1.1%)	0	7/701	1.0%
Infrastructure - staff knowledge	6/619 (1.0%)	0	6/701	0.9%
Tracheo-oesophageal fistula	6/619 (1.0%)	0	6/701	0.9%
Granuloma	1/619 (0.2%)	4/82 (4.9%)	5/701	0.7%
Infrastructure - Inadequate bedside equipment	5/619 (0.8%)	0	5/701	0.7%
Loss of airway	5/619 (0.8%)	0	5/701	0.7%
One-way valve used with cuff inflated	3/619 (0.5%)	0	3/701	0.4%
Chemical injury	2/619 (0.3%)	0	2/701	0.3%
Tracheo-cutaneous fistula	2/619 (0.3%)	0	2/701	0.3%
Illicit drug use by patient	1/619 (0.2%)	0	1/701	0.1%
Moving and handling (fall)	1/619 (0.2%)	0	1/701	0.1%
Tube adjustment	1/619 (0.2%)	0	1/701	0.1%
Insufficient details to classify further	103/619 (16.6%)	0	103/701	14.7%

Table 4. Breakdown of anxiety and depression cases over the three phases of the program from the HADS questionnaires (n=385).

		Phase of program							
		Baseline		Implementation		Evaluation		Change (baseline to evaluation)	
		Count	% of phase	Count	% of phase	Count	% of phase	% change	Fisher's exact p
Anxiety Classification	No anxiety case	65	45.8%	53	41.4%	72	62.6%	Reduction in anxiety cases	
	Borderline anxiety	26	18.3%	35	27.3%	20	17.4%	44.3% reduction (54.2% to 37.4%)	<0.01
	Anxiety case	51	35.9%	40	31.3%	23	20.0%		
Depression Classification	No depression case	63	44.4%	69	53.9%	81	70.4%	Reduction in depression cases	
	Borderline depression	24	16.9%	22	17.2%	13	11.3%	52.7% reduction (38.7% to 18.3%)	<0.01
	Depression case	55	38.7%	37	28.9%	21	18.3%		

1. McGrath BA, Wilkinson K. The NCEPOD study: on the right trach? lessons for the anaesthetist. *British Journal of Anaesthesia* 2015; **115**: 155–8
2. Martin IC, Freeth H, Kelly K, Mason M. NCEPOD: On the right Trach? [Internet]. A review of the care received by patients who underwent a tracheostomy. 2014. Available from: www.ncepod.org.uk/2014tc.htm
3. McGrath BA, Ramsaran R, Columb MO. Estimating the number of tracheostomies performed in critical care in England. *Br J Anaesth* 2012; **109**: 662P
4. McGrath BA, Wilkinson K, Shah RK. Notes from a Small Island: Lessons from the UK NCEPOD Tracheotomy Report. *Otolaryngology -- Head and Neck Surgery* 2015; **153**: 167–9
5. Thomas AN, McGrath BA. Patient safety incidents associated with airway devices in critical care: a review of reports to the UK National Patient Safety Agency. *Anaesthesia* 2009; **64**: 358–65
6. McGrath BA, Thomas AN. Patient safety incidents associated with tracheostomies occurring in hospital wards: a review of reports to the UK National Patient Safety Agency. *Postgrad Med J* 2010; **86**: 522–5
7. Cook TM, Woodall N, Harper J, Benger J. Major complications of airway management in the UK: results of the Fourth National Audit Project of the Royal College of Anaesthetists and the Difficult Airway Society. Part 2: intensive care and emergency departments†. *British Journal of Anaesthesia* 2011; **106**: 632–42
8. Halum SL, Ting JY, Plowman EK, et al. A multi-institutional analysis of tracheotomy complications. *Laryngoscope* 2012; **122**: 38–45
9. Shah RK, Lander L, Berry JG, Nussenbaum B, Merati A, Roberson DW. Tracheotomy outcomes and complications: a national perspective. *Laryngoscope* 2012; **122**: 25–9
10. Eibling DE, Roberson DW. Managing tracheotomy risk: time to look beyond hospital discharge. *Laryngoscope* 2012; **122**: 23–4
11. Bedwell JR, Pandian V, Roberson DW, McGrath BA, Cameron TS, Brenner MJ. Multidisciplinary Tracheostomy Care: How Collaboratives Drive Quality Improvement. *Otolaryngol Clin North Am* 2019; **52**: 135–47
12. McGrath BA, Lynch J, Bonvento B, et al. Evaluating the quality improvement impact of the Global Tracheostomy Collaborative in four diverse NHS hospitals. *BMJ Open Quality* 2017; **6**: [bmjqr.2017.022063](https://doi.org/10.1136/bmjqr.2017.022063)
13. Cameron TS, McKinstry A, Burt SK, et al. Outcomes of patients with spinal cord injury before and after introduction of an interdisciplinary tracheostomy team. *Crit Care Resusc* 2009; **11**: 14–9
14. Mirski MA, Pandian V, Bhatti N, et al. Safety, efficiency, and cost-effectiveness of a multidisciplinary percutaneous tracheostomy program. *Critical Care Medicine* 2012; **40**: 1827–34
15. Cetto R, Arora A, Hettige R, et al. Improving tracheostomy care: a prospective study of the multidisciplinary approach. *Clin Otolaryngol* 2011; **36**: 482–8
16. Roberson DW, Healy GB. The Global Tracheostomy Collaborative: Multidisciplinary quality improvement in tracheostomy care [Internet]. The Bulletin. 2017 [cited 2019 Dec 18]. Available from: <http://bulletin.facs.org/2017/09/the-global-tracheostomy-collaborative-multidisciplinary-quality-improvement-in-tracheostomy-care/>
17. McCormick ME, Ward E, Roberson DW, Shah RK, Stachler RJ, Brenner MJ. Life after

Tracheostomy: Patient and Family Perspectives on Teaching, Transitions, and Multidisciplinary Teams. *Otolaryngol Head Neck Surg* 2015; **153**: 914–20

18. Brenner MJ. GTC Global perspectives paper from Brenner et al. *British Journal of Anaesthesia* **TBC**
19. McGrath BA, Lynch J, Coe B, et al. Improving tracheostomy care: collaborative national consensus and prioritisation of quality improvements in the United Kingdom. 2018; 1–10
20. McGrath BA, Bates L, Atkinson D, Moore JA, National Tracheostomy Safety Project. Multidisciplinary guidelines for the management of tracheostomy and laryngectomy airway emergencies. *Anaesthesia* 2012; **67**: 1025–41
21. McGrath BA. *Comprehensive Tracheostomy Care: The National Tracheostomy Safety Project Manual*. Wiley; 2014.
22. Harris PA, Taylor R, Thielke R, Payne J, Gonzalez N, Conde JG. Research electronic data capture (REDCap)—A metadata-driven methodology and workflow process for providing translational research informatics support. *Journal of Biomedical Informatics* 2009; **42**: 377–81
23. Snaith RP, Zigmond AS. The hospital anxiety and depression scale. *Br Med J (Clin Res Ed)* 1986; **292**: 344
24. Finch TL, Girling M, May CR, et al. Improving the normalization of complex interventions: part 2 - validation of the NoMAD instrument for assessing implementation work based on normalization process theory (NPT). *BMC Med Res Methodol* 2018; **18**: 135
25. Braun V, Clarke V. Using thematic analysis in psychology. *Qualitative Research in Psychology* 2006; **3**: 77–101
26. NHS Improvement. NHS National schedule of reference cost [Internet]. 2018 [cited 2019 Oct 8]. Available from: <https://improvement.nhs.uk/resources/reference-costs/#rc1718>
27. McGrath B. By the patient, for the patient. Determining key quality of care measures for improving tracheostomy care. *Medical Research Archives* [Internet] 2019 [cited 2019 Dec 18]; **7** Available from: <https://journals.ke-i.org/index.php/mra/article/view/1989>
28. Ng FK, McGrath BA, Seth R, et al. Measuring multidisciplinary staff engagement in a national tracheostomy quality improvement project using the NoMAD instrument. *British Journal of Anaesthesia* 2019; **123**: e506
29. McGrath BA, Calder N, Laha S, et al. Reduction in harm from tracheostomy-related patient safety incidents following introduction of the National Tracheostomy Safety Project: our experience from two hundred and eighty-seven incidents. *Clin Otolaryngol* 2013; **38**: 541–5
30. Lavin J, Shah R, Greenlick H, Gaudreau P, Bedwell J. The Global Tracheostomy Collaborative: one institution's experience with a new quality improvement initiative. *Int J Pediatr Otorhinolaryngol* 2016; **80**: 106–8
31. Bonvento B, Wallace S, Lynch J, Coe B, McGrath BA. Role of the multidisciplinary team in the care of the tracheostomy patient. *JMDH* 2017; **Volume 10**: 391–8
32. McGrath BA, Wallace S. The UK National Tracheostomy Safety Project and the role of speech and language therapists. *Current Opinion in Otolaryngology & Head and Neck Surgery* 2014; **22**: 181–7
33. Pandian V, Miller CR, Schiavi AJ, et al. Utilization of a standardized tracheostomy capping

- and decannulation protocol to improve patient safety. *Laryngoscope* 2014; **124**: 1794–800
34. Pandian V, Miller CR, Mirski MA, et al. Multidisciplinary Team Approach in the Management of Tracheostomy Patients. *Otolaryngology -- Head and Neck Surgery* 2012; **147**: 684–91
35. Pandian V, Smith CP, Cole TK, et al. Optimizing Communication in Mechanically Ventilated Patients. *J Med Speech Lang Pathol* 2014; **21**: 309–18
36. Arora A, Hettige R, Ifeacho S, Narula A. Driving standards in tracheostomy care: a preliminary communication of the St Mary's ENT-led multi disciplinary team approach. *Clin Otolaryngol* 2008; **33**: 596–9
37. Hettige R, Arora A, Ifeacho S, Narula A. Improving tracheostomy management through design, implementation and prospective audit of a care bundle: how we do it. *Clin Otolaryngol* 2008; **33**: 488–91
38. [rcslt-position-statement-critical-care.pdf](https://www.rcslt.org/-/media/docs/clinical-guidance/rcslt-position-statement-critical-care.pdf?la=en&hash=42823C17957D4848818438CBCD5DC3998EF0CDF7) [Internet]. [cited 2019 Dec 18]. Available from: <https://www.rcslt.org/-/media/docs/clinical-guidance/rcslt-position-statement-critical-care.pdf?la=en&hash=42823C17957D4848818438CBCD5DC3998EF0CDF7>
39. Strychowsky JE, Albert D, Chan K, et al. International Pediatric Otolaryngology Group (IPOG) consensus recommendations: Routine peri-operative pediatric tracheotomy care. *Int J Pediatr Otorhinolaryngol* 2016; **86**: 250–5
40. Peden CJ, Stephens T, Martin G, et al. Effectiveness of a national quality improvement programme to improve survival after emergency abdominal surgery (EPOCH): a stepped-wedge cluster-randomised trial. *The Lancet* 2019; **393**: 2213–21