



# Direct dating of Neanderthal remains from the site of Vindija Cave and implications for the Middle to Upper Paleolithic transition

Thibaut Deviese<sup>a,1</sup>, Ivor Karvanić<sup>b,c</sup>, Daniel Comeskey<sup>a</sup>, Cara Kubiak<sup>a</sup>, Petra Korlević<sup>d</sup>, Mateja Hajdinjak<sup>d</sup>, Siniša Radović<sup>e</sup>, Noemi Procopio<sup>f</sup>, Michael Buckley<sup>f</sup>, Svante Pääbo<sup>d</sup>, and Tom Higham<sup>a</sup>

<sup>a</sup>Oxford Radiocarbon Accelerator Unit, Research Laboratory for Archaeology and the History of Art, University of Oxford, Oxford OX1 3QY, United Kingdom; <sup>b</sup>Department of Archaeology, Faculty of Humanities and Social Sciences, University of Zagreb, HR-10000 Zagreb, Croatia; <sup>c</sup>Department of Anthropology, University of Wyoming, Laramie, WY 82071; <sup>d</sup>Department of Evolutionary Genetics, Max-Planck-Institute for Evolutionary Anthropology, D-04103 Leipzig, Germany; <sup>e</sup>Institute for Quaternary Palaeontology and Geology, Croatian Academy of Sciences and Arts, HR-10000 Zagreb, Croatia; and <sup>f</sup>Manchester Institute of Biotechnology, University of Manchester, Manchester M1 7DN, United Kingdom

Edited by Richard G. Klein, Stanford University, Stanford, CA, and approved July 28, 2017 (received for review June 5, 2017)

Previous dating of the Vi-207 and Vi-208 Neanderthal remains from Vindija Cave (Croatia) led to the suggestion that Neanderthals survived there as recently as 28,000–29,000 B.P. Subsequent dating yielded older dates, interpreted as ages of at least ~32,500 B.P. We have redated these same specimens using an approach based on the extraction of the amino acid hydroxyproline, using preparative high-performance liquid chromatography (Prep-HPLC). This method is more efficient in eliminating modern contamination in the bone collagen. The revised dates are older than 40,000 B.P., suggesting the Vindija Neanderthals did not live more recently than others across Europe, and probably predate the arrival of anatomically modern humans in Eastern Europe. We applied zooarchaeology by mass spectrometry (ZooMS) to find additional hominin remains. We identified one bone that is Neanderthal, based on its mitochondrial DNA, and dated it directly to 46,200 ± 1,500 B.P. We also attempted to date six early Upper Paleolithic bone points from stratigraphic units G<sub>1</sub>, Fd/d+G<sub>1</sub> and Fd/d, Fd. One bone artifact gave a date of 29,500 ± 400 B.P., while the remainder yielded no collagen. We additionally dated animal bone samples from units G<sub>1</sub> and G<sub>1</sub>–G<sub>3</sub>. These dates suggest a co-occurrence of early Upper Paleolithic osseous artifacts, particularly split-based points, alongside the remains of Neanderthals is a result of postdepositional mixing, rather than an association between the two groups, although more work is required to show this definitively.

Vindija Cave (Croatia) | single-compound AMS dating | DNA analysis | zooarchaeology by mass spectrometry | Middle to Upper Paleolithic transition

The period between ~45,000 and 35,000 cal B.P. in Europe witnessed the so-called biocultural transition from the Middle to early Upper Paleolithic, when incoming anatomically modern humans displaced Neanderthal groups across the continent (1, 2). Significant questions still remain regarding the precise nature of this transition, the humans responsible for the various transitional early Upper Paleolithic industries, the degree of overlap between Neanderthals and modern humans, and the timing of the disappearance of the former. The European record for the transition retains its interest because it is the best-documented sequence for the disappearance of a hominin group available (3). The latest data, both radiometric and genetic, suggest Neanderthals and modern humans coexisted or overlapped for up to several thousand years in Europe until Neanderthal disappearance at around 40,000 cal B.P. (4, 5). Ascertaining the spatial attributes of Neanderthal and modern human populations in Europe is an area of active research, and a reliable chronology remains essential.

Our understanding of the biocultural processes involved in the transition have been greatly influenced by improved accelerator mass spectrometry (AMS) dating methods and their application

to directly dating the remains of late Neanderthals and early modern humans, as well as artifacts recovered from the sites they occupied. It has become clear that there have been major problems with dating reliability and accuracy across the Paleolithic in general, with studies highlighting issues with underestimation of the ages of different dated samples from previously analyzed sites (6). We have been working on redating some of the purported late-surviving Neanderthal sites from around Europe, which have included human and archaeological remains from sites such as Mezmaiskaya (Russia), where a previous directly dated Neanderthal infant yielded a radiocarbon age of ~29,000 B.P. (7), and Zafarraya (Spain), which was thought to contain Neanderthal remains clustering in age around a small group of U-series-dated animal bones between 33,400 and 28,900 B.P. (8). At Mezmaiskaya, the AMS dates obtained for the Neanderthal excavated above the previously dated individual were substantially older (9). This, along with other AMS dates from cut-marked fauna from the same archaeological horizons, suggested the original date of 29,000 B.P. could not be correct. At Zafarraya, Wood et al. (10) showed that, when redated using ultrafiltration methods, the bones that produced ages of ~33,000 B.P. were in fact beyond the radiocarbon limit, suggesting the

## Significance

Radiocarbon dating of Neanderthal remains recovered from Vindija Cave (Croatia) initially revealed surprisingly recent results: 28,000–29,000 B.P. This implied the remains could represent a late-surviving, refugial Neanderthal population and suggested they could have been responsible for producing some of the early Upper Paleolithic artefacts more usually produced by anatomically modern humans. This article presents revised radiocarbon dates of the human bones from this site obtained using a more robust purification method targeting the amino acid hydroxyproline. The data show that all the Neanderthal remains are from a much earlier period (>40,000 cal B.P.). These revised dates change our interpretation of this important site and demonstrate that the Vindija Neanderthals probably did not overlap temporally with early modern humans.

Author contributions: T.D., S.P., and T.H. designed research; T.D., I.K., D.C., C.K., P.K., M.H., S.R., N.P., M.B., S.P., and T.H. performed research; T.D., D.C., and T.H. contributed new reagents/analytic tools; T.D., I.K., D.C., C.K., P.K., M.H., S.R., N.P., M.B., S.P., and T.H. analyzed data; T.D. and T.H. wrote the paper; and I.K., D.C., C.K., P.K., M.H., S.R., N.P., M.B., and S.P. contributed to writing the paper.

The authors declare no conflict of interest.

This article is a PNAS Direct Submission.

Freely available online through the PNAS open access option.

<sup>1</sup>To whom correspondence should be addressed. Email: thibaut.deviese@rlaha.ox.ac.uk.

This article contains supporting information online at [www.pnas.org/lookup/suppl/doi:10.1073/pnas.1709235114/-DCSupplemental](http://www.pnas.org/lookup/suppl/doi:10.1073/pnas.1709235114/-DCSupplemental).

Neanderthal remains were unlikely to be as young as previously thought. In both cases, revised radiocarbon dates produced with more robust chemical pretreatment methods have illustrated significant underestimates in the previous dates that cannot be reconciled with a hypothesis of late-surviving refugial Neanderthals.

The Neanderthal fossil remains from level G<sub>1</sub> of Vindija Cave in northern Croatia have remained in the literature as potentially late individuals. Given the evidence from the Peštera cu Oase specimen, which demonstrates a recent Neanderthal ancestry in a 40,000 cal B.P. modern human from the Danube corridor (5), the renewed dating of the Vindija remains is overdue.

Two specimens, Vi-207 and Vi-208, were originally directly AMS dated in the late 1990s at the Oxford Radiocarbon Accelerator Unit (ORAU). Vi-207 is a right posterior mandible and Vi-208 is a parietal fragment, both showing Neanderthal-specific morphology (11, 12). The initial radiocarbon results were 29,080 ± 400 B.P. (OxA-8296) and 28,020 ± 360 B.P. (OxA-8295) (13). Higham et al. (14) attempted to redate these specimens by taking the very small amounts of collagen remaining from the original sample pretreatment and ultrafiltering the product before AMS dating. The revised measurements were 32,400 ± 1,800 B.P. (Vi-207: OxA-X-2089-07) and 32,400 ± 800 B.P. (Vi-208: OxA-X-2089-06), which indicated the previous dates were indeed too young. For sample Vi-208, after ultrafiltration, the C/N atomic ratio was 3.4, which indicates collagen of acceptable quality. However, for Vi-207, the >30-kDa fraction obtained produced a C/N ratio of 4.3, which indicates the presence of a high molecular weight contaminant. The radiocarbon date for this sample could therefore include a higher molecular weight noncollagenous contaminant, possibly cross-linked to the collagen. On the basis of the potential problems associated with the small size of the redated samples and the potential for remaining contaminants, OxA-X-2089-06 was considered to be a minimum age (14). If the dates are even approximately correct, however, it makes them the most recent known Neanderthals. This would imply a more extensive temporal overlap between Neanderthals and early modern humans in central Europe than has recently been documented (4).

In addition to the Neanderthal remains, level G<sub>1</sub> has yielded a small archaeological assemblage that contains techno-typologically Middle and Upper Paleolithic lithic artifacts plus several distinctively early Upper Paleolithic osseous points (12). It has been argued that the mix of Neanderthals, Middle Paleolithic tools, and Upper Paleolithic technology was the result of cryoturbation and *Ursus spelaeus* activity in level G<sub>1</sub>, with elements mixing into level G<sub>1</sub> from both the Upper Paleolithic unit F above and the Middle Paleolithic level G<sub>3</sub> below (15, 16). Zilhão (17) has suggested that the G<sub>1</sub> lithic assemblage has parallels with the Szeletian technocomplex, and further, that there is a mixture of elements of Szeletian and Aurignacian I and II within the level [see also

Svoboda (18)]. Karvanić and Smith (19) have suggested that the mixture of elements may represent the interaction and possible acculturation between modern humans and late Neanderthals. These alternatives are testable by selecting human and organic osseous points, as well as animal bones, for renewed AMS dating. This is what we have undertaken and describe here.

## Materials

A total of 10 samples from Vindija Cave were selected for AMS radiocarbon dating (Table 1). These included three previously dated Neanderthal specimens (Vi-207, Vi-208, and Vi-33.19), as well as a fourth Neanderthal bone (Vi-\*28) discovered using zooarchaeology by mass spectrometry (ZooMS) screening. In addition, to test the reality of the co-occurrence of earlier Upper Paleolithic bone and antler point artifacts with the Neanderthal remains, we selected six osseous points for dating (SI Appendix, Fig. S1). To test collagen preservation, we took ~3–5 mg bone powder, using tungsten carbide drills, and measured the %N content. This is an indicator of collagen preservation (20). The results show that only one bone point sample (Vi-3446) had sufficient levels of nitrogen to warrant full sampling for collagen extraction and AMS dating; the remainder failed and therefore were not sampled further (SI Appendix, Table S1). We also included the sample from a split-based bone point (Vi-3437) that had been analyzed in the laboratory previously, producing only a small collagen yield. We decided to attempt to redate it, using a larger starting mass of bone powder. Unfortunately, there was insufficient collagen remaining from this sample after pretreatment. We also selected four animal bones of *Cervidae* (Vi-\*17), *Panthera* sp. (Vi-\*6), *Ursus* sp. (Vi-\*7), and *Bovidae* (Vi-\*60), identified by ZooMS from stratigraphic units G<sub>1</sub> and G<sub>1</sub>–G<sub>3</sub> to explore the potential issue of postdepositional mixing in the deposits further.

## Results and Discussion


**ZooMS Collagen Fingerprinting.** We used ZooMS to identify potential hominin bone fragments among the unidentified faunal remains from the G<sub>1</sub> and G<sub>3</sub> levels, as well as the stratigraphic unit G<sub>1</sub>–G<sub>3</sub>. The majority of the 383 samples we analyzed yielded poor collagen preservation, which prevented any identification to genus or taxon. Only 101 samples produced identifiable spectra; a summary of all taxa identified by ZooMS is shown in SI Appendix, Table S2. This assemblage is dominated by *Ursus*, and only six of the 27 taxa identified by morphological study of the bones in Miracle et al. (21) could be identified here. We identified a single hominin specimen (Fig. 1 A and B), which again highlights the use of applying such techniques to groups of unidentified Paleolithic bone samples. The bone was analyzed using ancient DNA techniques to enable a formal species identification.

**DNA Analysis of the Human Bones.** Genomic analysis based on mitochondrial DNA revealed that all four human specimens fall into Neanderthal mitochondrial variation. Full mitochondrial genomes of Vi-207 and Vi-\*28 were reconstructed with an average coverage of 103-fold and 257-fold, respectively. The mitochondrial DNA sequence of Vi-207 was identical to Vi-33.25 and Feldhofer 1 mitochondrial genomes, whereas Vi-\*28 had an identical mitochondrial

**Table 1. Samples selected for AMS dating**

ORAU P Number	Sample Reference	Archeology Level	Species
39039	Vi-33.19 (SP2756)	G <sub>3</sub>	Neanderthal
41681	Vi-*28 (SP4162)	G <sub>1</sub>	Neanderthal
41415	Vi-208/Vi-11.29 (SP3563)	G <sub>1</sub>	Neanderthal
41416	Vi-207/Vi-11.41 (SP3562)	G <sub>1</sub>	Neanderthal
41417	Vi-3446	Fd/d+G1	Not known (point)
41687	Vi-3437	G <sub>1</sub>	Not known (split-based point)
41803	Vi-*17	G <sub>1</sub>	<i>Cervidae</i>
41801	Vi-*6	G <sub>1</sub>	<i>Panthera</i> sp
41802	Vi-*7	G <sub>1</sub>	<i>Ursus</i> sp
41804	Vi-*60	G <sub>1</sub> –G <sub>3</sub>	<i>Bovidae</i>

All samples produced enough collagen for radiocarbon dating with the exception of P41687, which failed to produce any collagen. See SI Appendix, Fig. S1 for images of the bone points included in this study.



**Fig. 1.** (A) High-resolution photographs of the Vi-\*28 Neanderthal bone found using ZooMS. The bone yields evidence for a probable cut and gauge marks (right upper part of the bone). The picture was taken after the bone had undergone sampling for ZooMS and before sampling for aDNA, radiocarbon, and stable isotope analysis. (B) MALDI-TOF mass spectrum of digested collagen from the Vi-\*28 bone. All tagged peaks (A, B, C, D, and F) denote sequence-matched peptides observed in human collagen (27, 28).

sequence to Vi-33.17 (*SI Appendix, Fig. S2*). Both Vi-33.25 and Vi-33.17 were found in layer I of Vindija Cave. As previously published, Vi-33.19 has the same mitochondrial sequence as Vi-33.16 (22). Because of lower endogenous DNA content in Vi-208, a full mitochondrial genome could not be reconstructed for the sample. However, from the limited amounts of mitochondrial sequences, we were able to trace most of the observed variants to variations found in previously sequenced Neanderthal mitochondrial genomes (*SI Appendix, Fig. S3*).

**AMS Dating.** Nine of the samples selected produced enough collagen (or hydroxyproline) to be dated by AMS. All dates obtained


on the four Neanderthal specimens at the ORAU are reported in Table 2. We also list the dates obtained on two other hominin samples: Vi-75-G3/h-203, analyzed at the Uppsala Radiocarbon Laboratory (Sweden) (23), and Vi-2291-18 (level G, sublayer unknown), prepared at the Max Planck Institute, Leipzig, and dated at the ORAU (24). The different sample pretreatments are also indicated in Table 2.

Vi-208 and Vi-207 produced hydroxyproline dates of  $42,700 \pm 1,600$  and  $43,900 \pm 2,000$  B.P., respectively. These ages are significantly older than any of the dates obtained previously for these specimens using the AG (gelatinized filtered collagen) and AF (ultrafiltered collagen) procedures, and this strongly suggests that

**Table 2. Radiocarbon dates of the Vindija Neanderthal remains**

P no.	OxA/OxA-X	CRA	±	P code	Used, mg	Yield, mg	%Yield	%C	$\delta^{13}\text{C}$ , ‰	$\delta^{15}\text{N}$ , ‰	C/N
Sample Vi-208 (SP3563)											
9663	8295	28,020	360	AG	233.9	15.2	6.5	37.1	-19.5	10.6	3.2
9663	2082-09	29,200	360	AG	229.9	10.9	4.7	42.7	-19.8	11.4	3.6
9663	2089-06	32,400	800	AF	n/a	n/a	n/a	42.3	-20.2	10.3	3.4
41415	X-2689-09	42,700	1,600	HYP	626.0	35.1	5.6	45.2	-26.1	10.2	5.6
Sample Vi-207 (SP3562)											
9665	8296	29,080	400	AG	229.2	9.7	1.5	36.6	-20.5	11.3	3.6
9665	2082-10	29,100	360	AG	128.8	6.7	5.2	41.6	-22.8	12.2	4.1
9665	2089-07	32,400	1,800	AF	n/a	n/a	n/a	39.0	-24.6	11.1	4.3
41416	X-2689-10	43,900	2,000	HYP	629.0	37.4	6.0	42.2	-24.7	11.5	5.6
Sample Vi-33.19 (SP2756)											
39039	32278	45,300	2,300	AF	560	62.2	11.1	45.2	-18.6	10.7	3.4
39039	X-2717-11	44,300	1,200	HYP	n/a	n/a	n/a	46.5	-24.4	12.3	4.9
Sample Vi-*28 (SP4162)											
41681	X-2687-57	46,200	1,500	HYP	1,000.0	54.0	5.4	34.3	-25.5	11.7	5.0
Sample Vi-2291-18 (level G, sublayer unknown)											
22966	V-2291-18	44,450	550	XB	/	/	/	39.3	-18.7	10.8	3.3
Sample Vi-75-G3/h-203											
/	Ua-13873	>42,000							-19.4	15.2	3.3

Radiocarbon dates of the Vindija Neanderthal remains. CRA is conventional radiocarbon age, expressed in years B.P. (49). Stable isotope ratios are expressed in per mil (‰) relative to Vienna Pee Dee Belemnite with a mass spectrometric precision of  $\pm 0.2\%$  (50). PCode refers to pretreatment code; AG is gelatinized filtered collagen, AF is ultrafiltered collagen, XB denotes collagen extracted at a different laboratory than the ORAU, and HYP denotes the extraction of hydroxyproline from hydrolyzed bone collagen (51, 52). Collagen yield represents the weight of gelatin or ultrafiltered gelatin in milligrams. %Yld is the percent yield of extracted collagen as a function of the starting weight of the bone analyzed. When further purification (AF or HYP) was performed on previously extracted collagen, the yields are not reported. %C is the carbon present in the combusted sample (gelatin or hydroxyproline). C/N is the atomic ratio of carbon to nitrogen and is acceptable if it ranges between 2.9–3.5 in the case of collagen or  $\sim 5.0$  in the case of hydroxyproline. All samples were prepared and dated at the ORAU with the exception of Vi-2291-18, which was extracted at the Max Planck Institute in Leipzig and dated in Oxford (24) and Vi-75-G3/h-203, which was prepared and dated at Uppsala in Sweden (23). The background carbon derived from the HPLC separation has been accounted for by using a correction where appropriate (details can be found in the *SI Appendix*). Slash indicates data not available.



**Fig. 2.** Bayesian age model showing the calibrated HYP ages of the four Neanderthal samples from Vindija Cave. The model is a simple phase model in OxCal 4.3 (47), in which all F14C determinations are assumed to have no relative order. “Start” and “End” correspond to the boundaries calculated by the model. The calibration curve of Reimer et al. (48) was used to calibrate the results. Details can be found in the [SI Appendix](#).

noncollagenous high molecular weight contaminants, probably crosslinked to the collagen, were still present in the sample previously dated. It is only by hydrolyzing the collagen and selecting the hydroxyproline that we were able to successfully remove these contaminants.

The AMS measurement of the third human bone from level G<sub>1</sub> (Vi-\*28), identified using the ZooMS method, gave a date of  $46,200 \pm 1,500$  B.P. For these three HYP (extraction of hydroxyproline from hydrolyzed bone collagen) dates obtained on the Neanderthal bones from level G<sub>1</sub>, we performed a  $\chi^2$  test using the modern fraction F14C and its error. The error weighted mean is  $0.0038 \pm 0.0005$  with a  $t$  value of 2.57. If  $t$  is  $<5.99$ , the value for  $\chi^2$ , the error weighted mean is not significant. This shows that the results obtained on the three Neanderthal bones from level G<sub>1</sub> are statistically in agreement. Vi-33.19 (level G<sub>3</sub>) was also dated using the hydroxyproline method and produced a date of  $44,300 \pm 1,200$  B.P., which is a more precise date than the one obtained using the AF procedure ( $45,300 \pm 2,300$  B.P.). The AF determination had a low target current in the AMS (85% of normal current) and was imprecise for this reason. The fact that the two dates overlap suggests no significant contamination in this bone. Taken together, these dates show a significantly older occupation of the site by Neanderthals, suggesting the site cannot be considered to be a refugium for late-surviving Neanderthals (Fig. 2, code in [SI Appendix](#)). The two other dates obtained on samples Vi-33.26 and Vi-75, even if obtained with a method that can be less efficient in removing contamination, confirm the dating of level G is more than 42,000 B.P.

Only one bone point (Vi-3446-P41417) produced enough nitrogen during prescreening to warrant pretreatment chemistry. Less than 500 mg bone was available, so we applied an AG (gelatin) treatment to maximize the collagen yield. We obtained a date of  $29,500 \pm 400$  B.P., which we consider to be a minimum age, as this pretreatment has been shown before to underestimate the age of ancient bones such as this (Table 3). To test possible perturbation in level G<sub>1</sub>, we also dated four faunal samples identified as originating from levels G<sub>1</sub> and G<sub>1</sub>–G<sub>3</sub>, using

ultrafiltration (AF). These samples produced dates that were all greater than or very close to the radiocarbon limit (Table 3). These samples are less likely to have been treated with conservation materials than the human bones, and therefore were not prepared using the HPLC method. Three of these faunal bones are from level G<sub>1</sub> (Table 1). We see more variation that would be expected statistically if we combine the three Neanderthal bones and three faunal bones from level G<sub>1</sub> for a  $\chi^2$  test (error weighted mean,  $0.0027 \pm 0.0004$ ;  $t = 22.52$ ;  $\chi^2 = 11.07$ ). This shows there is bone of variable radiocarbon age in the G<sub>1</sub> level and suggests the possibility of postdepositional mixing and movement of material. In consequence, the bone tools cannot be associated with the Neanderthal remains unless further direct radiocarbon dating using the HYP approach is undertaken. Clearly, the low collagen content of the points appears to preclude this at this time.

## Conclusions

Single-amino acid AMS dating of the Vindija Neanderthals has yielded results that are substantially older than the previous ages that were initially obtained. We have shown that the Neanderthals predate  $\sim 44,000$  cal B.P. The results suggest this group was not a late-surviving refugial Neanderthal population, as previously thought, and means the group almost certainly did not overlap with early anatomically modern humans in this part of Europe. Despite our best attempts, we were not able to date the bone industry associated with the archaeology of level G<sub>1</sub>. The one date we obtained from a later stratigraphic unit was younger than 30,000 B.P., but because the bone was not treated with the most rigorous pretreatment chemistry methods, it could potentially be older. The dating of other faunal materials from level G<sub>1</sub> highlighted a significant range in age, which could indicate a perturbation of the general sequence. The question, then, of whether some of the points could have been produced by Neanderthals remains open; however, it is parsimonious to conclude that the split-based point at least must have a maximum age of 32,000–34,000 B.P. based on evidence for its association with the Aurignacian in other regions, and so it likely postdates the Vindija Neanderthals significantly.

**Table 3. Radiocarbon determinations and analytical data from Vindija faunal remains and bone point**

OxA/OxA-X	CRA	±	PCode	Species	Used, mg	Yield, mg	%Yield	%C	$\delta^{13}\text{C}$ , ‰	$\delta^{15}\text{N}$ , ‰	C/N
34458	29,500	400	AG	Not known (bone point)	450	6.58	1.5	46.0	−19.4	3.5	3.4
2695-21	>46,700		AF	<i>Cervidae</i>	960	2.60	0.3	35.9	−20.7	5.6	3.3
34471	>48,400		AF	<i>Panthera</i> sp.	1,100	9.09	0.8	39.1	−18.9	10.1	3.2
34472	>49,000		AF	<i>Ursus</i> sp.	1,000	4.53	0.5	37.1	−23.0	8.8	3.2
34473	47,200	2,900	AF	<i>Bovidae</i>	950	40.41	4.3	42.2	−20.8	4.4	3.2

See [SI Appendix](#), [Table S2](#) caption for details of the terms used.

Bone points have been recovered throughout Eurasia with dates as early as ~37,000 B.P. (e.g., from the Aurignacian site of Pes'ko and the Châtelperronian site of Arcy-sur-Cure) (6, 25, 26). It is therefore clear that both anatomically modern humans and Neanderthals produced bone points, with only split-based bone/antler points being diagnostic of the earlier facies of the former.

Our perception of the biological transition between Neanderthals and modern humans has changed radically during the last decade. Evidence suggests interbreeding and a significant temporal overlap between the two from ~44,000–40,000 cal B.P. On the basis of our hydroxyproline dates and the DNA results, the Vindija Neanderthals date before the period when the first modern humans arrived into Europe and interbred with Neanderthals.

## Methods

**Bone Screening by ZooMS.** We used collagen peptide mass fingerprinting or ZooMS to analyze preserved type 1 collagen (COL1) from small unidentified bone fragments from the Vindija site. COL1 is characterized by three polypeptide (alpha) chains, and within these chains there are small differences in the amino acid sequences among different species. These offer the possibility of identification of species-specific amino acid markers. We used matrix-assisted laser desorption/ionization mass spectrometry to determine the relative mass-to-charge ( $m/z$ ) values of the tryptic digest peptides in each bone sample. By comparing these values with a library of known fauna, we identified bones to genus or species in some instances (27–31). We screened a total of 383 unidentified bone samples from Vindija Cave (levels G1, G3, and stratigraphic unit G1–G3).

**Radiocarbon Dating.** We used two different methods to prepare the samples for AMS dating. First, samples from artifacts were pretreated following the routine ORAU procedure, comprising a decalcification, base wash, reacidification, gelatinization, and ultrafiltration (coded AF in the ORAU), as described by Brock et al. (20). Sample P41417 produced too little collagen for ultrafiltration, so the treatment was halted after gelatinization (coded AG). The four human bone samples were dated using the single amino acid radiocarbon dating method developed at ORAU (32). This method involves separation of the underivatized amino acids from hydrolyzed bone collagen samples using preparative HPLC. Details of the method are described in the *SI Appendix*. Hydroxyproline (HYP), was isolated by Prep-HPLC, combusted, graphitized, and AMS-dated. This pretreatment approach is the most efficient technique to remove contaminants, including conservation materials (unless collagen-based glue has been applied).

## DNA Analysis.

**DNA extraction and library preparation.** Genomic analysis of the four human bones was performed at the Max Planck Institute for Evolutionary Anthropology in Leipzig, Germany. Three samples (Vi-207, Vi-208, and Vi-\*28) were

prepared specifically for this project, whereas Vi-33.19 was previously analyzed as part of a separate study (22). Bones were sampled using a sterile dentistry drill, and between 15 and 30 mg bone powder was used for DNA extraction, with the exception of Vi-\*28, in which three separate parts of the bone were sampled (*SI Appendix*, Table S3). DNA extraction was performed using a method for ultrashort DNA fragment retrieval (33) with modifications (34). The bone powder of Vi-207 and Vi-208 was treated with 0.5% hypochlorite solution before DNA extraction (34) to remove microbial and modern human DNA contamination. Next-generation sequencing libraries were prepared from 10  $\mu$ L of each extract, using two single-stranded library preparation schemes: Vi-207 and Vi-208 with a previously published method (35), with modifications from Korlević et al. (34), and Vi-\*28 with an automated library preparation scheme (36). Libraries were amplified and labeled with a pair of unique index sequences (37, 38).

**Enrichment of the mitochondrial DNA and sequencing.** Amplified libraries were enriched for human mitochondrial DNA, using a bead-based hybridization capture (39) with modifications (40). Enriched libraries were sequenced on an Illumina MiSeq platform with a double index configuration (41). Overlapping paired-end reads were merged (42) and mapped to the revised Cambridge Reference Sequence mitochondrial genome (rCRS; NC\_012092), using Burrows-Wheeler Aligner (43) with parameters “-n 0.01 -o 2 -l 16500” (44). Analyses were restricted to sequences with perfect index combinations. PCR duplicates were removed using bam-rmdup (<https://bitbucket.org/ustenzel/biohazard-tools>), and mapped sequences longer than 35 bases with a mapping quality of at least 25 were retained for subsequent analyses. To assess whether the results could be affected by present-day modern human contamination, sequences with ancient DNA specific cytosine to thymine substitutions (C to T) at the 5' or 3' molecule end were selected and analyzed separately.

**Phylogenetic analyses.** To reconstruct the mitochondrial genomes of the three previously unanalyzed hominin samples, DNA sequences were realigned to the Vindija 33.16 mitochondrial genome (AM948965) (45). Data were processed as described earlier. Data analysis was performed as previously published (ref. 28 for full mitochondrial genomes, ref. 46 for partial mitochondrial data).

**ACKNOWLEDGMENTS.** We thank Fred H. Smith (Illinois State University), Jadranka Mauch Lenardić (Institute for Quaternary Palaeontology and Geology, Croatian Academy of Sciences and Arts), and Pavao Rudan (Croatian Academy of Sciences and Arts) for their assistance in sampling and analyzing the Neanderthal and faunal bone from Vindija Cave. Ian Cartwright (Institute of Archaeology, University of Oxford) took the photographs of the Vi-\*28 Neanderthal bone. Rachel Hopkins ran the %N tests on some of the bones. We also thank the staff of the Oxford Radiocarbon Accelerator Unit at the University of Oxford and the Department of Evolutionary Genetics at the Max Planck Institute for Evolutionary Anthropology, Leipzig, for their support. The research leading to these results has received funding from the European Research Council (ERC) under the European Union's Seventh Framework Programme (FP7/2007-2013); ERC Grant 324139 “PalaeoChron” (to T.H.) and ERC Grant 694707 “100 Archaic Genomes” (to S.P.). We also acknowledge the support of the Royal Society for funding a University Research Fellowship (to M.B.).

- Mellars P (2006) A new radiocarbon revolution and the dispersal of modern humans in Eurasia. *Nature* 439:931–935.
- Hublin J-J (2015) The modern human colonization of western Eurasia: When and where? *Quat Sci Rev* 118:194–210.
- Mellars P (2004) Neanderthals and the modern human colonization of Europe. *Nature* 432:461–465.
- Higham T, et al. (2014) The timing and spatiotemporal patterning of Neanderthal disappearance. *Nature* 512:306–309.
- Fu Q, et al. (2015) An early modern human from Romania with a recent Neanderthal ancestor. *Nature* 524:216–219.
- Higham T (2011) European Middle and Upper Palaeolithic radiocarbon dates are often older than they look: Problems with previous dates and some remedies. *Antiquity* 85:235–249.
- Ovchinnikov IV, et al. (2000) Molecular analysis of Neanderthal DNA from the northern Caucasus. *Nature* 404:490–493.
- Hublin J-J, Barroso Ruiz C, Medina Lara P, Fontugne M, Reyss J-L (1995) The Mousterian site of Zafarraya (Andalucía, Spain): Dating and implications on the Palaeolithic peopling processes of western Europe. *C R Acad Sci II* 321:931–937.
- Pinhasi R, Higham TFG, Golovanova LV, Doronichev VB (2011) Revised age of late Neanderthal occupation and the end of the Middle Paleolithic in the northern Caucasus. *Proc Natl Acad Sci USA* 108:8611–8616.
- Wood RE, et al. (2013) Radiocarbon dating casts doubt on the late chronology of the Middle to Upper Palaeolithic transition in southern Iberia. *Proc Natl Acad Sci USA* 110:2781–2786.
- Wolpoff MH, Smith FH, Males M, Radović J, Rukavina D (1981) Upper Pleistocene human remains from Vindija cave, Croatia, Yugoslavia. *Am J Phys Anthropol* 54:499–545.
- Karavanić I, Smith FH (1998) The Middle/Upper Paleolithic interface and the relationship of Neanderthals and early modern humans in the Hrvatsko Zagorje, Croatia. *J Hum Evol* 34:223–248.
- Smith FH, Trinkaus E, Pettitt PB, Karavanić I, Pauunović M (1999) Direct radiocarbon dates for Vindija G(1) and Velika Pecina late Pleistocene hominid remains. *Proc Natl Acad Sci USA* 96:12281–12286.
- Higham T, Ramsey CB, Karavanić I, Smith FH, Trinkaus E (2006) Revised direct radiocarbon dating of the Vindija G1 Upper Paleolithic Neanderthals. *Proc Natl Acad Sci USA* 103:553–557.
- Kozłowski JK (1996) Cultural context of the last Neanderthals and early modern humans in central-eastern Europe. *The Lower and the Middle Palaeolithic*, eds Bar-Yosef O, Cavalli-Sforza LL, March RJ, Piperno M (International Union for Prehistoric and Protohistoric Sciences, Forlì, Italy), Vol 5, pp 205–218.
- Zilhao J, D'Errico F (1999) Reply, in the Neanderthal problem continued. *Curr Anthropol* 40:355–364.
- Zilhao J (2009) Szeletian, not Aurignacian: A review of the chronology and cultural associations of the Vindija G1 Neanderthals. *Sourcebook of Paleolithic Transitions: Methods, Theories, and Interpretations*, eds Camps M, Chauhan P (Springer New York, New York), pp 407–426.
- Svoboda J (2001) La question szélétienne. *Les industries a outils bifaciaux du paléolithique moyen d'Europe occidentale*, ed Cliquet D (Eraul, Liège, Belgium), pp 221–330.
- Karavanić I, Smith FH (2013) Alternative interpretations of the Middle/Upper Paleolithic interface at Vindija Cave (northwestern Croatia) in the context of central Europe and the Adriatic. *Archaeol Ethnol Anthropol Eurasia* 41:11–20.

20. Brock F, Higham T, Ramsey CB (2010) Pre-screening techniques for identification of samples suitable for radiocarbon dating of poorly preserved bones. *J Archaeol Sci* 37: 855–865.
21. Miracle PT, Mauch Lenardić J, Brajković D (2010) Last glacial climates, “Refugia”, and faunal change in southeastern Europe: Mammalian assemblages from Veternica, Velika pećina, and Vindija caves (Croatia). *Quat Int* 212:137–148.
22. Gansauge M-T, Meyer M (2014) Selective enrichment of damaged DNA molecules for ancient genome sequencing. *Genome Res* 24:1543–1549.
23. Krings M, et al. (2000) A view of Neandertal genetic diversity. *Nat Genet* 26:144–146.
24. Green RE, et al. (2010) A draft sequence of the Neandertal genome. *Science* 328: 710–722.
25. Davies W, Hedges REM (2008–2009) Dating a type site: Fitting Szeleta Cave into its regional chronometric context. *Praehistoria* 9–10:35–45.
26. Markó A (2013) Leaf-shaped lithic and osseous tools from old excavated cave sites: Demonstrating associations. *The Sound of Bones*, ed Lang F (Universität Salzburg, Salzburg, Austria), pp 191–202.
27. Buckley M, Collins M, Thomas-Oates J, Wilson JC (2009) Species identification by analysis of bone collagen using matrix-assisted laser desorption/ionisation time-of-flight mass spectrometry. *Rapid Commun Mass Spectrom* 23:3843–3854.
28. Brown S, et al. (2016) Identification of a new hominin bone from Denisova cave, Siberia using collagen fingerprinting and mitochondrial DNA analysis. *Sci Rep* 6:23559.
29. Richter KK, et al. (2011) Fish ‘n chips: ZooMS peptide mass fingerprinting in a 96 well plate format to identify fish bone fragments. *J Archaeol Sci* 38:1502–1510.
30. Buckley M, Gu M, Shameer S, Patel S, Chamberlain AT (2016) High-throughput collagen fingerprinting of intact microfaunal remains; a low-cost method for distinguishing between murine rodent bones. *Rapid Commun Mass Spectrom* 30: 805–812.
31. Buckley M, Harvey VL, Chamberlain AT (2017) Species identification and decay assessment of Late Pleistocene fragmentary vertebrate remains from Pin Hole cave (Creswell Crags, UK) using collagen fingerprinting. *Boreas* 46:402–411.
32. Marom A, McCullagh JSO, Higham TFG, Sinitsyn AA, Hedges REM (2012) Single amino acid radiocarbon dating of Upper Paleolithic modern humans. *Proc Natl Acad Sci USA* 109:6878–6881.
33. Dabney J, et al. (2013) Complete mitochondrial genome sequence of a Middle Pleistocene cave bear reconstructed from ultrashort DNA fragments. *Proc Natl Acad Sci USA* 110:15758–15763.
34. Korlević P, et al. (2015) Reducing microbial and human contamination in DNA extractions from ancient bones and teeth. *Biotechniques* 59:87–93.
35. Gansauge M-T, Meyer M (2013) Single-stranded DNA library preparation for the sequencing of ancient or damaged DNA. *Nat Protoc* 8:737–748.
36. Gansauge M-T, et al. (2017) Single-stranded DNA library preparation from highly degraded DNA using T4 DNA ligase. *Nucleic Acids Res* 45:e79.
37. Meyer M, Kircher M (2010) Illumina sequencing library preparation for highly multiplexed target capture and sequencing. *Cold Spring Harb Protoc*, 10.1101/pdb.prot5448.
38. Dabney J, Meyer M (2012) Length and GC-biases during sequencing library amplification: A comparison of various polymerase-buffer systems with ancient and modern DNA sequencing libraries. *Biotechniques* 52:87–94.
39. Maricic T, Whitten M, Pääbo S (2010) Multiplexed DNA sequence capture of mitochondrial genomes using PCR products. *PLoS One* 5:e14004.
40. Fu Q, et al. (2013) DNA analysis of an early modern human from Tianyuan cave, China. *Proc Natl Acad Sci USA* 110:2223–2227.
41. Kircher M, Sawyer S, Meyer M (2012) Double indexing overcomes inaccuracies in multiplex sequencing on the Illumina platform. *Nucleic Acids Res* 40:e3.
42. Renaud G, Stenzel U, Kelso J (2014) leehom: Adaptor trimming and merging for Illumina sequencing reads. *Nucleic Acids Res* 42:e141.
43. Li H, Durbin R (2009) Fast and accurate short read alignment with Burrows-Wheeler transform. *Bioinformatics* 25:1754–1760.
44. Meyer M, et al. (2012) A high-coverage genome sequence from an archaic Denisovan individual. *Science* 338:222–226.
45. Green RE, et al. (2008) A complete Neandertal mitochondrial genome sequence determined by high-throughput sequencing. *Cell* 134:416–426.
46. Welker F, et al. (2016) Palaeoproteomic evidence identifies archaic hominins associated with the Châtelperronian at the Grotte du Renne. *Proc Natl Acad Sci USA* 113: 11162–11167.
47. Bronk Ramsey C (2001) Development of the radiocarbon calibration program. *Radiocarbon* 43:355–363.
48. Reimer PJ, et al. (2013) IntCal13 and Marine13 radiocarbon age calibration curves 0–50,000 years cal BP. *Radiocarbon* 55:1869–1887.
49. Stuiver M, Polach HA (1977) Discussion: Reporting of C-14 data. *Radiocarbon* 19: 355–363.
50. Coplen TB (1994) Reporting of stable hydrogen, carbon, and oxygen isotopic abundances (technical report). *Pure Appl Chem* 66:273–276.
51. Brock F, Higham T, Ditchfield P, Bronk Ramsey C (2010) Current pretreatment methods for AMS radiocarbon dating at the Oxford Radiocarbon Accelerator Unit (ORAU). *Radiocarbon* 52:103–112.
52. Marom A, McCullagh JSO, Higham TFG, Hedges REM (2013) Hydroxyproline dating: Experiments on the <sup>14</sup>C analysis of contaminated and low-collagen bones. *Radiocarbon* 55:698–708.