



(12) **EUROPEAN PATENT SPECIFICATION**

(45) Date of publication and mention of the grant of the patent:  
**01.06.2005 Bulletin 2005/22**

(21) Application number: **99943085.3**

(22) Date of filing: **01.09.1999**

(51) Int Cl.7: **D06L 3/02**

(86) International application number:  
**PCT/GB1999/002878**

(87) International publication number:  
**WO 2000/012808 (09.03.2000 Gazette 2000/10)**

(54) **METHOD OF TREATING A TEXTILE**

VERFAHREN ZUR TEXTILBEHANDLUNG  
PROCEDE DE TRAITEMENT D'UN TEXTILE

(84) Designated Contracting States:  
**AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU  
MC NL PT SE**

(30) Priority: **01.09.1998 GB 9819046**  
**19.03.1999 GB 9906474**  
**01.04.1999 GB 9907713**

(43) Date of publication of application:  
**27.06.2001 Bulletin 2001/26**

(73) Proprietors:  
• **UNILEVER PLC**  
**London EC4P 4BQ (GB)**  
Designated Contracting States:  
**GB IE CY**  
• **UNILEVER N.V.**  
**3013 AL Rotterdam (NL)**  
Designated Contracting States:  
**AT BE CH DE DK ES FI FR GR IT LU NL SE PT  
MC LI**

(72) Inventors:  
• **APPEL, Adrianus, Cornelis, Maria**  
**NL-3068 GB Rotterdam (NL)**  
• **CARINA, Riccardo, Filippo**  
**1203 Geneve (CH)**  
• **DELROISSE, Michel Gilbert Jose**  
**Bromborough CH62 7FB (GB)**  
• **FERINGA, Bernard, Lucas**  
**NL-9765 EP Paterswolde (NL)**  
• **GIRERD, Jean-Jacques**  
**F-91190 Gif-sur-Yvette (FR)**  
• **HAGE, Ronald**  
**NL-2313 GD Leiden (NL)**

- **KALMEIJER, Robertus, Everardus**  
**NL-2803 VK Gouda (NL)**
- **MARTENS, Constantinus, Franciscus**  
**NL-3317 HG Dordrecht (NL)**
- **PEELEN, Jacobus, Carolina, Johannes**  
**NL-4926 RA Lage Zwaluwe (NL)**
- **QUE, Lawrence**  
**Roseville, MN 55113 (US)**
- **SWARTHOFF, Ton**  
**NL-3222 KC Hellevoetsluis (NL)**
- **TETARD, David**  
**Spital, Wirral CH63 9LS (GB)**
- **THORNTHWAITE, David**  
**Neston CH64 3SF (GB)**
- **TIWARI, Laxmikant**  
**Ellesmere Port S. Wirral CH65 4AN (GB)**
- **THIJSSSEN, Rob**  
**NL-5912 CA Venlo (NL)**
- **TWISKER, Robin, Stefan**  
**NL-2807 LN Gouda (NL)**
- **VEERMAN, Simon, Marinus**  
**NL-3067 EZ Rotterdam (NL)**
- **VAN DER VOET, Gerrit**  
**NL-3134 VR Vlaardingen (NL)**
- **SMITH, Richard, George**  
**Port Sunlight Merseyside CH62 5DT (GB)**

(74) Representative: **Elliott, Peter William**  
**Unilever PLC**  
**Unilever Intellectual Property Group**  
**Colworth House**  
**Sharnbrook Bedford, MK44 1LQ (GB)**

(56) References cited:  
**WO-A-95/27772**                      **WO-A-96/06154**  
**WO-A-97/07124**                      **DE-A- 19 714 122**

Note: Within nine months from the publication of the mention of the grant of the European patent, any person may give notice to the European Patent Office of opposition to the European patent granted. Notice of opposition shall be filed in a written reasoned statement. It shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

**Description**

5 [0001] This invention relates to a method of treating textiles such as laundry fabrics, more specifically to a method whereby bleaching by atmospheric oxygen is catalysed after the treatment. This invention also relates to textiles thus treated.

[0002] In a conventional bleaching treatment, a substrate such as a laundry fabric or other textile is contacted is subjected to hydrogen peroxide, or to substances which can generate hydroperoxyl radicals, such as inorganic or organic peroxides.

10 [0003] A preferred approach to generating hydroperoxyl bleach radicals is the use of inorganic peroxides coupled with organic precursor compounds. These systems are employed for many commercial laundry powders. For example, various European systems are based on tetraacetyl ethylenediamine (TAED) as the organic precursor coupled with sodium perborate or sodium percarbonate, whereas in the United States laundry bleach products are typically based on sodium nonanoyloxybenzenesulphonate (SNOBS) as the organic precursor coupled with sodium perborate. Alternatively, or additionally, hydrogen peroxide and peroxy systems can be activated by bleach catalysts, such as by complexes of iron and the ligand N4Py (*i.e.* N, N-bis(pyridin-2-yl-methyl)-bis(pyridin-2-yl)methylamine) disclosed in WO95/34628, or the ligand Tpen (*i.e.* N, N, N', N'-tetra(pyridin-2-yl-methyl)ethylenediamine) disclosed in WO97/48787. It has long been thought desirable to be able to use atmospheric oxygen (air) as the source for a bleaching species, as this would avoid the need for costly hydroperoxyl generating systems. Unfortunately, air as such is kinetically inert towards bleaching substrates and exhibits no bleaching ability. Recently some progress has been made in this area. For example, WO 97/38074 reports the use of air for oxidising stains on fabrics by bubbling air through an aqueous solution containing an aldehyde and a radical initiator, whereas according to WO95/34628 and WO97/48787 referred to above, molecular oxygen may be used as the oxidant with the iron catalysts, as an alternative to peroxide generating systems.

25 [0004] However, the known art teaches a bleaching effect only as long as the substrate is being subjected to the bleaching treatment. Thus, there is no expectation that hydrogen peroxide or peroxy bleach systems could continue to provide a bleaching effect on a treated substrate, such as a laundry fabric after washing and drying, since the bleaching species themselves or any activators necessary for the bleaching systems would be assumed to be removed from the substrate, or consumed or deactivated, on completing the wash cycle and drying.

30 [0005] It would be desirable to be able to treat a textile such that, after the treatment is completed, a bleaching effect is observed on the textile. Furthermore, it would be desirable to be able to provide a bleach treatment for textiles such as laundry fabrics whereby residual bleaching occurs when the treated fabric has been treated and is dry.

[0006] We have now found this can be achieved by a treatment method in accordance with the present invention, by catalysing bleaching of the substrate by atmospheric oxygen after treatment of the substrate.

[0007] Accordingly, the present invention provides a method according to claim 1.

35 [0008] The present invention further provides a dry textile having an organic substance as defined above applied or deposited thereon, whereby bleaching by atmospheric oxygen is catalysed on the textile.

[0009] Advantageously, by enabling a bleaching effect even after the textile has been treated, the benefits of bleaching can be prolonged on the textile. Furthermore, since a bleaching effect is conferred to the textile after the treatment, the treatment itself, such as a laundry wash cycle, may for example be shortened. Moreover, since a bleaching effect is achieved by atmospheric oxygen after treatment of the textile, hydrogen peroxide or peroxy-based bleach systems can be omitted from the treatment substance.

40 [0010] The organic substance may be contacted to the textile fabric in any suitable manner. For example, it may be applied in dry form, such as in powder form, or in a liquor that is then dried, for example as an aqueous spray-on fabric treatment fluid or a wash liquor for laundry cleaning, or a non-aqueous dry cleaning fluid or spray-on aerosol fluid. Other suitable means of contacting the organic substance to the textile may be used, as further explained below.

[0011] Any suitable textile that is susceptible to bleaching or one that one might wish to subject to bleaching may be used. Preferably the textile is a laundry fabric or garment.

45 [0012] In a preferred embodiment, the method according to the present invention is carried out on a laundry fabric using an aqueous treatment liquor. In particular, the treatment may be effected in a wash cycle for cleaning laundry. More preferably, the treatment is carried out in an aqueous detergent bleach wash liquid.

[0013] In a preferred embodiment, the treated textile is dried, by allowing it to dry under ambient temperature or at elevated temperatures.

50 [0014] The bleaching method may be carried out by simply leaving the substrate in contact with the organic substance for a sufficient period of time. Preferably, however, the organic substance is in an aqueous medium, and the aqueous medium on or containing the substrate is agitated.

55 [0015] The organic substance can be contacted to the textile fabric in any conventional manner. For example it may be applied in dry form, such as in powder form, or in a liquor that is then dried, for example in an aqueous spray-on fabric treatment fluid or a wash liquor for laundry cleaning, or a non-aqueous dry cleaning fluid or spray-on aerosol fluid.

[0016] In a preferred embodiment, the treated textile is dried, by allowing it to dry under ambient temperature or at elevated temperatures.

[0017] In a particularly preferred embodiment the method according to the present invention is carried out on a laundry fabric using aqueous treatment liquor. In particular the treatment may be effected in, or as an adjunct to, an essentially conventional wash cycle for cleaning laundry. More preferably, the treatment is carried out in an aqueous detergent wash liquor. The organic substance can be delivered into the wash liquor from a powder, granule, pellet, tablet, block, bar or other such solid form. The solid form can comprise a carrier, which can be particulate, sheet-like or comprise a three-dimensional object. The carrier can be dispersible or soluble in the wash liquor or may remain substantially intact. In other embodiments, the organic substance can be delivered into the wash liquor from a paste, gel or liquid concentrate.

[0018] The organic substance used in the method of the present invention makes use of atmospheric oxygen in its bleaching activity. This avoids the requirement that peroxygen bleaches and/or other relatively large quantities of reactive substances need be used in the treatment process. Consequently, only a relatively small quantity of bleach active substance need be employed and this allows dosage routes to be exploited which could previously not be used. Thus, while it is preferable to include the organic substance in a composition that is normally used in a washing process, such as a pre-treatment, main-wash, conditioning composition or ironing aid, other means for ensuring that the organic substance is present in the wash liquor may be envisaged.

[0019] For example, it is envisaged that the organic substance can be presented in the form of a body from which it is slowly released during the whole or part of the laundry process. Such release can occur over the course of a single wash or over the course of a plurality of washes. In the latter case it is envisaged that the organic substance can be released from a carrier substrate used in association with the wash process, e.g. from a body placed in the dispenser drawer of a washing machine, elsewhere in the delivery system or in the drum of the washing machine. When used in the drum of the washing machine the carrier can be freely moving or fixed relative to the drum. Such fixing can be achieved by mechanical means, for example by barbs that interact with the drum wall, or employ other forces, for example a magnetic force. The modification of a washing machine to provide for means to hold and retain such a carrier is envisaged similar means being known from the analogous art of toilet block manufacture. Freely moving carriers such as shuttles for dosage of surfactant materials and/or other detergent ingredients into the wash can comprise means for the release of the organic substance into the wash.

[0020] In the alternative, the organic substance can be presented in the form of a wash additive that preferably is soluble. The additive can take any of the physical forms used for wash additives, including powder, granule, pellet, sheet, tablet, block, bar or other such solid form or take the form of a paste, gel or liquid. Dosage of the additive can be unitary or in a quantity determined by the user. While it is envisaged that such additives can be used in the main washing cycle, the use of them in the conditioning or drying cycle is not hereby excluded.


[0021] The present invention is not limited to those circumstances in which a washing machine is employed, but can be applied where washing is performed in some alternative vessel. In these circumstances it is envisaged that the organic substance can be delivered by means of slow release from the bowl, bucket or other vessel which is being employed, or from any implement which is being employed, such as a brush, bat or dolly, or from any suitable applicator.

[0022] Suitable pre-treatment means for application of the organic substance to the textile material prior to the main wash include sprays, pens, roller-ball devices, bars, soft solid applicator sticks and impregnated cloths or cloths containing microcapsules. Such means are well known in the analogous art of deodorant application and/or in spot treatment of textiles. Similar means for application are employed in those embodiments where the organic substance is applied after the main washing and/or conditioning steps have been performed, e.g. prior to or after ironing or drying of the cloth. For example, the organic substance may be applied using tapes, sheets or sticking plasters coated or impregnated with the substance, or containing microcapsules of the substance. The organic substance may for example be incorporated into a drier sheet so as to be activated or released during a tumble-drier cycle, or the substance can be provided in an impregnated or microcapsule-containing sheet so as to be delivered to the textile when ironed.

[0023] The organic substance may comprise a preformed complex of a ligand and a transition metal. Alternatively, the organic substance may comprise a free ligand that complexes with a transition metal already present in the water or that complexes with a transition metal present in the substrate. The organic substance may also be included in the form of a composition of a free ligand or a transition metal-substitutable metal-ligand complex, and a source of transition metal, whereby the complex is formed *in situ* in the medium.

[0024] The organic substance forms a complex with one or more transition metals, in the latter case for example as a dinuclear complex. Suitable transition metals include for example: manganese in oxidation states II-V, iron I-IV, copper I-III, cobalt I-III, nickel I-III, chromium II-VII, silver I-II, titanium II-IV, tungsten IV-VI, palladium II, ruthenium II-V, vanadium II-V and molybdenum II-VI.


[0025] In a preferred embodiment, the organic substance forms a complex of the general formula (A1):



in which:


- 5 M represents a metal selected from Mn(II)-(III)-(IV)-(V), Cu(I)-(II)-(III), Fe(I)-(II)-(III)-(IV), Co(I)-(II)-(III), Ni(I)-(II)-(III), Cr(II)-(III)-(IV)-(V)-(VI)-(VII), Ti(II)-(III)-(IV), V(II)-(III)-(IV)-(V), Mo(II)-(III)-(IV)-(V)-(VI), W(IV)-(V)-(VI), Pd(II), Ru(II)-(III)-(IV)-(V) and Ag(I)-(II), and preferably selected from Mn(II)-(III)-(IV)-(V), Cu(I)-(II), Fe(II)-(III)-(IV) and Co(I)-(II)-(III);
- 10 L represents a ligand as herein defined, or its protonated or deprotonated analogue;
- X represents a coordinating species selected from any mono, bi or tri charged anions and any neutral molecules able to coordinate the metal in a mono, bi or tridentate manner, preferably selected from  $O^{2-}$ ,  $RBO_2^{2-}$ ,  $RCOO^-$ ,  $RCONR^-$ ,  $OH^-$ ,  $NO_3^-$ ,  $NO_2^-$ ,  $NO$ ,  $CO$ ,  $S^{2-}$ ,  $RS^-$ ,  $PO_3^{4-}$ , STP-derived anions,  $PO_3OR^{3-}$ ,  $H_2O$ ,  $CO_3^{2-}$ ,  $HCO_3^-$ ,  $ROH$ ,  $NRR'R''$ ,  $RCN$ ,  $Cl^-$ ,  $Br^-$ ,  $OCN^-$ ,  $SCN^-$ ,  $CN^-$ ,  $N_3^-$ ,  $F^-$ ,  $I^-$ ,  $RO^-$ ,  $ClO_4^-$ ,  $SO_4^{2-}$ ,  $HSO_4^-$ ,  $SO_3^{2-}$  and  $RSO_3^-$ , and more preferably selected from  $O^{2-}$ ,  $RBO_2^{2-}$ ,  $RCOO^-$ ,  $OH^-$ ,  $NO_3^-$ ,  $NO_2^-$ ,  $NO$ ,  $CO$ ,  $CN^-$ ,  $S^{2-}$ ,  $RS^-$ ,  $PO_3^{4-}$ ,  $H_2O$ ,  $CO_3^{2-}$ ,  $HCO_3^-$ ,  $ROH$ ,  $NRR'R''$ ,  $Cl^-$ ,  $Br^-$ ,  $OCN^-$ ,  $SCN^-$ ,  $RCN$ ,  $N_3^-$ ,  $F^-$ ,  $I^-$ ,  $RO^-$ ,  $ClO_4^-$ ,  $SO_4^{2-}$ ,  $HSO_4^-$ ,  $SO_3^{2-}$  and  $RSO_3^-$  (preferably  $CF_3SO_3^-$ );
- 15 Y represents any non-coordinated counter ion, preferably selected from  $ClO_4^-$ ,  $BR_4^-$ ,  $[FeCl_4]^-$ ,  $PF_6^-$ ,  $RCOO^-$ ,  $NO_3^-$ ,  $NO_2^-$ ,  $RO^-$ ,  $N^+RR'R''R'''$ ,  $Cl^-$ ,  $Br^-$ ,  $F^-$ ,  $I^-$ ,  $RSO_3^-$ ,  $S_2O_6^{2-}$ ,  $OCN^-$ ,  $SCN^-$ ,  $Li^+$ ,  $Ba^{2+}$ ,  $Na^+$ ,  $Mg^{2+}$ ,  $K^+$ ,  $Ca^{2+}$ ,  $Cs^+$ ,  $PR_4^+$ ,  $RBO_2^{2-}$ ,  $SO_4^{2-}$ ,  $HSO_4^-$ ,  $SO_3^{2-}$ ,  $SbCl_6^-$ ,  $CuCl_4^{2-}$ ,  $CN$ ,  $PO_4^{3-}$ ,  $HPO_4^{2-}$ ,  $H_2PO_4^-$ , STP-derived anions,  $CO_3^{2-}$ ,  $HCO_3^-$  and  $BF_4^-$ , and more preferably selected from  $ClO_4^-$ ,  $BR_4^-$ ,  $[FeCl_4]^-$ ,  $PF_6^-$ ,  $RCOO^-$ ,  $NO_3^-$ ,  $NO_2^-$ ,  $RO^-$ ,  $N^+RR'R''R'''$ ,  $Cl^-$ ,  $Br^-$ ,  $F^-$ ,  $I^-$ ,  $RSO_3^-$  (preferably  $CF_3SO_3^-$ ),  $S_2O_6^{2-}$ ,  $OCN^-$ ,  $SCN^-$ ,  $Li^+$ ,  $Ba^{2+}$ ,  $Na^+$ ,  $Mg^{2+}$ ,  $K^+$ ,  $Ca^{2+}$ ,  $PR_4^+$ ,  $SO_4^{2-}$ ,  $HSO_4^-$ ,  $SO_3^{2-}$ , and  $BF_4^-$ ;
- 20 R, R', R'', R''' independently represent a group selected from hydrogen, hydroxyl, -OR (wherein R= alkyl, alkenyl, cycloalkyl, heterocycloalkyl, aryl, heteroaryl or carbonyl derivative group), -OAr, alkyl, alkenyl, cycloalkyl, heterocycloalkyl, aryl, heteroaryl and carbonyl derivative groups, each of R, Ar, alkyl, alkenyl, cycloalkyl, heterocycloalkyl, aryl, heteroaryl and carbonyl derivative groups being optionally substituted by one or more functional groups E, or R6 together with R7 and independently R8 together with R9 represent oxygen, wherein E is selected from functional groups containing oxygen, sulphur, phosphorus, nitrogen, selenium, halogens, and any electron donating and/or withdrawing groups, and preferably R, R', R'', R''' represent hydrogen, optionally substituted alkyl or optionally substituted aryl, more preferably hydrogen or optionally substituted phenyl, naphthyl or  $C_{1-4}$ -alkyl;
- 25 a represents an integer from 1 to 10, preferably from 1 to 4;
- k represents an integer from 1 to 10;
- 30 n represents zero or an integer from 1 to 10, preferably from 1 to 4;
- 35 m represents zero or an integer from 1 to 20, preferably from 1 to 8.

**[0026]** Preferably, the ligand L is of the general formula (BI):



wherein

- 45 g represents zero or an integer from 1 to 6;
- r represents an integer from 1 to 6;
- s represents zero or an integer from 1 to 6;
- Z1 and Z2 independently represent a heteroatom or a heterocyclic or heteroaromatic ring, Z1 and/or Z2 being optionally substituted by one or more functional groups E as defined below;
- 50 Q 1 and Q2 independently represent a group of the formula:



wherein

$10 > d + e + f > 1$ ;  $d = 0-9$ ;  $e = 0-9$ ;  $f = 0-9$ ;

each Y1 is independently selected from -O-, -S-, -SO-, -SO<sub>2</sub>-, -(G<sup>1</sup>)N-, -(G<sup>1</sup>)(G<sup>2</sup>)N- (wherein G<sup>1</sup> and G<sup>2</sup> are as defined below), -C(O)-, arylene, alkylene, heteroarylene, -P- and -P(O)-;

if  $s > 1$ , each -[Z1(R1)-(Q1)<sub>r</sub>]- group is independently defined;

R1, R2, R6, R7, R8, R9 independently represent a group selected from hydrogen, hydroxyl, -OR (wherein R = alkyl, alkenyl, cycloalkyl, heterocycloalkyl, aryl, heteroaryl or carbonyl derivative group), -OAr, alkyl, alkenyl, cycloalkyl, heterocycloalkyl, aryl, heteroaryl and carbonyl derivative groups, each of R, Ar, alkyl, alkenyl, cycloalkyl, heterocycloalkyl, aryl, heteroaryl and carbonyl derivative groups being optionally substituted by one or more functional groups E, or R6 together with R7 and independently R8 together with R9 represent oxygen;


E is selected from functional groups containing oxygen, sulphur, phosphorus, nitrogen, selenium, halogens, and any electron donating and/or withdrawing groups (preferably E is selected from hydroxy, mono- or polycarboxylate derivatives, aryl, heteroaryl, sulphonate, thiol (-RSH), thioethers (-R-S-R'), disulphides (-RSSR'), dithiolenes, mono- or polyphosphonates, mono- or polyphosphates, electron donating groups and electron withdrawing groups, and groups of formulae (G<sup>1</sup>)(G<sup>2</sup>)N-, (G<sup>1</sup>)(G<sup>2</sup>)(G<sup>3</sup>)N-, (G<sup>1</sup>)(G<sup>2</sup>)N-C(O)-, G<sup>3</sup>O- and G<sup>3</sup>C(O)-, wherein each of G<sup>1</sup>, G<sup>2</sup> and G<sup>3</sup> is independently selected from hydrogen, alkyl, electron donating groups and electron withdrawing groups (in addition to any amongst the foregoing));

or one of R1-R9 is a bridging group bound to another moiety of the same general formula;

T1 and T2 independently represent groups R4 and R5, wherein R4 and R5 are as defined for R1-R9, and if  $g = 0$  and  $s > 0$ , R1 together with R4, and/or R2 together with R5, may optionally independently represent =CH-R10, wherein R10 is as defined for R1-R9, or

T1 and T2 may together (-T2-T1-) represent a covalent bond linkage when  $s > 1$  and  $g > 0$ ;

if Z1 and/or Z2 represent N and T1 and T2 together represent a single bond linkage and R1 and/or R2 are absent, Q1 and/or Q2 may independently represent a group of the formula:



optionally any two or more of R1, R2, R6, R7, R8, R9 independently are linked together by a covalent bond;

if Z1 and/or Z2 represents O, then R1 and/or R2 do not exist;

if Z1 and/or Z2 represents S, N, P, B or Si then R1 and/or R2 may be absent;

if Z1 and/or Z2 represents a heteroatom substituted by a functional group E then R1 and/or R2 and/or R4 and/or R5 may be absent.

**[0027]** The groups Z1 and Z2 preferably independently represent an optionally substituted heteroatom selected from N, P, O, S, B and Si or an optionally substituted heterocyclic ring or an optionally substituted heteroaromatic ring selected from pyridine, pyrimidines, pyrazine, pyrimidine, pyrazole, pyrrole, imidazole, benzimidazole, quinoline, isoquinoline, carbazole, indole, isoindole, furane, thiophene, oxazole and thiazole.


**[0028]** The groups R1-R9 are preferably independently selected from -H, hydroxy-C<sub>0</sub>-C<sub>20</sub>-alkyl, halo-C<sub>0</sub>-C<sub>20</sub>-alkyl, nitroso, formyl-C<sub>0</sub>-C<sub>20</sub>-alkyl, carboxyl-C<sub>0</sub>-C<sub>20</sub>-alkyl and esters and salts thereof, carbamoyl-C<sub>0</sub>-C<sub>20</sub>-alkyl, sulpho-C<sub>0</sub>-C<sub>20</sub>-alkyl and esters and salts thereof, sulphamoyl-C<sub>0</sub>-C<sub>20</sub>-alkyl, amino-C<sub>0</sub>-C<sub>20</sub>-alkyl, aryl-C<sub>0</sub>-C<sub>20</sub>-alkyl, heteroaryl-C<sub>0</sub>-C<sub>20</sub>-alkyl, C<sub>0</sub>-C<sub>20</sub>-alkyl, alkoxy-C<sub>0</sub>-C<sub>8</sub>-alkyl, carbonyl-C<sub>0</sub>-C<sub>6</sub>-alkoxy, and aryl-C<sub>0</sub>-C<sub>6</sub>-alkyl and C<sub>0</sub>-C<sub>20</sub>-alkylamide.

**[0029]** One of R1-R9 may be a bridging group which links the ligand moiety to a second ligand moiety of preferably the same general structure. In this case the bridging group may have the formula -C<sub>n</sub>(R11)(R12)-(D)<sub>p</sub>-C<sub>m</sub>(R11)(R12)- bound between the two moieties, wherein p is zero or one, D is selected from a heteroatom or a heteroatom-containing group, or is part of an aromatic or saturated homonuclear and heteronuclear ring, n' is an integer from 1 to 4, m' is an integer from 1 to 4, with the proviso that  $n' + m' \leq 4$ , R11 and R12 are each independently preferably selected from -H, NR13 and OR14, alkyl, aryl, optionally substituted, and R13 and R14 are each independently selected from -H, alkyl, aryl, both optionally substituted. Alternatively, or additionally, two or more of R1-R9 together represent a bridging group linking atoms, preferably hetero atoms, in the same moiety, with the bridging group preferably being alkylene or hydroxy-alkylene or a heteroaryl-containing bridge.

**[0030]** In a first variant according to formula (BI), the groups T1 and T2 together form a single bond linkage and  $s > 1$ , according to general formula (BII):

5

10



15

wherein Z3 independently represents a group as defined for Z1 or Z2; R3 independently represents a group as defined for R1-R9; Q3 independently represents a group as defined for Q1, Q2; h represents zero or an integer from 1 to 6; and s=s-1.

**[0031]** In a first embodiment of the first variant, in general formula (BII), s'=1, 2 or 3; r=g=h=1; d=2 or 3; e=f=0; R6=R7=H, preferably such that the ligand has a general formula selected from:

20

25


30

35

40


45

50



55

and more preferably selected from:



**[0032]** In these preferred examples, R1, R2, R3 and R4 are preferably independently selected from -H, alkyl, aryl, heteroaryl, and/or one of R1-R4 represents a bridging group bound to another moiety of the same general formula and/or two or more of R1-R4 together represent a bridging group linking N atoms in the same moiety, with the bridging group being alkylene or hydroxy-alkylene or a heteroaryl-containing bridge, preferably heteroarylene. More preferably, R1, R2, R3 and R4 are independently selected from -H, methyl, ethyl, isopropyl, nitrogen-containing heteroaryl, or a bridging group bound to another moiety of the same general formula or linking N atoms in the same moiety with the bridging group being alkylene or hydroxy-alkylene.

**[0033]** According to this first embodiment, in the complex  $[M_a L_k X_n] Y_m$  preferably:

M= Mn(II)-(IV), Cu(I)-(III), Fe(II)-(III), Co(II)-(III);

X=  $CH_3CN$ ,  $OH_2$ ,  $Cl^-$ ,  $Br^-$ ,  $OCN^-$ ,  $N_3^-$ ,  $SCN^-$ ,  $OH^-$ ,  $O^{2-}$ ,  $PO_4^{3-}$ ,  $C_6H_5BO_2^{2-}$ ,  $RCOO^-$ ;

Y=  $ClO_4^-$ ,  $BPh_4^-$ ,  $Br^-$ ,  $Cl^-$ ,  $[FeCl_4]^-$ ,  $PF_6^-$ ,  $NO_3^-$


a= 1, 2, 3, 4;

n= 0, 1, 2, 3, 4, 5, 6, 7, 8, 9;

m= 1, 2, 3, 4; and

k= 1, 2, 4.

**[0034]** In a second embodiment of the first variant, in general formula (BII),  $s'=2$ ;  $r=g=h=1$ ;  $d=f=0$ ;  $e=1$ ; and each Y1 is independently alkylene or heteroarylene. The ligand preferably has the general formula:



wherein

$A_1, A_2, A_3, A_4$  are independently selected from  $C_{1-9}$ -alkylene or heteroarylene groups; and

$N_1$  and  $N_2$  independently represent a hetero atom or a heteroarylene group.

**[0035]** In a preferred second embodiment,  $N_1$  represents an aliphatic nitrogen,  $N_2$  represents a heteroarylene group, R1, R2, R3, R4 each independently represent -H, alkyl, aryl or heteroaryl, and  $A_1, A_2, A_3, A_4$  each represent  $-CH_2-$ .

**[0036]** One of R1-R4 may represent a bridging group bound to another moiety of the same general formula and/or two or more of R1-R4 may together represent a bridging group linking N atoms in the same moiety, with the bridging group being alkylene or hydroxy-alkylene or a heteroaryl-containing bridge. Preferably, R1, R2, R3 and R4 are independently selected from -H, methyl, ethyl, isopropyl, nitrogen-containing heteroaryl, or a bridging group bound to another moiety of the same general formula or linking N atoms in the same moiety with the bridging group being alkylene or hydroxy-alkylene.

**[0037]** Particularly preferably, the ligand has the general formula:



5

10

wherein R<sub>1</sub>, R<sub>2</sub> each independently represent -H, alkyl, aryl or heteroaryl.

**[0038]** According to this second embodiment, in the complex [M<sub>a</sub>L<sub>k</sub>X<sub>n</sub>]Y<sub>m</sub> preferably:

15

M= Fe(II)-(III), Mn(II)-(IV), Cu(II), Co(II)-(III);

X= CH<sub>3</sub>CN, OH<sub>2</sub>, Cl<sup>-</sup>, Br<sup>-</sup>, OCN<sup>-</sup>, N<sub>3</sub><sup>-</sup>, SCN<sup>-</sup>, OH<sup>-</sup>, O<sup>2-</sup>, PO<sub>4</sub><sup>3-</sup>, C<sub>6</sub>H<sub>5</sub>BO<sub>2</sub><sup>2-</sup>, RCOO<sup>-</sup>;

Y= ClO<sub>4</sub><sup>-</sup>, BPh<sub>4</sub><sup>-</sup>, Br<sup>-</sup>, Cl<sup>-</sup>, [FeCl<sub>4</sub>]<sup>-</sup>, PF<sub>6</sub><sup>-</sup>, NO<sub>3</sub><sup>-</sup>;

a= 1, 2, 3, 4;

20


n= 0, 1, 2, 3, 4, 5, 6, 7, 8, 9;

m= 1, 2, 3, 4; and

k= 1, 2, 4.

**[0039]** In a third embodiment of the first variant, in general formula (BII), s'=2 and r=g=h=1, according to the general formula:

25



30

35

**[0040]** In this third embodiment, preferably each Z1-Z4 represents a heteroaromatic ring; e=f=0; d=1; and R7 is absent, with preferably R1=R2=R3=R4= 2,4,6-trimethyl-3-SO<sub>3</sub>Na-phenyl, 2,6-diCl-3(or 4)-SO<sub>3</sub>Na-phenyl.

**[0041]** Alternatively, each Z1-Z4 represents N; R1-R4 are absent; both Q1 and Q3 represent =CH—[—Y1—]<sub>e</sub>—CH=; and both Q2 and Q4 represent —CH<sub>2</sub>—[—Y1—]<sub>n</sub>—CH<sub>2</sub>—.


**[0042]** Thus, preferably the ligand has the general formula:

45

50

55





15 wherein A represents optionally substituted alkylene optionally interrupted by a heteroatom; and n is zero or an integer from 1 to 5.

20 **[0043]** Preferably, R1-R6 represent hydrogen, n=1 and A= -CH<sub>2</sub>-, -CHOH-, -CH<sub>2</sub>N(R)CH<sub>2</sub>- or -CH<sub>2</sub>CH<sub>2</sub>N(R)CH<sub>2</sub>CH<sub>2</sub>- wherein R represents hydrogen or alkyl, more preferably A= -CH<sub>2</sub>-, -CHOH- or -CH<sub>2</sub>CH<sub>2</sub>NHCH<sub>2</sub>CH<sub>2</sub>-.

25 **[0044]** According to this third embodiment, in the complex [M<sub>a</sub>L<sub>k</sub>X<sub>n</sub>]Y<sub>m</sub> preferably:

M= Mn(II)-(IV), Co(II)-(III), Fe(II)-(III);

X= CH<sub>3</sub>CN, OH<sub>2</sub>, Cl<sup>-</sup>, Br<sup>-</sup>, OCN<sup>-</sup>, N<sub>3</sub><sup>-</sup>, SCN<sup>-</sup>, OH<sup>-</sup>, O<sup>2-</sup>, PO<sub>4</sub><sup>3-</sup>, C<sub>6</sub>H<sub>5</sub>BO<sub>2</sub><sup>2-</sup>, RCOO<sup>-</sup>;

Y= ClO<sub>4</sub><sup>-</sup>, BPh<sub>4</sub><sup>-</sup>, Br<sup>-</sup>, Cl<sup>-</sup>, [FeCl<sub>4</sub>]<sup>-</sup>, PF<sub>6</sub><sup>-</sup>, NO<sub>3</sub><sup>-</sup>;


a= 1, 2, 3, 4;

n= 0, 1, 2, 3, 4, 5, 6, 7, 8, 9;


m= 1, 2, 3, 4; and

k= 1, 2, 4.


30 **[0045]** In a second variant according to formula (BI), T1 and T2 independently represent groups R4, R5 as defined for R1-R9, according to the general formula (BIII):



40 **[0046]** In a first embodiment of the second variant, in general formula (BIII), s=1; r=1; g=0; d=f=1; e=1-4; Y1= -CH<sub>2</sub>-; and R1 together with R4, and/or R2 together with R5, independently represent =CH-R10, wherein R10 is as defined for R1-R9. In one example, R2 together with R5 represents =CH-R10, with R1 and R4 being two separate groups. Alternatively, both R1 together with R4, and R2 together with R5 may independently represent =CH-R10. Thus, preferred ligands may for example have a



55 **[0047]** Preferably, the ligand is selected from:



5

10 wherein R1 and R2 are selected from optionally substituted phenols, heteroaryl-C<sub>0</sub>-C<sub>20</sub>-alkyls, R3 and R4 are selected from -H, alkyl, aryl, optionally substituted phenols, heteroaryl-C<sub>0</sub>-C<sub>20</sub>-alkyls, alkylaryl, aminoalkyl, alkoxy, more preferably R1 and R2 being selected from optionally substituted phenols, heteroaryl-C<sub>0</sub>-C<sub>2</sub>-alkyls, R3 and R4 are selected from -H, alkyl, aryl, optionally substituted phenols, nitrogen-heteroaryl-C<sub>0</sub>-C<sub>2</sub>-alkyls.

**[0048]** According to this first embodiment, in the complex [M<sub>a</sub>L<sub>k</sub>X<sub>n</sub>]Y<sub>m</sub> preferably:

15

M= Mn(II)-(IV), Co(II)-(III), Fe(II)-(III);

X= CH<sub>3</sub>CN, OH<sub>2</sub>, Cl<sup>-</sup>, Br<sup>-</sup>, OCN<sup>-</sup>, N<sub>3</sub><sup>-</sup>, SCN<sup>-</sup>, OH<sup>-</sup>, O<sup>2-</sup>, PO<sub>4</sub><sup>3-</sup>, C<sub>6</sub>H<sub>5</sub>BO<sub>2</sub><sup>2-</sup>, RCOO<sup>-</sup>;

Y= ClO<sub>4</sub><sup>-</sup>, BPh<sub>4</sub><sup>-</sup>, Br<sup>-</sup>, Cl<sup>-</sup>, [FeCl<sub>4</sub>]<sup>-</sup>, PF<sub>6</sub><sup>-</sup>, NO<sub>3</sub><sup>-</sup>;

a= 1, 2, 3, 4;

n= 0, 1, 2, 3, 4, 5, 6, 7, 8, 9;


20

m= 1, 2, 3, 4; and

k=1, 2, 4.

**[0049]** In a second embodiment of the second variant, in general formula (BIII), s=1; r=1; g=0; d=f=1; e=1-4; Y1= -C(R')(R''), wherein R' and R'' are independently as defined for R1-R9. Preferably, the ligand has the general formula:

25



30

35

**[0050]** The groups R1, R2, R3, R4, R5 in this formula are preferably -H or C<sub>0</sub>-C<sub>20</sub>-alkyl, n=0 or 1, R6 is -H, alkyl, -OH or -SH, and R7, R8, R9, R10 are preferably each independently selected from -H, C<sub>0</sub>-C<sub>20</sub>-alkyl, heteroaryl-C<sub>0</sub>-C<sub>20</sub>-alkyl, alkoxy-C<sub>0</sub>-C<sub>8</sub>-alkyl and amino-C<sub>0</sub>-C<sub>20</sub>-alkyl.

**[0051]** According to this second embodiment, in the complex [M<sub>a</sub>L<sub>k</sub>X<sub>n</sub>]Y<sub>m</sub> preferably:

40

M= Mn(II)-(IV), Fe(II)-(III), Cu(II), Co(II)-(III);

X= CH<sub>3</sub>CN, OH<sub>2</sub>, Cl<sup>-</sup>, Br<sup>-</sup>, OCN<sup>-</sup>, N<sub>3</sub><sup>-</sup>, SCN<sup>-</sup>, OH<sup>-</sup>, O<sup>2-</sup>, PO<sub>4</sub><sup>3-</sup>, C<sub>6</sub>H<sub>5</sub>BO<sub>2</sub><sup>2-</sup>, RCOO<sup>-</sup>;

Y= ClO<sub>4</sub><sup>-</sup>, BPh<sub>4</sub><sup>-</sup>, Br<sup>-</sup>, Cl<sup>-</sup>, [FeCl<sub>4</sub>]<sup>-</sup>, PF<sub>6</sub><sup>-</sup>, NO<sub>3</sub><sup>-</sup>;

a= 1, 2, 3, 4;

45

n= 0, 1, 2, 3, 4;


m= 0, 1, 2, 3, 4, 5, 6, 7, 8; and

k= 1, 2, 3, 4.

**[0052]** In a third embodiment of the second variant, in general formula (BIII), s=0; g=1; d=e=0; f=1-4. Preferably, the ligand has the general formula:


50

55



5

10 **[0053]** More preferably, the ligand has the general formula:



15

20

wherein R1, R2, R3 are as defined for R2, R4, R5.

**[0054]** According to this third embodiment, in the complex  $[M_a L_k X_n] Y_m$  preferably:

25

M = Mn(II)-(IV), Fe(II)-(III), Cu(II), Co(II)-(III);

X = CH<sub>3</sub>CN, OH<sub>2</sub>, Cl<sup>-</sup>, Br<sup>-</sup>, OCN<sup>-</sup>, N<sub>3</sub><sup>-</sup>, SCN<sup>-</sup>, OH<sup>-</sup>, O<sup>2-</sup>, PO<sub>4</sub><sup>3-</sup>, C<sub>6</sub>H<sub>5</sub>BO<sub>2</sub><sup>2-</sup>, RCOO<sup>-</sup>;

Y = ClO<sub>4</sub><sup>-</sup>, BPh<sub>4</sub><sup>-</sup>, Br<sup>-</sup>, Cl<sup>-</sup>, [FeCl<sub>4</sub>]<sup>-</sup>, PF<sub>6</sub><sup>-</sup>, NO<sub>3</sub><sup>-</sup>;

a = 1, 2, 3, 4;

n = 0, 1, 2, 3, 4;


30

m = 0, 1, 2, 3, 4, 5, 6, 7, 8; and

k = 1, 2, 3, 4.

**[0055]** In a fourth embodiment of the second variant, the organic substance forms a complex of the general formula (A):

35



in which

40

M represents iron in the II, III, IV or V oxidation state, manganese in the II, III, IV, VI or VII oxidation state, copper in the I, II or III oxidation state, cobalt in the II, III or IV oxidation state, or chromium in the II-VI oxidation state;

X represents a coordinating species;

n represents zero or an integer in the range from 0 to 3;

z represents the charge of the complex and is an integer which can be positive, zero or negative;

45

Y represents a counter ion, the type of which is dependent on the charge of the complex;

q = z/[charge Y]; and

L represents a pentadentate ligand of the general formula (B):

50



55



wherein

each R<sup>1</sup>, R<sup>2</sup> independently represents -R<sup>4</sup>-R<sup>5</sup>,

R<sup>3</sup> represents hydrogen, optionally substituted alkyl, aryl or arylalkyl, or -R<sup>4</sup>-R<sup>5</sup>,

5 each R<sup>4</sup> independently represents a single bond or optionally substituted alkylene, alkenylene, oxyalkylene, aminoalkylene, alkylene ether, carboxylic ester or carboxylic amide, and

each R<sup>5</sup> independently represents an optionally N-substituted aminoalkyl group or an optionally substituted heteroaryl group selected from pyridinyl, pyrazinyl, pyrazolyl, pyrrolyl, imidazolyl, benzimidazolyl, pyrimidinyl, triazolyl and thiazolyl.

10 **[0056]** The ligand L having the general formula (B), as defined above, is a pentadentate ligand. By 'pentadentate' herein is meant that five hetero atoms can coordinate to the metal M ion in the metal-complex.

**[0057]** In formula (B), one coordinating hetero atom is provided by the nitrogen atom in the methylamine backbone, and preferably one coordinating hetero atom is contained in each of the four R<sup>1</sup> and R<sup>2</sup> side groups. Preferably, all the coordinating hetero atoms are nitrogen atoms.

15 **[0058]** The ligand L of formula (B) preferably comprises at least two substituted or unsubstituted heteroaryl groups in the four side groups. The heteroaryl group is preferably a pyridin-2-yl group and, if substituted, preferably a methyl- or ethyl-substituted pyridin-2-yl group. More preferably, the heteroaryl group is an unsubstituted pyridin-2-yl group. Preferably, the heteroaryl group is linked to methylamine, and preferably to the N atom thereof, *via* a methylene group. Preferably, the ligand L of formula (B) contains at least one optionally substituted amino-alkyl side group, more preferably two amino-ethyl side groups, in particular 2-(N-alkyl)amino-ethyl or 2-(N,N-dialkyl)amino-ethyl.

20 **[0059]** Thus, in formula (B) preferably R<sup>1</sup> represents pyridin-2-yl or R<sup>2</sup> represents pyridin-2-yl-methyl. Preferably R<sup>2</sup> or R<sup>1</sup> represents 2-amino-ethyl, 2-(N-(m)ethyl)amino-ethyl or 2-(N,N-di(m)ethyl)amino-ethyl. If substituted, R<sup>5</sup> preferably represents 3-methyl pyridin-2-yl. R<sup>3</sup> preferably represents hydrogen, benzyl or methyl.

**[0060]** Examples of preferred ligands L of formula (B) in their simplest forms are:

25 (i) pyridin-2-yl containing ligands such as:

N,N-bis(pyridin-2-yl-methyl)-bis(pyridin-2-yl)methylamine;

N,N-bis(pyrazol-1-yl-methyl)-bis(pyridin-2-yl)methylamine;

N,N-bis(imidazol-2-yl-methyl)-bis(pyridin-2-yl)methylamine;

30 N,N-bis(1,2,4-triazol-1-yl-methyl)-bis(pyridin-2-yl)methylamine;

N,N-bis(pyridin-2-yl-methyl)-bis(pyrazol-1-yl)methylamine;

N,N-bis(pyridin-2-yl-methyl)-bis(imidazol-2-yl)methylamine;

N,N-bis(pyridin-2-yl-methyl)-bis(1,2,4-triazol-1-yl)methylamine;

N,N-bis(pyridin-2-yl-methyl)-1,1-bis(pyridin-2-yl)-1-aminoethane;

35 N,N-bis(pyridin-2-yl-methyl)-1,1-bis(pyridin-2-yl)-2-phenyl-1-aminoethane;

N,N-bis(pyrazol-1-yl-methyl)-1,1-bis(pyridin-2-yl)-1-aminoethane;

N,N-bis(pyrazol-1-yl-methyl)-1,1-bis(pyridin-2-yl)-2-phenyl-1-aminoethane;

N,N-bis(imidazol-2-yl-methyl)-1,1-bis(pyridin-2-yl)-1-aminoethane;

N,N-bis(imidazol-2-yl-methyl)-1,1-bis(pyridin-2-yl)-2-phenyl-1-aminoethane;

40 N,N-bis(1,2,4-triazol-1-yl-methyl)-1,1-bis(pyridin-2-yl)-1-aminoethane;

N,N-bis(1,2,4-triazol-1-yl-methyl)-1,1-bis(pyridin-2-yl)-2-phenyl-1-aminoethane;

N,N-bis(pyridin-2-yl-methyl)-1,1-bis(pyrazol-1-yl)-1-aminoethane;

N,N-bis(pyridin-2-yl-methyl)-1,1-bis(pyrazol-1-yl)-2-phenyl-1-aminoethane;

N,N-bis(pyridin-2-yl-methyl)-1,1-bis(imidazol-2-yl)-1-aminoethane;

45 N,N-bis(pyridin-2-yl-methyl)-1,1-bis(imidazol-2-yl)-2-phenyl-1-aminoethane;

N,N-bis(pyridin-2-yl-methyl)-1,1-bis(1,2,4-triazol-1-yl)-1-aminoethane;

N,N-bis(pyridin-2-yl-methyl)-1,1-bis(1,2,4-triazol-1-yl)-1-aminoethane;

N,N-bis(pyridin-2-yl-methyl)-1,1-bis(pyridin-2-yl)-1-aminoethane;

N,N-bis(pyridin-2-yl-methyl)-1,1-bis(pyridin-2-yl)-1-aminoethane;

50 N,N-bis(pyridin-2-yl-methyl)-1,1-bis(pyridin-2-yl)-2-phenyl-1-aminoethane;

N,N-bis(pyridin-2-yl-methyl)-1,1-bis(pyridin-2-yl)-2-(4-sulphonic acid-phenyl)-1-aminoethane;

N,N-bis(pyridin-2-yl-methyl)-1,1-bis(pyridin-2-yl)-2-(pyridin-2-yl)-1-aminoethane;

N,N-bis(pyridin-2-yl-methyl)-1,1-bis(pyridin-2-yl)-2-(pyridin-3-yl)-1-aminoethane;

N,N-bis(pyridin-2-yl-methyl)-1,1-bis(pyridin-2-yl)-2-(pyridin-4-yl)-1-aminoethane;

55 N,N-bis(pyridin-2-yl-methyl)-1,1-bis(pyridin-2-yl)-2-(1-alkyl-pyridinium-4-yl)-1-aminoethane;

N,N-bis(pyridin-2-yl-methyl)-1,1-bis(pyridin-2-yl)-2-(1-alkyl-pyridinium-3-yl)-1-aminoethane;

N,N-bis(pyridin-2-yl-methyl)-1,1-bis(pyridin-2-yl)-2-(1-alkyl-pyridinium-2-yl)-1-aminoethane;

(ii) 2-amino-ethyl containing ligands such as:


5 N,N-bis(2-(N-alkyl)amino-ethyl)-bis(pyridin-2-yl)methylamine;  
 N,N-bis(2-(N-alkyl)amino-ethyl)-bis(pyrazol-1-yl)methylamine;  
 N,N-bis(2-(N-alkyl)amino-ethyl)-bis(imidazol-2-yl)methylamine;  
 N,N-bis(2-(N-alkyl)amino-ethyl)-bis(1,2,4-triazol-1-yl)methylamine;  
 N,N-bis(2-(N,N-dialkyl)amino-ethyl)-bis(pyridin-2-yl)methylamine;  
 N,N-bis(2-(N,N-dialkyl)amino-ethyl)-bis(pyrazol-1-yl)methylamine;  
 N,N-bis(2-(N,N-dialkyl)amino-ethyl)-bis(imidazol-2-yl)methylamine;  
 10 N,N-bis(2-(N,N-dialkyl)amino-ethyl)-bis(1,2,4-triazol-1-yl)methylamine;  
 N,N-bis(pyridin-2-yl-methyl)-bis(2-amino-ethyl)methylamine;  
 N,N-bis(pyrazol-1-yl-methyl)-bis(2-amino-ethyl)methylamine;  
 N,N-bis(imidazol-2-yl-methyl)-bis(2-amino-ethyl)methylamine;  
 15 N,N-bis(1,2,4-triazol-1-yl-methyl)-bis(2-amino-ethyl)methylamine.

[0061] More preferred ligands are:

20 N,N-bis(pyridin-2-yl-methyl)-bis(pyridin-2-yl)methylamine, hereafter referred to as N4Py.  
 N,N-bis(pyridin-2-yl-methyl)-1,1-bis(pyridin-2-yl)-1-aminoethane, hereafter referred to as MeN4Py,  
 N,N-bis(pyridin-2-yl-methyl)-1,-bis(pyridin-2-yl)-2-phenyl-1-aminoethane, hereafter referred to as BzN4Py.

[0062] In an alternative fourth embodiment, the organic substance forms a complex of the general formula (A) including a ligand (B) as defined above, but with the proviso that R<sup>3</sup> does not represent hydrogen.

25 [0063] In a fifth embodiment of the second variant, the organic substance forms a complex of the general formula (A) as defined above, but wherein L represents a pentadentate or hexadentate ligand of general formula (C):



30 wherein

each R<sup>1</sup> independently represents -R<sup>3</sup>-V, in which R<sup>3</sup> represents optionally substituted alkylene, alkenylene, oxyalkylene, aminoalkylene or alkylene ether, and V represents an optionally substituted heteroaryl group selected from pyridinyl, pyrazinyl, pyrazolyl, pyrrolyl, imidazolyl, benzimidazolyl, pyrimidinyl, triazolyl and thiazolyl;

35 W represents an optionally substituted alkylene bridging group selected from -CH<sub>2</sub>CH<sub>2</sub>-, -CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>-, -CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>-, -CH<sub>2</sub>-C<sub>6</sub>H<sub>4</sub>-CH<sub>2</sub>-, -CH<sub>2</sub>-C<sub>6</sub>H<sub>10</sub>-CH<sub>2</sub>-, and -CH<sub>2</sub>-C<sub>10</sub>H<sub>6</sub>-CH<sub>2</sub>-; and

R<sup>2</sup> represents a group selected from R<sup>1</sup>, and alkyl, aryl and arylalkyl groups optionally substituted with a substituent selected from hydroxy, alkoxy, phenoxy, carboxylate, carboxamide, carboxylic ester, sulphonate, amine, alkylamine and N<sup>+</sup>(R<sup>4</sup>)<sub>3</sub>, wherein R<sup>4</sup> is selected from hydrogen, alkanyl, alkenyl, arylalkanyl, arylalkenyl, oxyalkanyl, oxyalkenyl, aminoalkanyl, aminoalkenyl, alkanyl ether and alkenyl ether.

40 [0064] The ligand L having the general formula (C), as defined above, is a pentadentate ligand or, if R<sup>1</sup>=R<sup>2</sup>, can be a hexadentate ligand. As mentioned above, by 'pentadentate' is meant that five hetero atoms can coordinate to the metal M ion in the metal-complex. Similarly, by 'hexadentate' is meant that six hetero atoms can in principle coordinate to the metal M ion. However, in this case it is believed that one of the arms will not be bound in the complex, so that the hexadentate ligand will be penta coordinating.

45 [0065] In the formula (C), two hetero atoms are linked by the bridging group W and one coordinating hetero atom is contained in each of the three R<sup>1</sup> groups. Preferably, the coordinating hetero atoms are nitrogen atoms.

[0066] The ligand L of formula (C) comprises at least one optionally substituted heteroaryl group in each of the three R<sup>1</sup> groups. Preferably, the heteroaryl group is a pyridin-2-yl group, in particular a methyl- or ethyl-substituted pyridin-2-yl group. The heteroaryl group is linked to an N atom in formula (C), preferably *via* an alkylene group, more preferably a methylene group. Most preferably, the heteroaryl group is a 3-methyl-pyridin-2-yl group linked to an N atom *via* methylene.

[0067] The group R<sup>2</sup> in formula (C) is a substituted or unsubstituted alkyl, aryl or arylalkyl group, or a group R<sup>1</sup>. However, preferably R<sup>2</sup> is different from each of the groups R<sup>1</sup> in the formula above. Preferably, R<sup>2</sup> is methyl, ethyl, benzyl, 2-hydroxyethyl or 2-methoxyethyl. More preferably, R<sup>2</sup> is methyl or ethyl.

55 [0068] The bridging group W may be a substituted or unsubstituted alkylene group selected from -CH<sub>2</sub>CH<sub>2</sub>-, -CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>-, -CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>-, -CH<sub>2</sub>-C<sub>6</sub>H<sub>4</sub>-CH<sub>2</sub>-, -CH<sub>2</sub>-C<sub>6</sub>H<sub>10</sub>-CH<sub>2</sub>-, and -CH<sub>2</sub>-C<sub>10</sub>H<sub>6</sub>-CH<sub>2</sub>- (wherein -C<sub>6</sub>H<sub>4</sub>-, -C<sub>6</sub>H<sub>10</sub>-, -C<sub>10</sub>H<sub>6</sub>- can be *ortho*-, *para*-, or *meta*-C<sub>6</sub>H<sub>4</sub>-, -C<sub>6</sub>H<sub>10</sub>-, -C<sub>10</sub>H<sub>6</sub>-). Preferably, the bridging group W is an ethylene or 1,4-butylene group, more preferably an ethylene group.

## EP 1 109 965 B1

**[0069]** Preferably, V represents substituted pyridin-2-yl, especially methyl-substituted or ethyl-substituted pyridin-2-yl, and most preferably V represents 3-methyl pyridin-2-yl.

**[0070]** Examples of preferred ligands of formula (C) in their simplest forms are:

5 N-methyl-N,N',N'-tris(3-methyl-pyridin-2-ylmethyl)ethylene-1,2-diamine;  
N-ethyl-N,N',N'-tris(3-methyl-pyridin-2-ylmethyl)ethylene-1,2-diamine;  
N-benzyl-N,N',N'-tris(3-methyl-pyridin-2-ylmethyl)ethylene-1,2-diamine;  
N-(2-hydroxyethyl)-N,N',N'-tris(3-methyl-pyridin-2-ylmethyl)ethylene-1,2-diamine;  
N-(2-methoxyethyl)-N,N',N'-tris(3-methyl-pyridin-2-ylmethyl)ethylene-1,2-diamine;

10 N-methyl-N,N',N'-tris(5-methyl-pyridin-2-ylmethyl)ethylene-1,2-diamine;  
N-ethyl-N,N',N'-tris(5-methyl-pyridin-2-ylmethyl)ethylene-1,2-diamine;  
N-benzyl-N,N',N'-tris(5-methyl-pyridin-2-ylmethyl)ethylene-1,2-diamine;  
N-(2-hydroxyethyl)-N,N',N'-tris(5-methyl-pyridin-2-ylmethyl)ethylene-1,2-diamine;  
N-(2-methoxyethyl)-N,N',N'-tris(5-methyl-pyridin-2-ylmethyl)ethylene-1,2-diamine;

15 N-methyl-N,N',N'-tris(3-ethyl-pyridin-2-ylmethyl)ethylene-1,2-diamine;  
N-ethyl-N,N',N'-tris(3-ethyl-pyridin-2-ylmethyl)ethylene-1,2-diamine;  
N-benzyl-N,N',N'-tris(3-ethyl-pyridin-2-ylmethyl)ethylene-1,2-diamine;  
N-(2-hydroxyethyl)-N,N',N'-tris(3-ethyl-pyridin-2-ylmethyl)ethylene-1,2-diamine;  
N-(2-methoxyethyl)-N,N',N'-tris(3-ethyl-pyridin-2-ylmethyl)ethylene-1,2-diamine;

20 N-methyl-N,N',N'-tris(5-ethyl-pyridin-2-ylmethyl)ethylene-1,2-diamine;  
N-ethyl-N,N',N'-tris(5-ethyl-pyridin-2-ylmethyl)ethylene-1,2-diamine;  
N-benzyl-N,N',N'-tris(5-ethyl-pyridin-2-ylmethyl)ethylene-1,2-diamine; and  
N-(2-methoxyethyl)-N,N',N'-tris(5-ethyl-pyridin-2-ylmethyl)ethylene-1,2-diamine.

**[0071]** More preferred ligands are:

30 N-methyl-N,N',N'-tris(3-methyl-pyridin-2-ylmethyl)ethylene-1,2-diamine;  
N-ethyl-N,N',N'-tris(3-methyl-pyridin-2-ylmethyl)ethylene-1,2-diamine;  
N-benzyl-N,N',N'-tris(3-methyl-pyridin-2-ylmethyl)ethylene-1,2-diamine;  
N-(2-hydroxyethyl)-N,N',N'-tris(3-methyl-pyridin-2-ylmethyl)ethylene-1,2-diamine;  
and  
35 N-(2-methoxyethyl)-N,N',N'-tris(3-methyl-pyridin-2-ylmethyl)ethylene-1,2-diamine.

**[0072]** The most preferred ligands are:

40 N-methyl-N,N',N'-tris(3-methyl-pyridin-2-ylmethyl)ethylene-1,2-diamine; and  
N-ethyl-N,N',N'-tris(3-methyl-pyridin-2-ylmethyl)ethylene-1,2-diamine.

**[0073]** Preferably, the metal M in formula (A) is Fe or Mn, more preferably Fe.

**[0074]** Preferred coordinating species X in formula (A) may be selected from  $R^6OH$ ,  $NR^6_3$ ,  $R^6CN$ ,  $R^6OO^-$ ,  $R^6S^-$ ,  $R^6O^-$ ,  $R^6COO^-$ ,  $OCN^-$ ,  $SCN^-$ ,  $N_3^-$ ,  $CN^-$ ,  $F^-$ ,  $Cl^-$ ,  $Br^-$ ,  $I^-$ ,  $O^{2-}$ ,  $NO_3^-$ ,  $NO_2^-$ ,  $SO_4^{2-}$ ,  $SO_3^{2-}$ ,  $PO_4^{3-}$  and aromatic N donors selected from pyridines, pyrazines, pyrazoles, pyrroles, imidazoles, benzimidazoles, pyrimidines, triazoles and thiazoles, with  $R^6$  being selected from hydrogen, optionally substituted alkyl and optionally substituted aryl. X may also be the species  $LMO^-$  or  $LMOO^-$ , wherein M is a transition metal and L is a ligand as defined above. The coordinating species X is preferably selected from  $CH_3CN$ ,  $H_2O$ ,  $F^-$ ,  $Cl^-$ ,  $Br^-$ ,  $OOH^-$ ,  $R^6COO^-$ ,  $R^6O^-$ ,  $LMO^-$ , and  $LMOO^-$  wherein  $R^6$  represents hydrogen or optionally substituted phenyl, naphthyl, or  $C_1$ - $C_4$  alkyl.


50 **[0075]** The counter ions Y in formula (A) balance the charge z on the complex formed by the ligand L, metal M and coordinating species X. Thus, if the charge z is positive, Y may be an anion such as  $R^7COO^-$ ,  $BPh_4^-$ ,  $ClO_4^-$ ,  $BF_4^-$ ,  $PF_6^-$ ,  $R^7SO_3^-$ ,  $R^7SO_4^-$ ,  $SO_4^{2-}$ ,  $NO_3^-$ ,  $F^-$ ,  $Cl^-$ ,  $Br^-$ , or  $I^-$ , with  $R^7$  being hydrogen, optionally substituted alkyl or optionally substituted aryl. If z is negative, Y may be a common cation such as an alkali metal, alkaline earth metal or (alkyl)ammonium cation.

55 **[0076]** Suitable counter ions Y include those which give rise to the formation of storage-stable solids. Preferred counter ions for the preferred metal complexes are selected from  $R^7COO^-$ ,  $ClO_4^-$ ,  $BF_4^-$ ,  $PF_6^-$ ,  $R^7SO_3^-$  (in particular  $CF_3SO_3^-$ ),  $R^7SO_4^-$ ,  $SO_4^{2-}$ ,  $NO_3^-$ ,  $F^-$ ,  $Cl^-$ ,  $Br^-$ , and  $I^-$ , wherein  $R^7$  represents hydrogen or optionally substituted phenyl, naphthyl or  $C_1$ - $C_4$  alkyl.

EP 1 109 965 B1

[0077] It will be appreciated that the complex (A) can be formed by any appropriate means, including *in situ* formation whereby precursors of the complex are transformed into the active complex of general formula (A) under conditions of storage or use. Preferably, the complex is formed as a well-defined complex or in a solvent mixture comprising a salt of the metal M and the ligand L or ligand L-generating species. Alternatively, the catalyst may be formed *in situ* from suitable precursors for the complex, for example in a solution or dispersion containing the precursor materials. In one such example, the active catalyst may be *formed in situ* in a mixture comprising a salt of the metal M and the ligand L, or a ligand L-generating species, in a suitable solvent. Thus, for example, if M is iron, an iron salt such as FeSO<sub>4</sub> can be mixed in solution with the ligand L, or a ligand L-generating species, to form the active complex. In another such example, the ligand L, or a ligand L-generating species, can be mixed with metal M ions present in the substrate or wash liquor to form the active catalyst *in situ*. Suitable ligand L-generating species include metal-free compounds or metal coordination complexes that comprise the ligand L and can be substituted by metal M ions to form the active complex according to the formula (A).

[0078] Therefore, in alternative fourth and fifth embodiments, the organic substance is a compound of the general formula (D):



in which

M' represents hydrogen or a metal selected from Ti, V, Co, Zn, Mg, Ca, Sr, Ba, Na, K, and Li;

X represents a coordinating species;

a represents an integer in the range from 1 to 5;

b represents an integer in the range from 1 to 4;

c represents zero or an integer in the range from 0 to 5;


z represents the charge of the compound and is an integer which can be positive, zero or negative;

Y represents a counter ion, the type of which is dependent on the charge of the compound;

q = z/[charge Y]; and

L represents a pentadentate ligand of general formula (B) or (C) as defined above.

[0079] In a fourth embodiment of the first variant, the organic substance comprises a macrocyclic ligand of formula (E):



wherein


Z<sup>1</sup> and Z<sup>2</sup> are independently selected from monocyclic or polycyclic aromatic ring structures optionally containing one or more heteroatoms, each aromatic ring structure being substituted by one or more substituents;

Y<sup>1</sup> and Y<sup>2</sup> are independently selected from C, N, O, Si, P and S atoms;

A<sup>1</sup> and A<sup>2</sup> are independently selected from hydrogen, alkyl, alkenyl and cycloalkyl (each of alkyl, alkenyl and cycloalkyl) being optionally substituted by one or more groups selected from hydroxy, aryl, heteroaryl, sulphonate, phosphate, electron donating groups and electron withdrawing groups, and groups of formulae (G<sup>1</sup>)(G<sup>2</sup>)N-, G<sup>3</sup>OC(O)-, G<sup>3</sup>O- and G<sup>3</sup>C(O)-, wherein each of G<sup>1</sup>, G<sup>2</sup> and G<sup>3</sup> is independently selected from hydrogen and alkyl, and electron donating and/or withdrawing groups (in addition to any amongst the foregoing);

i and j are selected from 0, 1 and 2 to complete the valency of the groups Y<sup>1</sup> and Y<sup>2</sup>;

each of Q<sup>1</sup>-Q<sup>4</sup> is independently selected from groups of formula




- 5
- 10 wherein  $10 > a + b + c > 2$  and  $d \geq 1$ ;
- each  $Y^3$  is independently selected from -O-, -S-, -SO-, -SO<sub>2</sub>-, -(G<sup>1</sup>)N- (wherein G<sup>1</sup> is hereinbefore defined), -C(O)-, arylene, heteroarylene, -P- and -P(O)-;
- each of A<sup>3</sup>-A<sup>6</sup> is independently selected from the groups hereinbefore defined for A<sup>1</sup> and A<sup>2</sup>; and
- 15 wherein any two or more of A<sup>1</sup>-A<sup>6</sup> together form a bridging group, provided that if A<sup>1</sup> and A<sup>2</sup> are linked without simultaneous linking also to any of A<sup>3</sup>-A<sup>6</sup>, then the bridging group linking A<sup>1</sup> and A<sup>2</sup> must contain at least one carbonyl group.

**[0080]** In the ligands of formula (E), unless specifically stated to the contrary, all alkyl, hydroxyalkyl alkoxy, and alkenyl groups preferably have from 1 to 6, more preferably from 1 to 4 carbon atoms.

**[0081]** Moreover, preferred electron donating groups include alkyl (e.g. methyl), alkoxy (e.g. methoxy), phenoxy, and unsubstituted, monosubstituted and disubstituted amine groups. Preferred electron withdrawing groups include nitro, carboxy, sulphonyl and halo groups.

**[0082]** The ligands of formula (E) may be used in the form of complexes with an appropriate metal or, in some cases, in non-complexed form. In the non-complexed form, they rely upon complexing with a metal supplied in the form of a separate ingredient in the composition, specifically provided for supplying that metal, or upon complexing with a metal found as a trace element in tap water. However, where the ligand alone or in complex form carries a (positive) charge, a counter anion is necessary. The ligand or complex may be formed as a neutral species but it is often advantageous, for reasons of stability or ease of synthesis, to have a charged species with appropriate anion.


**[0083]** Therefore, in an alternative fourth embodiment, the ligand of formula (E) is ion-paired with a counter ion, which ion-pairing is denoted by formula (F):



wherein

- 35 H is a hydrogen atom;
- Y is a counter anion, the type of which is dependent on the charge of the complex;
- x is an integer such that one or more nitrogen atoms in L is protonated;
- z represents the charge of the complex and is an integer which can be positive or zero;
- q = z/[charge of Y]; and
- 40 L is a ligand of formula (E) as defined above.

**[0084]** In a further alternative fourth embodiment, the organic substance forms a metal complex of formula (G) based on the ion pairing of formula (F) thus:



wherein L, Y, x, z and q are as defined for formula (F) above and M is a metal selected from manganese in oxidation states II-V, iron II-V, copper I-III, cobalt I-III, nickel I-III, chromium II-VI, tungsten IV-VI, palladium V, ruthenium II-IV, vanadium III-IV and molybdenum IV-VI.

**[0085]** Especially preferred are the complexes of formula (G) wherein M represents manganese, cobalt, iron or copper.


**[0086]** In a preferred fourth embodiment, the organic substance forms a complex of the formula (H):



5

10

15



20

wherein M represents an iron atom in oxidation state II or III, a manganese atom in oxidation state II, III, IV or V, a copper atom in oxidation state I, II or III or a cobalt atom in oxidation state II, III or IV, X is a group which is either a bridge or is not a bridge between iron atoms, Y is a counter ion, x and y being  $\geq 1$ ,  $0 \leq n \leq 3$ , and z being the charge of the metal complex, and  $p = z / \text{charge of Y}$ ;  $R_1$  and  $R_2$  being independently one or more ring substituents selected from hydrogen and electron donating and withdrawing groups,  $R_3$  to  $R_8$  being independently hydrogen, alkyl, hydroxy-alkyl, alkenyl or variants of any of these when substituted by one or more electron donating or withdrawing groups.

25

[0087] For the avoidance of doubt, " $\leq$ " means "less than or equal to" and " $\geq$ " means "greater than or equal to".

[0088] Preferably, in the complex of formula (H), M represents an iron atom in oxidation state II or III or a manganese atom in oxidation state II, III, IV, or V. Preferably the oxidation state of M is III.

30

[0089] When M is iron, preferably the complex of formula (H) is in the form of a salt of iron (in oxidised state) dihalo-2,11-diazo[3.3](2,6)pyridinophane, dihalo-4-methoxy-2,11-diazo[3.3](2,6)pyridinophane and mixtures thereof, especially in the form of the chloride salt.

[0090] When M is manganese, preferably the complex of formula (H) is in the form of a salt of manganese (in oxidised state) N, N'-dimethyl-2,11-diazo[3.3](2,6)pyridinophane, especially in the form of the monohexafluorophosphate salt.

35

[0091] Preferably, X is selected from  $H_2O$ ,  $OH^-$ ,  $O^{2-}$ ,  $SH^-$ ,  $S^{2-}$ ,  $SO_4^{2-}$ ,  $NR_9R_{10}^-$ ,  $RCOO^-$ ,  $NR_9R_{10}R_{11}$ ,  $Cl^-$ ,  $Br^-$ ,  $F^-$ ,  $N_3^-$  and combinations thereof, wherein  $R_9$ ,  $R_{10}$  and  $R_{11}$  are independently selected from -H,  $C_{1-4}$  alkyl and aryl optionally substituted by one or more electron withdrawing and/or donating groups. More preferably, X is a halogen, especially a fluoride ion.

[0092] In the formulae (F), (G) and (H), the anionic counter ion equivalent Y is preferably selected from  $Cl^-$ ,  $Br^-$ ,  $I^-$ ,  $NO_3^-$ ,  $ClO_4^-$ ,  $SCN^-$ ,  $PF_6^-$ ,  $RSO_3^-$ ,  $RSO_4^-$ ,  $CF_3SO_3^-$ ,  $BPh_4^-$ , and  $OAc^-$ . A cationic counter ion equivalent is preferably absent.

40

[0093] In formula (H),  $R_1$  and  $R_2$  are preferably both hydrogen.  $R_3$  and  $R_4$  are preferably  $C_{1-4}$  alkyl, especially methyl.  $R_5$ - $R_8$  are each preferably hydrogen.

[0094] According to the values of x and y, the aforementioned preferred iron or manganese catalysts of formula (H) may be in the form of a monomer, dimer or oligomer. Without being bound by any theory, it has been conjectured that in the raw material or detergent composition state, the catalyst exists mainly or solely in monomer form but could be converted to dimer, or even oligomeric form, in the wash solution.

45

[0095] In typical washing compositions the level of the organic substance is such that the in-use level is from  $1 \mu M$  to 50mM, with preferred in-use levels for domestic laundry operations falling in the range 10 to  $100 \mu M$ . Higher levels may be desired and applied in industrial textile bleaching processes.

50

[0096] Preferably, the aqueous medium has a pH in the range from pH 6 to 13, more preferably from pH 6 to 11, still more preferably from pH 8 to 11, and most preferably from pH 8 to 10, in particular from pH 9 to 10.

[0097] The method of the present invention has particular application in detergent bleaching, especially for laundry cleaning. Accordingly, in another preferred embodiment, the method uses the organic substance in a liquor that additionally contains a surface-active material, optionally together with detergency builder.

55

[0098] The bleach liquor may for example contain a surface-active material in an amount of from 10 to 50% by weight. The surface-active material may be naturally derived, such as soap, or a synthetic material selected from anionic, nonionic, amphoteric, zwitterionic, cationic actives and mixtures thereof. Many suitable actives are commercially avail-

able and are fully described in the literature, for example in "Surface Active Agents and Detergents", Volumes I and II, by Schwartz, Perry and Berch.

**[0099]** Typical synthetic anionic surface-actives are usually water-soluble alkali metal salts of organic sulphates and sulphonates having alkyl groups containing from about 8 to about 22 carbon atoms, the term "alkyl" being used to include the alkyl portion of higher aryl groups. Examples of suitable synthetic anionic detergent compounds are sodium and ammonium alkyl sulphates, especially those obtained by sulphating higher (C<sub>8</sub>-C<sub>18</sub>) alcohols produced, for example, from tallow or coconut oil; sodium and ammonium alkyl (C<sub>9</sub>-C<sub>20</sub>) benzene sulphonates, particularly sodium linear secondary alkyl (C<sub>10</sub>-C<sub>15</sub>) benzene sulphonates; sodium alkyl glyceryl ether sulphates, especially those ethers of the higher alcohols derived from tallow or coconut oil fatty acid monoglyceride sulphates and sulphonates; sodium and ammonium salts of sulphuric acid esters of higher (C<sub>9</sub>-C<sub>18</sub>) fatty alcohol alkylene oxide, particularly ethylene oxide, reaction products; the reaction products of fatty acids such as coconut fatty acids esterified with isethionic acid and neutralised with sodium hydroxide; sodium and ammonium salts of fatty acid amides of methyl taurine; alkane mono-sulphonates such as those derived by reacting alpha-olefins (C<sub>8</sub>-C<sub>20</sub>) with sodium bisulphite and those derived by reacting paraffins with SO<sub>2</sub> and Cl<sub>2</sub> and then hydrolysing with a base to produce a random sulphonate; sodium and ammonium (C<sub>7</sub>-C<sub>12</sub>) dialkyl sulphosuccinates; and olefin sulphonates, which term is used to describe material made by reacting olefins, particularly (C<sub>10</sub>-C<sub>20</sub>) alpha-olefins, with SO<sub>3</sub> and then neutralising and hydrolysing the reaction product. The preferred anionic detergent compounds are sodium (C<sub>10</sub>-C<sub>15</sub>) alkylbenzene sulphonates, and sodium (C<sub>16</sub>-C<sub>18</sub>) alkyl ether sulphates.

**[0100]** Examples of suitable nonionic surface-active compounds which may be used, preferably together with the anionic surface-active compounds, include, in particular, the reaction products of alkylene oxides, usually ethylene oxide, with alkyl (C<sub>6</sub>-C<sub>22</sub>) phenols, generally 5-25 EO, *i.e.* 5-25 units of ethylene oxides per molecule; and the condensation products of aliphatic (C<sub>8</sub>-C<sub>18</sub>) primary or secondary linear or branched alcohols with ethylene oxide, generally 2-30 EO. Other so-called nonionic surface-actives include alkyl polyglycosides, sugar esters, long-chain tertiary amine oxides, long-chain tertiary phosphine oxides and dialkyl sulphoxides.

**[0101]** Amphoteric or zwitterionic surface-active compounds can also be used in the compositions of the invention but this is not normally desired owing to their relatively high cost. If any amphoteric or zwitterionic detergent compounds are used, it is generally in small amounts in compositions based on the much more commonly used synthetic anionic and nonionic actives.

**[0102]** The detergent bleach liquor will preferably comprise from 1 to 15 % wt of anionic surfactant and from 10 to 40 % by weight of nonionic surfactant. In a further preferred embodiment, the detergent active system is free from C<sub>16</sub>-C<sub>12</sub> fatty acid soaps.

**[0103]** The bleach liquor may also contain a detergency builder, for example in an amount of from about 5 to 80 % by weight, preferably from about 10 to 60 % by weight.

**[0104]** Builder materials may be selected from 1) calcium sequestrant materials, 2) precipitating materials, 3) calcium ion-exchange materials and 4) mixtures thereof.

**[0105]** Examples of calcium sequestrant builder materials include alkali metal polyphosphates, such as sodium tripolyphosphate; nitrilotriacetic acid and its water-soluble salts; the alkali metal salts of carboxymethyloxy succinic acid, ethylene diamine tetraacetic acid, oxydisuccinic acid, mellitic acid, benzene polycarboxylic acids, citric acid; and polyacetal carboxylates as disclosed in US-A-4,144,226 and US-A-4,146,495.

**[0106]** Examples of precipitating builder materials include sodium orthophosphate and sodium carbonate.

**[0107]** Examples of calcium ion-exchange builder materials include the various types of water-insoluble crystalline or amorphous aluminosilicates, of which zeolites are the best known representatives, *e.g.* zeolite A, zeolite B (also known as zeolite P), zeolite C, zeolite X, zeolite Y and also the zeolite P-type as described in EP-A-0,384,070.

**[0108]** In particular, the bleach liquor may contain any one of the organic and inorganic builder materials, though, for environmental reasons, phosphate builders are preferably omitted or only used in very small amounts. Typical builders usable in the present invention are, for example, sodium carbonate, calcite/carbonate, the sodium salt of nitrilotriacetic acid, sodium citrate, carboxymethyloxy malonate, carboxymethyloxy succinate and water-insoluble crystalline or amorphous aluminosilicate builder materials, each of which can be used as the main builder, either alone or in admixture with minor amounts of other builders or polymers as co-builder.

**[0109]** It is preferred that the composition contains not more than 5% by weight of a carbonate builder, expressed as sodium carbonate, more preferably not more than 2.5 % by weight to substantially nil, if the composition pH lies in the lower alkaline region of up to 10.

**[0110]** Apart from the components already mentioned, the bleach liquor can contain any of the conventional additives in amounts of which such materials are normally employed in fabric washing detergent compositions. Examples of these additives include buffers such as carbonates, lather boosters, such as alkanolamides, particularly the monoethanol amides derived from palmkernel fatty acids and coconut fatty acids; lather depressants, such as alkyl phosphates and silicones; anti-redeposition agents, such as sodium carboxymethyl cellulose and alkyl or substituted alkyl cellulose ethers; stabilisers, such as phosphonic acid derivatives (*i.e.* Dequest® types); fabric softening agents; inorganic salts

and alkaline buffering agents, such as sodium sulphate and sodium silicate; and, usually in very small amounts, fluorescent agents; perfumes; enzymes, such as proteases, cellulases, lipases, amylases and oxidases; germicides and colourants.

**[0111]** Transition metal sequestrants such as EDTA, and phosphonic acid derivatives such as EDTMP (ethylene diamine tetra(methylene phosphonate)) may also be included, in addition to the organic substance specified, for example to improve the stability sensitive ingredients such as enzymes, fluorescent agents and perfumes, but provided the composition remains bleaching effective. However, the treatment composition containing the organic substance, is preferably substantially, and more preferably completely, devoid of transition metal sequestrants (other than the organic substance).

**[0112]** Whilst the present invention is based on the catalytic bleaching of a substrate by atmospheric oxygen or air, it will be appreciated that small amounts of hydrogen peroxide or peroxy-based or -generating systems may be included in the treatment composition, if desired. Preferably, however, the composition will be devoid of peroxygen bleach or peroxy-based or -generating bleach systems.

**[0113]** The invention will now be further illustrated by way of the following non-limiting examples:

## EXAMPLES

### Example 1

**[0114]** This example describes a synthesis of a catalyst according to formula (A):

#### **(i) Preparation of MeN4Py ligand:**

**[0115]** The precursor N4Py.HClO<sub>4</sub> was prepared as follows:

To pyridyl ketone oxim (3 g, 15.1 mmol) was added ethanol (15 ml), concentrated ammonia solution (15 mL) and NH<sub>4</sub>OAc (1.21 g, 15.8 mmol). The solution was warmed until reflux. To this solution was added 4.64 g Zn in small portions. After the addition of all Zn, the mixture was refluxed for 1 hour and allowed to cool to ambient temperature. The solution was filtered and water (15 ml) was added. Solid NaOH was added until pH > 10 and the solution was extracted with CH<sub>2</sub>Cl<sub>2</sub> (3 x 20 ml). The organic layers were dried over Na<sub>2</sub>SO<sub>4</sub> and evaporated until dryness. Bis(pyridin-2-yl)methylamine (2.39 g, 12.9 mmol) was obtained as a colourless oil in 86% yield, showing the following analytical characteristics:

<sup>1</sup>H NMR (360 MHz, CDCl<sub>3</sub>): δ 2.64 (s, 2H, NH<sub>2</sub>), 5.18 (s, 1H, CH), 6.93 (m, 2H, pyridine), 7.22 (m, 2H, pyridine), 7.41 (m, 2H, pyridine), 8.32 (m, 2H, pyridine); <sup>13</sup>C NMR (CDCl<sub>3</sub>): δ 62.19 (CH), 121.73 (CH), 122.01 (CH), 136.56 (CH), 149.03 (CH), 162.64 (Cq).

**[0116]** To picolylchloride hydrochloride (4.06 g, 24.8 mmol) was added, at 0°C, 4.9 ml of a 5N NaOH solution. This emulsion was added by means of a syringe to bis(pyridin-2-yl)methylamine (2.3 g, 12.4 mmol) at 0°C. Another 5 ml of a 5N NaOH solution was added to this mixture. After warming to ambient temperature, the mixture was stirred vigorously for 40 hrs. The mixture was put in an ice bath and HClO<sub>4</sub> was added until pH < 1, whereupon a brown solid precipitated. The brown precipitate was collected by filtration and recrystallized from water. While stirring, this mixture was allowed to cool to ambient temperature, whereupon a light-brown solid precipitated which was collected by filtration and washed with cold water and air-dried (1.47 g).

**[0117]** From 0.5 g of the perchlorate salt of N4Py prepared as described above, the free amine was obtained by precipitating the salt with 2N NaOH and subsequently by extraction with CH<sub>2</sub>Cl<sub>2</sub>. To the free amine was added under argon 20 ml of dry tetrahydrofuran freshly distilled from LiAlH<sub>4</sub>. The mixture was stirred and cooled to -70 °C by an alcohol / dry ice bath. Now 1 ml of 2.5 N butyllithium solution in hexane was added giving an immediate dark red colour. The mixture was allowed to warm to -20 °C and now 0.1 ml of methyl iodide was added. The temperature was kept to -10 °C for 1 hour. Subsequently 0.5 g of ammonium chloride was added and the mixture was evaporated in vacuo. To the residue water was added and the aqueous layer was extracted with dichloromethane. The dichloromethane layer was dried on sodium sulphate, filtered and evaporated giving 0.4 g residue. The residue was purified by crystallisation from ethyl acetate and hexane giving 0.2 g of creamish powder (50% yield) showing the following analytical characteristics:

<sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>): δ (ppm) 2.05 (s, 3H, CH<sub>3</sub>), 4.01 (s, 4H, CH<sub>2</sub>), 6.92 (m, 2H, pyridine), 7.08 (m, 2H, pyridine), 7.39 (m, 4H, pyridine), 7.60 (m, 2H, pyridine), 7.98 (d, 2H, pyridine), 8.41 (m, 2H, pyridine), 8.57 (m, 2H, pyridine). <sup>13</sup>C NMR (100.55 MHz, CDCl<sub>3</sub>): δ (ppm) 21.7 (CH<sub>3</sub>), 58.2 (CH<sub>2</sub>), 73.2 (Cq), 121.4 (CH), 121.7 (CH), 123.4 (CH), 123.6 (CH), 136.0 (CH), 148.2 (Cq), 148.6 (Cq), 160.1 (Cq), 163.8 (Cq).

**(ii) Synthesis of the complex [(MeN4Py)Fe(CH<sub>3</sub>CN)](ClO<sub>4</sub>)<sub>2</sub>, Fe(MeN4Py):**

**[0118]** To a solution of 0.27 g of MeN4Py in 12 ml of a mixture of 6 ml acetonitrile and 6 ml methanol was added 350 mg Fe(ClO<sub>4</sub>)<sub>2</sub>·6H<sub>2</sub>O immediately a dark red colour formed. To the mix was added now 0.5 g of sodium perchlorate and a orange red precipitate formed immediately. After 5 minutes stirring and ultrasonic treatment the precipitate was isolated by filtration and dried in vacuo at 50°C. In this way 350 mg of an orange red powder was obtained in 70% yield showing the following analytical characteristics:

<sup>1</sup>H NMR (400 MHz, CD<sub>3</sub>CN): δ (ppm) 2.15, (CH<sub>3</sub>CN), 2.28 (s, 3H, CH<sub>3</sub>), 4.2 (ab, 4H, CH<sub>2</sub>), 7.05 (d, 2H, pyridine), 7.38 (m, 4H, pyridine), 7.71 (2t, 4H pyridine), 7.98 (t, 2H, pyridine), 8.96 (d, 2H pyridine), 9.06 (m, 2H, pyridine).

UV/V is (acetonitrile) [λ<sub>max</sub>, nm (ε, M<sup>-1</sup> cm<sup>-1</sup>): 381 (8400), 458 nm (6400).

Anal. Calcd for C<sub>25</sub>H<sub>26</sub>Cl<sub>2</sub>FeN<sub>6</sub>O<sub>8</sub>: C, 46.11; H, 3.87; N, 12.41; Cl, 10.47; Fe, 8.25. Found: C, 45.49; H, 3.95; N, 12.5; Cl, 10.7; Fe, 8.12.

Mass-ESP (cone voltage 17V in CH<sub>3</sub>CN): m/z 218.6 [MeN4PyFe]<sup>2+</sup>; 239.1 [MeN4PyFeCH<sub>3</sub>CN]<sup>2+</sup>.

**Example 2**

**[0119]** This example describes a synthesis of a catalyst according to formula (A):

**(i) Synthesis of BzN4Py ligand:**

**[0120]** To 1 g of the N4Py ligand prepared as described above, 20 ml of dry tetrahydrofuran freshly distilled from LiAlH<sub>4</sub>, was added under argon. The mixture was stirred and cooled to -70 °C by an alcohol / dry ice bath. Now 2 ml of 2.5 N butyllithium solution in hexane was added giving an immediate dark red colour. The mix was allowed to warm to -20°C and now 0.4 ml of benzyl bromide was added. The mixture was allowed to warm up to 25 °C and stirring was continued over night. Subsequently 0.5 g of ammonium chloride was added and the mixture was evaporated in vacuo. To the residue water was added and the aqueous layer was extracted with dichloromethane. The dichloromethane layer was dried on sodium sulphate, filtered and evaporated giving 1 g brown oily residue. According to NMR spectroscopy, the product was not pure but contained no starting material (N4Py). The residue was used without further purification.

**(ii) Synthesis of the complex [(BzN4Py)Fe(CH<sub>3</sub>CN)](ClO<sub>4</sub>)<sub>2</sub>, Fe(BzN4Py):**

**[0121]** To a solution of 0.2 g of the residue obtained by the previous described procedure in 10 ml of a mixture of 5 ml acetonitrile and 5 ml methanol was added 100 mg Fe(ClO<sub>4</sub>)<sub>2</sub>·6H<sub>2</sub>O immediately a dark red colour formed. To the mix was added now 0.25 g of sodium perchlorate and ethylacetate was allowed to diffuse into the mixture overnight. Some red crystals were formed which were isolated by filtration and washed with methanol. In this way 70 mg of a red powder was obtained showing the following analytical characteristics:

<sup>1</sup>H NMR (400 MHz, CD<sub>3</sub>CN): δ (ppm) 2.12, (s, 3H, CH<sub>3</sub>CN), 3.65 + 4.1 (ab, 4H, CH<sub>2</sub>), 4.42 (s, 2H, CH<sub>2</sub>-benzyl), 6.84 (d, 2H, pyridine), 7.35 (m, 4H, pyridine), 7.45 (m, 3 H, benzene) 7.65 (m, 4H benzene + pyridine), 8.08(m, 4H, pyridine), 8.95 (m, 4H pyridine).

UV/Vis (acetonitrile) [λ<sub>max</sub>, nm (ε, M<sup>-1</sup> cm<sup>-1</sup>): 380 (7400), 458 nm (5500).

Mass-ESP (cone voltage 17V in CH<sub>3</sub>CN): m/z 256.4 [BzN4Py]<sup>2+</sup>; 612 [BzN4PyFeClO<sub>4</sub>]<sup>+</sup>

**Example 3:**

**[0122]** This example describes syntheses of catalysts according to formula (C):

**[0123]** All reactions were performed under a nitrogen atmosphere, unless indicated otherwise. All reagents and solvents were obtained from Aldrich or Across and used as received, unless stated otherwise. Petroleum ether 40-60 was distilled using a rotavapor before using it as eluent. Flash column chromatography was performed using Merck silica gel 60 or aluminium oxide 90 (activity II-III according to Brockmann). <sup>1</sup>H NMR (300 MHz) and <sup>13</sup>C NMR (75 MHz) were recorded in CDCl<sub>3</sub>, unless stated otherwise.

Multiplicities were addressed with the normal abbreviations using p for quintet.

**Synthesis of starting materials for ligand synthesis:**

**[0124]** **Synthesis of N-benzyl amino acetonitrile.** N-benzyl amine (5.35 g, 50 mmol) was dissolved in a water :

methanol mixture (50 mL, 1:4). Hydrochloric acid (aq., 30 %) was added until the pH reached 7.0. Added was NaCN (2.45 g, 50 mmol). After cooling to 0 °C, formaline (aq. 35 %, 4.00 g, 50 mmol) was added. The reaction was followed by TLC (aluminium oxide; EtOAc : Et<sub>3</sub>N = 9:1) until benzylamine could be detected. Subsequently the methanol was evaporated *in vacuo* and the remaining oil "dissolved" in water. The aqueous phase was extracted with methylene chloride (3 x 50 mL). The organic layers were collected and the solvent removed *in vacuo*. The residue was purified by Kugelrohr distillation (p = 20 mm Hg, T = 120 °C) giving *N*-benzyl amino acetonitrile (4.39 g, 30 mmol, 60 %) as a colourless oil.

<sup>1</sup>H NMR: δ 7.37 - 7.30 (m, 5H), 3.94 (s, 2H), 3.57 (s, 2H), 1.67 (br s, 1H);

<sup>13</sup>C NMR: δ 137.74, 128.58, 128.46, 128.37, 127.98, 127.62, 117.60, 52.24, 36.19.

**[0125] Synthesis of *N*-ethyl amino acetonitrile.** This synthesis was performed analogously to the synthesis reported for *N*-benzyl amino acetonitrile. However, detection was done by dipping the TLC plate in a solution of KMnO<sub>4</sub> and heating the plate until bright spots appeared. Starting from ethylamine (2.25 g, 50 mmol), pure *N*-ethyl amino acetonitrile (0.68 g, 8.1 mmol, 16 %) was obtained as a slightly yellow oil.

<sup>1</sup>H NMR: δ 3.60 (s, 2H), 2.78 (q, *J* = 7.1, 2H), 1.22 (br s, 1H), 1.14 (t, *J* = 7.2, 3H);

<sup>13</sup>C NMR: δ 117.78, 43.08, 37.01, 14.53.

**[0126] Synthesis of *N*-ethyl ethylene-1,2-diamine.** The synthesis was performed according to Hageman; J.Org. Chem.; 14; 1949; 616, 634, starting from *N*-ethyl amino acetonitrile.

**[0127] Synthesis of *N*-benzyl ethylene-1,2-diamine.** Sodium hydroxide (890 mg; 22.4 mmol) was dissolved in ethanol (96 %, 20 mL), the process taking the better part of 2 hours. Added was *N*-benzyl amino acetonitrile (4, 2.92 g, 20 mmol) and Raney Nickel (approx. 0.5 g). Hydrogen pressure was applied (p = 3.0 atm.) until hydrogen uptake ceased.

The mixture was filtered over Cellite, washing the residue with ethanol. The filter should not run dry since Raney Nickel is relatively pyrophoric. The Cellite containing the Raney Nickel was destroyed by putting the mixture in dilute acid, causing gas formation). The ethanol was evaporated in *in vacuo* and the residue dissolved in water.

**[0128]** Upon addition of base (aq. NaOH, 5N) the product oiled out and was extracted with chloroform (3 x 20 mL). After evaporation of the solvent *in vacuo* the <sup>1</sup>H NMR showed the presence of benzylamine. Separation was enforced by column chromatography (silica gel; MeOH : EtOAc : Et<sub>3</sub>N = 1:8:1 ) yielding the benzyl amine, followed by the solvent mixture MeOH : EtOAc : Et<sub>3</sub>N = 5:4:1. Detection was done by using aluminium oxide as a solid phase in TLC, yielding pure *N*-benzyl ethylene-1,2-diamine (2.04 g, 13.6 mmol, 69 %).

<sup>1</sup>H NMR: δ 7.33 - 7.24 (m, 5H), 3.80 (s, 2H), 2.82 (t, *J* = 5.7, 2H), 2.69 (t, *J* = 5.7, 2H), 1.46 (br s, 3H);

<sup>13</sup>C NMR: δ 140.37, 128.22, 127.93, 126.73, 53.73, 51.88, 41.66.

**[0129] Synthesis of 2-acetoxymethyl-5-methyl pyridine.** 2,5-Lutidine (31.0 g, 290 mmol), acetic acid (180 mL) and hydrogen peroxide (30 mL, 30 %) were heated at 70-80 °C for 3 hours. Hydrogen peroxide (24 mL, 30 %) was added and the subsequent mixture heated for 16 hours at 60-70 °C. Most of the mixture of (probably) hydrogen peroxide, water, acetic acid, and peracetic acid was removed *in vacuo* (rotavap, water bath 50 °C until p = 20 mbar). The resulting mixture containing the *N*-oxide was added dropwise to acetic anhydride heated under reflux. This reaction was highly exothermic, and was controlled by the dropping speed. After heating under reflux for an hour, methanol was added dropwise. This reaction was highly exothermic. The resulting mixture was heated under reflux for another 30 minutes. After evaporation of the methanol (rotavap, 50 °C until p = 20 mbar), the resulting mixture was purified by Kugelrohr distillation (p = 20 mm Hg, T = 150 °C). The clear oil that was obtained still contained acetic acid.

This was removed by extraction (CH<sub>2</sub>Cl<sub>2</sub>, NaHCO<sub>3</sub> (sat.)) yielding the pure acetate of 2-acetoxymethyl-5-methyl pyridine (34.35 g, 208 mmol, 72 %) as a slightly yellow oil.

<sup>1</sup>H NMR: δ 8.43 (s, 1H), 7.52 (dd, *J* = 7.8, *J* = 1.7, 1H), 7.26 (d, *J* = 7.2, 1H), 5.18 (s, 2H), 2.34 (s, 3H), 2.15 (s, 3H);

<sup>13</sup>C NMR: δ 170.09, 152.32, 149.39, 136.74, 131.98, 121.14, 66.31, 20.39, 17.66.

**[0130] Synthesis of 2-acetoxymethyl-5-ethyl pyridine.** This synthesis was performed analogously to the synthesis reported for 2-acetoxymethyl-5-methyl pyridine. Starting from 5-ethyl-2-methyl pyridine (35.10 g, 290 mmol), pure 2-acetoxymethyl-5-ethyl pyridine (46.19 g, 258 mmol, 89%) was obtained as a slightly yellow oil.

<sup>1</sup>H NMR: δ 8.47 (s, 1H), 7.55 (d, *J* = 7.8, 1H), 7.29 (d, *J* = 8.1, 1H), 2.67 (q, *J* = 7.8, 2H), 2.14 (s, 3H), 1.26 (t, *J* = 7.77, 3H);

<sup>13</sup>C NMR: δ 170.56, 152.80, 149.11, 138.47, 135.89, 121.67, 66.72, 25.65, 20.78, 15.13.

**[0131] Synthesis of 2-acetoxymethyl-3-methyl pyridine.** This synthesis was performed analogously to the synthesis reported for 2-acetoxymethyl-5-methyl pyridine. The only difference was the reversal of the Kugelrohr distillation and the extraction. According to <sup>1</sup>H NMR a mixture of the acetate and the corresponding alcohol was obtained. Starting from 2,3-picoline (31.0 g, 290 mmol), pure 2-acetoxymethyl-3-methyl pyridine (46.19 g, 258 mmol, 89%, calculated for pure acetate) was obtained as a slightly yellow oil.

<sup>1</sup>H NMR: δ 8.45 (d, *J* = 3.9, 1H), 7.50 (d, *J* = 8.4, 1H), 7.17 (dd, *J* = 7.8, *J* = 4.8, 1H), 5.24 (s, 2H), 2.37 (s, 3H), 2.14 (s, 3H).

**[0132] Synthesis of 2-hydroxymethyl-5-methyl pyridine.** 2-Acetoxymethyl-5-methyl pyridine (30 g, 182 mmol) was dissolved in hydrochloric acid (100 mL, 4 N). The mixture was heated under reflux, until TLC (silica gel; triethylamine:ethyl acetate:petroleum ether 40-60 = 1:9:19) showed complete absence of the acetate (normally 1 hour). The

mixture was cooled, brought to pH > 11, extracted with dichloromethane (3 x 50 mL) and the solvent removed *in vacuo*. Pure 2-hydroxymethyl-5-methyl pyridine (18.80 g, 152 mmol, 84 %) was obtained by Kugelrohr distillation (p = 20 mm Hg, T = 130 °C) as a slightly yellow oil.

<sup>1</sup>H NMR: δ 8.39 (s, 1H), 7.50 (dd, J = 7.8, J = 1.8, 1H), 7.15 (d, J = 8.1, 1H), 4.73 (s, 2H), 3.83 (br s, 1H), 2.34 (s, 3H);  
<sup>13</sup>C NMR: δ 156.67, 148.66, 137.32, 131.62, 120.24, 64.12, 17.98.

**[0133] Synthesis of 2-hydroxymethyl-5-ethyl pyridine.** This synthesis was performed analogously to the synthesis reported for 2-hydroxymethyl-5-methyl pyridine. Starting from 2-acetoxymethyl-5-ethyl pyridine (40 g, 223 mmol), pure 2-hydroxymethyl-5-ethyl pyridine (26.02 g, 189 mmol, 85 %) was obtained as a slightly yellow oil.

<sup>1</sup>H NMR: δ 8.40 (d, J = 1.2, 1H), 7.52 (dd, J = 8.0, J = 2.0, 1H), 7.18 (d, J = 8.1, 1H), 4.74 (s, 2H), 3.93 (br s, 1H), 2.66 (q, J = 7.6, 2H), 1.26 (t, J = 7.5, 3H);

<sup>13</sup>C NMR: δ 156.67, 148.00, 137.87, 136.13, 120.27, 64.07, 25.67, 15.28.

**[0134] Synthesis of 2-hydroxymethyl-3-methyl pyridine.** This synthesis was performed analogously to the synthesis reported for 2-hydroxymethyl-5-methyl pyridine. Starting from 2-acetoxymethyl-3-methyl pyridine (25g (recalculated for the mixture), 152 mmol), pure 2-hydroxymethyl-3-methyl pyridine (15.51 g, 126 mmol, 83 %) was obtained as a slightly yellow oil.

<sup>1</sup>H NMR: δ 8.40 (d, J = 4.5, 1H), 7.47 (d, J = 7.2, 1H), 7.15 (dd, J = 7.5, J = 5.1, 1H), 4.85 (br s, 1H), 4.69 (s, 1H), 2.22 (s, 3H);

<sup>13</sup>C NMR: δ 156.06, 144.97, 137.38, 129.53, 121.91, 61.38, 16.30.

#### (i) Synthesis of ligands:

**[0135] Synthesis of *N*-methyl-*N,N,N*-tris(pyridin-2-ylmethyl)ethylene-1,2-diamine (L1).** The ligand L1 (comparative) was prepared according to Bernal, Ivan; Jensen, Inge Margrethe; Jensen, Kenneth B.; McKenzie, Christine J.; Toftlund, Hans; Tuchagues, Jean-Pierre; J.Chem.Soc.Dalton Trans.; 22; 1995; 3667-3676.

**[0136] Synthesis of *N*-methyl-*N,N,N*-tris(3-methylpyridin-2-ylmethyl)ethylene-1,2-diamine (L2, MeTrilen).** 2-Hydroxymethyl-3-methyl pyridine (5.00 g, 40.7 mmol) was dissolved in dichloromethane (30 mL). Thionyl chloride (30 mL) was added dropwise under cooling (ice bath). The resulting mixture was stirred for 1 hour and the solvents removed *in vacuo* (rotavap, until p = 20 mm Hg, T = 50 °C). To the resultant mixture was added dichloromethane (25 mL). Subsequently NaOH (5 N, aq.) was added dropwise until the pH (aqua) ≥ 11. The reaction was quite vigorous in the beginning, since part of the thionyl chloride was still present. *N*-methyl ethylene-1,2-diamine (502 mg, 6.8 mmol) and additional NaOH (5 N, 10 mL) were added. The reaction mixture was stirred at room temperature for 45 hours. The mixture was poured into water (200 mL), and the pH checked (≥ 14, otherwise addition of NaOH (aq. 5N)). The reaction mixture was extracted with dichloromethane (3 or 4 x 50 mL, until no product could be detected by TLC). The combined organic phases were dried and the solvent removed *in vacuo*. Purification was enforced as described before, yielding *N*-methyl-*N,N,N*-tris(3-methylpyridin-2-ylmethyl)ethylene-1,2-diamine as a slightly yellow oil. Purification was enforced by column chromatography (aluminium oxide 90 (activity II-III according to Brockmann); triethylamine : ethyl acetate : petroleum ether 40-60 = 1:9:10) until the impurities were removed according to TLC (aluminium oxide, same eluent, R<sub>f</sub> ≈ 0.9). The compound was eluted using ethylacetate : triethyl amine = 9:1. *N*-methyl-*N,N,N*-tris(3-methylpyridin-2-ylmethyl)ethylene-1,2-diamine (L2, 1.743 g, 4.30 mmol, 63 %) was obtained.

<sup>1</sup>H NMR: δ 8.36 (d, J = 3.0, 3H), 7.40 - 7.37 (m, 3H), 7.11-7.06 (m, 3H), 3.76 (s, 4H), 3.48 (s, 2H), 2.76 - 2.71 (m, 2H), 2.53 - 2.48 (m, 2H), 2.30 (s, 3H), 2.12 (s, 6H), 2.05 (s, 3H);

<sup>13</sup>C NMR: δ 156.82, 156.77, 145.83, 145.67, 137.61, 133.14, 132.72, 122.10, 121.88, 62.32, 59.73, 55.19, 51.87, 42.37, 18.22, 17.80.

**[0137] Synthesis of *N*-ethyl-*N,N,N*-tris(3-methylpyridin-2-ylmethyl)ethylene-1,2-diamine (L3, EtTrilen).** This synthesis is performed analogously to the synthesis for L2. Starting from 2-hydroxymethyl-3-methyl pyridine (25.00 g, 203 mmol) and *N*-ethyl ethylene-1,2-diamine (2.99 g, 34.0 mmol), *N*-ethyl-*N,N,N*-tris(methylpyridin-2-ylmethyl)ethylene-1,2-diamine (L3, 11.49 g, 28.5 mmol, 84 %) was obtained. Column chromatography (aluminium oxide; Et<sub>3</sub>N : EtOAc : petroleum ether 40-60 = 1:9:30, followed by Et<sub>3</sub>N : EtOAc = 1:9).

<sup>1</sup>H NMR: δ 8.34 - 8.30 (m, 3H), 7.40 - 7.34 (m, 3H), 7.09 - 7.03 (m, 3H), 3.71 (s, 4H), 3.58 (s, 2H), 2.64 - 2.59 (m, 2H), 2.52 - 2.47 (m, 2H), 2.43 - 2.36 (m, 2H), 2.31 (s, 3H), 2.10 (s, 6H), 0.87 (t, J = 7.2, 3H);

<sup>13</sup>C NMR: δ 157.35, 156.92, 145.65, 137.61, 133.14, 132.97, 122.09, 121.85, 59.81, 59.28, 51.98, 50.75, 48.02, 18.27, 17.80, 11.36.

**[0138] Synthesis of *N*-benzyl-*N,N,N*-tris(3-methylpyridin-2-ylmethyl)ethylene-1,2-diamine (L4, BzTrilen).** This synthesis is performed analogously to the synthesis for L2. Starting from 2-hydroxymethyl-3-methylpyridine (3.00 g 24.4 mmol), and *N*-benzyl ethylene-1,2-diamine (610 mg, 4.07 mmol), *N*-benzyl-*N,N,N*-tris(3-methylpyridin-2-ylmethyl)ethylene-1,2-diamine (L4, 1.363 g, 2.93 mmol, 72 %) was obtained. Column chromatography (aluminium oxide; Et<sub>3</sub>N : EtOAc : petroleum ether 40-60 = 1:9:10).

<sup>1</sup>H NMR: δ 8.33 - 8.29 (m, 3H), 7.37 - 7.33 (m, 3H), 7.21 - 7.03 (m, 8H), 3.66 (s, 4H), 3.60 (s, 2H), 3.42 (s, 2H), 2.72

- 2.67 (m, 2H), 2.50 - 2.45 (m, 2H), 2.23 (s, 3H), 2.03 (s, 6H);

$^{13}\text{C}$  NMR:  $\delta$  157.17, 156.96, 145.83, 145.78, 139.29, 137.91, 137.80, 133.45, 133.30, 128.98, 127.85, 126.62, 122.28, 122.22, 59.99, 58.83, 51.92, 51.54, 18.40, 17.95.

**[0139] Synthesis of *N*-hydroxyethyl-*N,N,N*-tris(3-methylpyridin-2-ylmethyl)ethylene-1,2-diamine (L5).** This synthesis is performed analogously to the synthesis for L6. Starting from 2-hydroxymethyl-3-methyl pyridine (3.49 g, 28.4 mmol), and *N*-hydroxyethyl ethylene-1,2-diamine (656 mg 6.30 mmol), after 7 days *N*-hydroxyethyl-*N,N,N*-tris(3-methylpyridin-2-ylmethyl)ethylene-1,2-diamine (L5, 379 mg, 0.97 mmol, 14 %) was obtained.

$^1\text{H}$  NMR:  $\delta$  8.31 - 8.28 (m, 3H), 7.35 - 7.33 (m, 3H), 7.06 - 7.00 (m, 3H), 4.71 (br s, 1H), 3.73 (s, 4H), 3.61 (s, 2H), 3.44 (t,  $J$  = 5.1, 2H), 2.68 (s, 4H), 2.57 (t,  $J$  = 5.0, 2H), 2.19 (s, 3H), 2.10 (s, 6H);

$^{13}\text{C}$  NMR:  $\delta$  157.01, 156.88, 145.91, 145.80, 137.90, 137.83, 133.30, 131.89, 122.30, 121.97, 59.60, 59.39, 57.95, 56.67, 51.95, 51.22, 18.14, 17.95.

**[0140] Synthesis of *N*-methyl-*N,N,N*-tris(5-methylpyridin-2-ylmethyl)ethylene-1,2-diamine (L6).** 2-hydroxymethyl-5-methyl pyridine (2.70 g, 21.9 mmol) was dissolved in dichloromethane (25 mL). Thionyl chloride (25 mL) was added dropwise under cooling (ice bath). The resulting mixture was stirred for 1 hour and the solvents removed *in vacuo* (rotavap, until  $p = 20$  mm Hg,  $T \pm 35^\circ\text{C}$ ). The remaining oil was used directly in the synthesis of the ligands, since it was known from the literature that the free picolyl chlorides are somewhat unstable and are highly lachrymatory. To the resultant mixture was added dichloromethane (25 mL) and *N*-methyl ethylene-1,2-diamine (360 mg, 4.86 mmol). Subsequently NaOH (5 N, aq.) was added dropwise. The reaction was quite vigorous in the beginning, since part of the thionyl chloride was still present. The aqueous layer was brought to  $\text{pH} = 10$ , and additional NaOH (5 N, 4.38 mL) was added. The reaction mixture was stirred until a sample indicated complete conversion (7 days). The reaction mixture was extracted with dichloromethane (3 x 25 mL). The combined organic phases were dried and the solvent removed *in vacuo*. Purification was enforced by column chromatography (aluminium oxide 90 (activity II-III according to Brockmann); triethylamine : ethyl acetate : petroleum ether 40-60 = 1:9:10) until the impurities were removed according to TLC (aluminium oxide, same eluent,  $R_f \approx 0.9$ ). The compound was eluted using ethyl acetate : triethyl amine = 9:1, yielding *N*-methyl-*N,N,N*-tris(5-methylpyridin-2-ylmethyl)ethylene-1,2-diamine (L6, 685 mg, 1.76 mmol, 36 %) as a slightly yellow oil.

$^1\text{H}$  NMR:  $\delta$  8.31 (s, 3H) 7.43 - 7.35 (m, 5H), 7.21 (d,  $J$  = 7.8, 1H), 3.76 (s, 4H), 3.56 (s, 2H), 2.74 - 2.69 (m, 2H), 2.63 - 2.58 (m, 2H), 2.27 (s, 6H), 2.16 (s, 3H);

$^{13}\text{C}$  NMR:  $\delta$  156.83, 156.43, 149.23, 149.18, 136.85, 136.81, 131.02, 122.41, 122.30, 63.83, 60.38, 55.53, 52.00, 42.76, 18.03.

**[0141] Synthesis of *N*-methyl-*N,N,N*-tris(5-ethylpyridin-2-ylmethyl)ethylene-1,2-diamine (L7).** This synthesis is performed analogously to the synthesis for L6. Starting from 2-hydroxymethyl-5-ethyl pyridine (3.00 g, 21.9 mmol), and *N*-methyl ethylene-1,2-diamine (360 mg, 4.86 mmol), after 7 days *N*-methyl-*N,N,N*-tris(5-ethylpyridin-2-ylmethyl)ethylene-1,2-diamine (L7, 545 mg, 1.26 mmol, 26 %) was obtained.  $^1\text{H}$  NMR:  $\delta$  8.34 (s, 3H), 7.44 - 7.39 (m, 5H), 7.26 (d,  $J$  = 6.6, 1H), 3.80 (s, 4H), 3.59 (s, 2H), 2.77 - 2.72 (m, 2H), 2.66 - 2.57 (m, 8H), 2.18 (s, 3H), 1.23 (t,  $J$  = 7.5, 9H);  $^{13}\text{C}$  NMR:  $\delta$  157.14, 156.70, 148.60, 148.53, 137.25, 135.70, 122.59, 122.43, 63.91, 60.48, 55.65, 52.11, 42.82, 25.73, 15.36.

(ii) Synthesis of metal-ligand complexes:

**[0142] Synthesis of *N*-methyl-*N,N,N*-tris(3-methylpyridin-2-ylmethyl)ethylene-1,2-diamine iron(II)chloride.  $\text{PF}_6$  ([L2 Fe(II)Cl]PF<sub>6</sub>).**  $\text{FeCl}_2 \cdot 4\text{H}_2\text{O}$  (51.2 mg, 257  $\mu\text{mol}$ ) was dissolved in MeOH :  $\text{H}_2\text{O} = 1:1$  (2.5 mL). The solution was heated to  $50^\circ\text{C}$ . Added was *N*-methyl-*N,N,N*-tris(3-methylpyridin-2-ylmethyl)ethylene-1,2-diamine (L2, 100 mg, 257  $\mu\text{mol}$ ) in MeOH :  $\text{H}_2\text{O} = 1:1$  (2.0 mL). Subsequently  $\text{NaPF}_6$  (86.4 mg, 514  $\mu\text{mol}$ ) in  $\text{H}_2\text{O}$  (2.5 mL) was added dropwise. Cooling to room temperature, filtration and drying *in vacuo* ( $p = 0.05$  mm Hg,  $T =$  room temperature) yielded the complex [L2 Fe(II)Cl]PF<sub>6</sub> (149 mg, 239  $\mu\text{mol}$ , 93 %) as a yellow solid.

$^1\text{H}$  NMR ( $\text{CD}_3\text{CN}$ , paramagnetic):  $\delta$  167.17, 142.18, 117.01, 113.34, 104.79, 98.62, 70.77, 67.04, 66.63, 58.86, 57.56, 54.49, 51.68, 48.56, 45.90, 27.99, 27.36, 22.89, 20.57, 14.79, 12.14, 8.41, 8.16, 7.18, 6.32, 5.78, 5.07, 4.29, 3.82, 3.43, 2.91, 2.05, 1.75, 1.58, 0.94, 0.53, -0.28, -1.25, -4.82, -18.97, -23.46.

**[0143] Synthesis of *N*-ethyl-*N,N,N*-tris(3-methylpyridin-2-ylmethyl)ethylene-1,2-diamine iron(II)chloride.  $\text{PF}_6$  ([L3 Fe(II)Cl]PF<sub>6</sub>).** This synthesis was performed analogously to the synthesis for [L2 Fe(II)Cl]PF<sub>6</sub>. Starting from *N*-ethyl-*N,N,N*-tris(3-methylpyridin-2-ylmethyl)ethylene-1,2-diamine (L3, 104 mg, 257  $\mu\text{mol}$ ) gave the complex [L3 Fe(II)Cl]PF<sub>6</sub> (146 mg, 229  $\mu\text{mol}$ , 89%) as a yellow solid.

$^1\text{H}$  NMR ( $\text{CD}_3\text{CN}$ , paramagnetic):  $\delta$  165.61, 147.20, 119.23, 112.67, 92.92, 63.14, 57.44, 53.20, 50.43, 47.80, 28.59, 27.09, 22.48, 8.55, 7.40, 3.63, 2.95, 2.75, 2.56, 2.26, 1.75, 1.58, 0.92, 0.74, -0.28, -1.68, -2.68, -12.36, -28.75.


**[0144] Synthesis of *N*-benzyl-*N,N,N*-tris(3-methylpyridin-2-ylmethyl)ethylene-1,2-diamine iron(II)chloride.  $\text{PF}_6$  ([L4 Fe(II)Cl]PF<sub>6</sub>).** This synthesis was performed analogously to the synthesis for [L2 Fe(II)Cl]PF<sub>6</sub>. Starting from *N*-benzyl-*N,N,N*-tris(3-methylpyridin-2-ylmethyl)ethylene-1,2-diamine (L4, 119.5 mg, 257  $\mu\text{mol}$ ) gave the complex

(172 mg, 229  $\mu\text{mol}$ , 95 %) as a yellow solid.

$^1\text{H}$  NMR ( $\text{CD}_3\text{CN}$ , paramagnetic):  $\delta$  166.33, 145.09, 119.80, 109.45, 92.94, 57.59, 52.83, 47.31, 28.40, 27.89, 16.28, 11.05, 8.70, 8.45, 7.69, 6.99, 6.01, 4.12, 2.89, 2.71, 1.93, 1.56, -0.28, -1.68, -2.58, -11.40, -25.32.

#### 5 **Example 4**

[0145] This example describes a synthesis of a catalyst of formula (H) wherein:-



$\text{R}_2\text{-R}_8\text{=H}$ ;  $\text{R}_1\text{=4-MeO}$ ;  $x\text{=1}$ ;  $y\text{=1}$ ;  $z\text{=1}$ ;  $\text{X=Cl}$ ,  $n\text{=2}$ ;  $\text{Y=Cl}$ ,  $p\text{=1}$ .

#### 30 **(i) Synthesis of the ligand 2,11-diaza[3.3]-(4-methoxy)(2,6)pyridinophane ((4OMe)LN<sub>4</sub>H<sub>2</sub>):**

[0146] *4-chloro-2,6-pyridyl dimethyl ester (2)*. A mixture of 4-hydroxy-2,6-pyridine dicarboxylic acid (12.2 g, 60 mmoles) and  $\text{PCl}_5$  (41.8g, 200 mmoles) in 100 ml of  $\text{CCl}_4$  was refluxed until the evolution of HCl ceased. Absolute methanol (50ml) was slowly added. After cooling, all the volatile material was removed. The mixture was then poured into 200 ml of water and ice. The diester crystallised immediately and was collected by filtration (70%).  $^1\text{H}$  NMR (200MHz,  $\text{H}_2\text{O}$ )  $\delta$  7.60 (2H,s), 4.05 (6H, s).

[0147] *4-methoxy-2,6-pyridine dimethanol (4)*. Metallic sodium (1g, 44 mmoles) was dissolved into 200 ml of dry methanol. 4-chloro-2,6-pyridyl dimethyl ester (9.2 g, 40 mmoles) was then added and the mixture was refluxed for 3 hours to obtain pure 4-methoxy-2,6-pyridyl dimethyl ester. To this solution, at RT,  $\text{NaBH}_4$  (9.1 g, 240 mmoles) was added in small portions and the mixture was refluxed for 16 hours. Acetone (30 ml) was then added and the solution refluxed for an additional 1 hour. After all the volatile material was removed, the residue was heated with 60 ml of a saturated  $\text{NaHCO}_3/\text{Na}_2\text{CO}_3$  solution. After dilution with 80 ml of water, the product was continuously extracted with  $\text{CHCl}_3$  for 2-3 days. Evaporation of the  $\text{CHCl}_3$  yielded 83 % of 4-methoxy-2,6-pyridine dimethanol.  $^1\text{H}$  NMR (200MHz,  $\text{H}_2\text{O}$ )  $\delta$  6.83 (2H,s), 5.30 (2H,s), 4.43 (4H,s), 3.82 (3H, s).

*4-methoxy-2,6-dichloromethylpyridine (5)*. This synthesis is carried out according literature.

[0148] *N,N'-ditosyl-2,11-diaza[3.3]-(4-methoxy)(2,6)pyridinophane*. the procedure is similar to that described in the literature. The crude product obtained is practically pure (yield=95%).

$^1\text{H}$ -NMR ( $\text{CDCl}_3$ , 250 MHz): 7.72 (4H, d,  $J\text{=7Hz}$ ), 7.4 (1H, t,  $J\text{=6Hz}$ ), 7.35 (4H, d,  $J\text{=7Hz}$ ), 7.1 (1H, d,  $J\text{=6Hz}$ ), 6.57 (2H, s), 4.45 (4H, s), 4.35 (4H, s), 3.65 (3H, s), 2.4 (6H, s).

[0149] *2,11-diaza[3.3]-(4-methoxy)(2,6)pyridinophane*. The procedure is similar to the one described previously. The crude product obtained is purified by chromatography (alumina,  $\text{CH}_2\text{Cl}_2/\text{MeOH}$  95:5), yield = 65%.

$^1\text{H}$ -NMR ( $\text{CDCl}_3$ , 250 MHz): 7.15 (1H, t,  $J\text{=6Hz}$ ), 6.55 (1H, d,  $J\text{=6Hz}$ ), 6.05 (2H, s), 3.95 (4H, s), 3.87 (4H, s), 3.65 (3H, s).

Mass spectrum (EI):  $\text{M}^+ = 270$  (100%)




**(ii) Synthesis of the complex  $[\text{Fe}(\text{4OMeLN}_4\text{H}_2)\text{Cl}_2]\text{Cl}$ :**

**[0150]** 270 mg of 2,11-diaza[3.3]-(4-methoxy)(2,6)pyridinophane (1 mmole) were dissolved in 15 ml of dry THF. To this solution was added a solution of 270 mg of  $\text{FeCl}_3 \cdot 6\text{H}_2\text{O}$  (1 mmole) in 5 ml of MeOH. The resulting mixture is evaporated to dryness and the solid product is dissolved in 10 ml of AcN with a minimum of MeOH. Slow diffusion of THF give 300 mg of brown crystals, yield = 70%. Elemental analysis for  $\text{C}_{15}\text{H}_{18}\text{N}_4\text{Cl}_3\text{OFe} \cdot 0,5\text{MeOH}$  (found/theoretical): C=41.5/41.61 H=4.46/4.52 N=12.5/12.08

IR (KBr pellets,  $\text{cm}^{-1}$ ): 3545, 3414, 3235, 3075, 2883, 1615, 1477, 1437, 1340, 1157, 1049, 883, 628, 338.

**Example 5:**

**[0151]** This example describes a synthesis of a catalyst of formula (H) wherein:-



$R_1-R_8=H$ ;  $x=1$ ;  $y=1$ ;  $z=1$ ;  $X=C1$ ,  $n=2$ ;  $Y=C1$ ,  $p=1$


**Synthesis of the complex  $[\text{Fe}(\text{LN}_4\text{H}_2)\text{Cl}_2]\text{Cl}$ :**

**[0152]** 240 mg of  $\text{LN}_4\text{H}_2$  (1 mmole) were dissolved in 15 ml of dry THF. To this solution was added a solution of 270 mg of  $\text{FeCl}_3 \cdot 6\text{H}_2\text{O}$  (1 mmole) in 5 ml of MeOH. The resulting mixture is stirred and gives spontaneously 340 mg of yellow powder,

yield = 85%. IR (KBr pellets,  $\text{cm}^{-1}$ ): 3445, 3031, 2851, 1629, 1062, 1473, 1427, 1335, 1157, 1118, 1045, 936, 796, 340, 318

**Example 6:**

**[0153]** This Example describes a synthesis of a catalyst of formula (H) wherein:-



$R_1=R_2=R_{5-8}=H$ ;  $R_3=R_4=Me$ ;  $x=1$ ;  $y=1$ ;  $n=2$ ;  $z=1$ ;  $X=F^-$ ;  $m=2$ ;  $Y=PF_6^-$ ;  $p=1$

**difluoro[N,N'-dimethyl-2,11-diaza[3.3](2,6)pyridinophane]manganese(III) hexafluorophosphate.**

**(i) Synthesis of the ligand N,N'-dimethyl-2,11-diaza[3.3](2,6)pyridinophane:**

**[0154]** 2,6-dichloromethylpyridine. A mixture of 2,6-dimethanolpyridine (5g, 36 mmoles) and 75 ml of  $SOCl_2$  was refluxed for 4 hours. The mixture was concentrated (half volume). Toluene was added (50 ml). The solid formed after cooling was then filtered and dissolved in water and the solution neutralised with  $NaHCO_3$ . The solid obtained is filtered and dried (65%).  $^1H$  NMR (200MHz,  $CDCl_3$ )  $\delta$  7.8 (1H,t,  $J=7$ Hz), 7.45 (2H,d,  $J=7$  Hz), 4.7 (4H, s).

**[0155]** Sodium p-toluenesulphonamidure. To a mixture of  $Na^+$  in dry EtOH (0.7 g, 29 mmoles) was added p-toluenesulphonamide (5 g, 29 mmoles) and the solution was refluxed for 2 hours. After cooling, the solid obtained was filtered, washed with EtOH and dried (quantitative yield).

**[0156]** N,N'-ditosyl-2,11-diaza[3.3](2,6)pyridinophane. To a solution of sodium p-toluenesulphonamidure (1.93 g, 10 mmoles) in 200 ml of dry DMF at  $80^\circ C$  was slowly added 2,6-dichloromethylpyridine (1.76 g, 10 mmoles). After 1 hour a new portion of sodium p-toluenesulphonamidure was added (1.93 g) and the final mixture stirred at  $80^\circ C$  for an addition 4 hours. The solution was then evaporated to dryness. The solid obtained was washed with water and then with EtOH and finally crystallised in an  $CHCl_3/MeOH$  mixture. The solid obtained is filtered and dried. The yield of (15) was 55 %.  $^1H$  NMR (200MHz,  $CDCl_3$ )  $\delta$  7.78 (4H,d,  $J=6$ Hz), 7.45 (6H,m), 7.15 (4H,d,  $J=6$ Hz), 4.4 (8H, s), 2.4 (6H,s)

**[0157]** 2,11-diaza[3.3](2,6)pyridinophane. A mixture of N,N'-ditosyl-2,11-diaza[3.3](2,6)pyridinophane (1.53 g, 2.8 mmoles) and 14 ml of  $H_2SO_4$  90 % was heated at  $110^\circ C$  for 2 hours. The solution, cooled and diluted with 14 ml of water, was then carefully poured into a saturated NaOH solution. The solid formed is extracted with chloroform. The organic layer is evaporated to dryness to yield 85 % of 2,11-diaza[3.3](2,6)pyridinophane.  $^1H$  NMR (200MHz,  $CDCl_3$ )  $\delta$  7.1 (2H,t,  $J=7$ Hz), 6.5 (4H,d,  $J=7$  Hz), 3.9 (8H, s).

**[0158]** N,N'-dimethyl-2,11-diaza[3.3](2,6)pyridinophane. A mixture of 2,11-diaza[3.3](2,6)pyridinophane (0.57 g, 2.4 mmoles), 120 ml of formic acid and 32 ml of formaldehyde (32% in water) was refluxed for 24 hours. Concentrated HCl (10 ml) were added and the solution evaporated to dryness. The solid was dissolved in water and basified with NaOH 5M, and the resulting solution was extracted with  $CHCl_3$ . The solid obtained was purified by chromatography on alox ( $CH_2Cl_2+1\%$  MeOH) to yield 51 % of N,N'-dimethyl-2,11-diaza[3.3](2,6)pyridinophane.  $^1H$  NMR (200MHz,  $CDCl_3$ )  $\delta$  7.15 (2H,t,  $J=7$ Hz), 6.8 (4H,d,  $J=7$  Hz), 3.9 (8H, s), 2.73 (6H,s).

**(ii) Synthesis of the complex:**

**[0159]**  $MnF_3$  (41.8 mg, 373 mmoles) was dissolved in 5 ml of MeOH, and N,N'-dimethyl-2,11-diaza[3.3](2,6)pyridinophane (0.1 g, 373 mmoles) was added with 5 ml of THF. After 30 minutes of stirring at RT, 4 ml of THF saturated in  $NBu_4PF_6$  were added, and the solution left without stirring until the crystallisation was finished. The product was collected by filtration to yield 80% of complex. Elemental analysis (found, theoretical): %C (38.35, 37.94), %N (11.32, 11.1), %H (3.75, 3.95). IR (KBr pellet,  $cm^{-1}$ ): 3086, 2965, 2930, 2821, 1607, 1478, 1444, 1425, 1174, 1034, 1019, 844, 796, 603, 574, 555. UV-Vis ( $CH_3CN$ ,  $\lambda$  in nm,  $\epsilon$ ): 500, 110; 850, 30; ( $CH_3CN/H_2O$ :1/1,  $\lambda$  in nm,  $\epsilon$ ): 465, 168; 850, 30.

**Example 7:**

Bleaching of tomato-oil stained cloths without and with addition of  $[\text{Fe}(\text{MeN4Py})(\text{CH}_3\text{CN})](\text{ClO}_4)_2$ , immediately after the wash (t=0) and after 24 h storage (t=1 day).

**[0160]** In an aqueous solution containing 10 mM carbonate buffer (pH 10) without and with 0.6 g/l LAS (linear alkylbenzene sulphonate) or containing 10 mM borate buffer (pH 8) without and with 0.6 g/l LAS, tomato-soya oil stained cloths (6x6 cm) were added and stirred for 30 minutes at 30 °C. In a second series of experiments, the same tests were done in the presence of 10  $\mu\text{M}$   $[\text{Fe}(\text{MeN4Py})(\text{CH}_3\text{CN})](\text{ClO}_4)_2$ , referred to in the table below as Fe(MeN4Py).

**[0161]** After the wash, the cloths were dried in a tumble drier and the reflectance was measured with a Minolta 3700d spectrophotometer at 460 nm. The difference in reflectance before and after the wash is defined as  $\Delta\text{R}460$  value.

**[0162]** The cloths were measured immediately after the wash (t=0), and after 24 h storage in a dark room under ambient conditions (t=1d). The results obtained are listed in the table below:

	$\Delta\text{R}$ value (t=0) blank (no cat)	$\Delta\text{R}$ value (t=0) + Fe (MeN4Py)	$\Delta\text{R}$ value (t=1d) blank	$\Delta\text{R}$ value (t=1d) + Fe (MeN4Py)
pH 8 no LAS	11.5	23	11.5	44
pH 8 with LAS	12.5	19	12.5	36
pH 10 no LAS	10.5	30	11.5	43
pH 10 with LAS	12.5	30	14	39

**[0163]** A clear bleaching effect can thus be observed after the treated fabric cloths were dried and stored.

**Example 8:**

Bleaching of tomato-oil stained cloths without and with addition of various metal catalysts measured after 24 hours storage in the dark under ambient conditions.

**[0164]** In an aqueous solution containing 10 mM carbonate buffer (pH 10) without and with 0.6 g/l LAS (linear alkylbenzene sulphonate) or containing 10 mM borate buffer (pH 8) without and with 0.6 g/l LAS, tomato-soya oil stained cloths were added and kept in contact with the solution under agitation for 30 minutes at 30 °C. In comparative experiments, the same experiments were done by addition of 5  $\mu\text{M}$  of dinuclear or 10  $\mu\text{M}$  mononuclear complex, referred to in the table below.

**[0165]** After the wash, the cloths were rinsed with water and subsequently dried at 30 °C and the change in colour was measured after leaving the cloths for 24 h in the dark with a Linotype-Hell scanner (ex Linotype). The change in colour (including bleaching) is expressed as the  $\Delta\text{E}$  value. The measured colour difference ( $\Delta\text{E}$ ) between the washed cloth and the unwashed cloth is defined as follows:

$$\Delta\text{E} = [(\Delta\text{L})^2 + (\Delta\text{a})^2 + (\Delta\text{b})^2]^{1/2}$$

wherein  $\Delta\text{L}$  is a measure for the difference in darkness between the washed and unwashed test cloth;  $\Delta\text{a}$  and  $\Delta\text{b}$  are measures for the difference in redness and yellowness respectively between both cloths. With regard to this colour measurement technique, reference is made to Commission International de l'Eclairage (CIE); Recommendation on Uniform Colour Spaces, colour difference equations, psychometric colour terms, supplement no 2 to CIE Publication, no 15, Colorimetry, Bureau Central de la CIE, Paris 1978.

**[0166]** The following complexes were used:

- i)  $[\text{Mn}_2(1,4,7\text{-trimethyl-1,4,7\text{-triazacyclononane})_2(\mu\text{-O})_3](\text{PF}_6)_2$  (**1**) *Synthesised according to EP-B-458397;*
- ii)  $[\text{Mn}(\text{LN4Me2})]$  (=difluoro[N,N'dimethyl-2,11-diaza[3.3](2,6)pyridinophane] manganese(III)hexafluorophosphate) (**2**)  
*Synthesised as described previously;*
- iii)  $[\text{Fe}(\text{OMe})\text{LN4H2Cl}_2]$  (=Fe(2,11-diaza[3.3]-(4-methoxy)(2,6)pyridinophane)Cl<sub>2</sub>) (**3**)

Synthesised as described previously;

iv) Cl<sub>2</sub>-CoCo (**4**)

Synthesised according to EP-A-408131;

v) Me<sub>2</sub>CoCo (**5**)

Synthesised according to EP-A-408131;

vi) [Fe(tpen)](ClO<sub>4</sub>)<sub>2</sub> (**6**)

Synthesised according to WO-A-9748787;

vii) [Fe(N,N,N'-tris(pyridin-2-ylmethyl)-N-methyl-1,2-ethylenediamine)Cl](PF<sub>6</sub>)<sub>2</sub> (**7**)

Synthesised according to I. Bernal, et al., J. Chem. Soc., Dalton Trans, 22, 3667 (1995);

viii) [Fe<sub>2</sub>(N,N,N,N'-tetrakis(benzimidazol-2-ylmethyl)-propan-2-ol-1,3-diamine)(μ-OH)(NO<sub>3</sub>)<sub>2</sub>](NO<sub>3</sub>)<sub>2</sub> (**8**)

Synthesised according to Brennan, et al., Inorg. Chem., 30, 1937 (1991);

ix) [Mn<sub>2</sub>(tpen)(μ-O)<sub>2</sub>(μ-OAc)](ClO<sub>4</sub>)<sub>2</sub> (**9**)

Synthesised according to Toftlund, H.; Markiewicz, A.; Murray, K.S.; Acta Chem. Scand., **44**, 443 (1990);

x) [Mn(N,N,N'-tris(pyridin-2-ylmethyl)-N'-methyl-1,2-ethylenediamine)Cl](PF<sub>6</sub>) (**10**)

Synthesised as follows:

To a solution of manganese chloride tetrahydrate in tetrahydrofuran (0.190g, 1 mmol of MnCl<sub>2</sub>·4H<sub>2</sub>O in 10 mL of THF) ligand trispicen(NMe) (0.347, 1 mmol) was added to give a brown precipitate (reference ligand: I. Bernal, et al., J. Chem. Soc., Dalton Trans, 22, 3667 (1995)). The mixture was stirred for 10 minutes and ammonium hexafluorophosphate (0.163g, 1 mmol) dissolved in THF was added to give a cream coloured precipitate. The mixture was filtered, the filtrate was washed with THF and dried under vacuum to furnish the complex (FW=522.21g·mol<sup>-1</sup>) as a white solid (0.499g, 86%). ESMS (m/z): 437 ([LMnCl]<sup>+</sup>)

xi) [Mn<sub>2</sub>(N,N'-bis(pyridin-2-ylmethyl)-1,2-ethylenediamine)<sub>2</sub>(μ-O)<sub>2</sub>](ClO<sub>4</sub>)<sub>3</sub> (**11**)

Synthesised according to Glerup, J.; Goodson, P. A.; Hazell, A.; Hazell, R.; Hodgson, D. J.; McKenzie, C. J.; Michelsen, K.; Rychlewska, U.; Toftlund, H. Inorg. Chem. (1994), 33(18), 4105-11;

xii) [Mn(N,N'-bis(pyridin-2-ylmethyl)-N,N'-dimethyl-1,2-ethylenediamine)<sub>2</sub>Cl<sub>2</sub>] (**12**)

Synthesised as follows:

Triethylamine (0.405g, 4 mmol) was a solution of salt of the ligand bispicen(NMe) (0.416g, 1 mmol) in tetrahydrofuran anhydrous (10 mL) (refligand: C. Li, et al, J. Chem. Soc., Dalton Trans. (1991), 1909-14). The mixture was stirred at room temperature for 30 minutes. A few drops of methanol were added. The mixture was filtered. Manganese chloride (0.198g, 1 mmol) dissolved in THF (1 mL) was added to the mixture to give, after a stirring of 30 minutes, a white precipitate. The solution was filtered, the filtrate was washed twice with dry ether and dried under vacuum. This gave 0.093g of complex (23% yield).

xiii) [Mn<sub>2</sub>(N,N,N,N'-tetrakis(pyridin-2-ylmethyl)-propan-1,3-diamine)(μ-O)(μ-OAc)<sub>2</sub>](ClO<sub>4</sub>)<sub>2</sub> (**13**)

Synthesised as follows:

To a stirred solution of 6.56 g 2-chloro-methylpyridine (40 mmol) and 0.75 ml 1,3-propanediamine (9 mmol) in 40 ml water, is added slowly at 70°C over a period of 10 minutes, 8 ml 10M NaOH-solution. The colour of the reaction turned from yellow to deep red. The reaction was stirred for an additional 30 minutes at 70°C, after which the reaction was cooled to room temperature. The reaction mixture was extracted with dichloromethane (totally 200 ml), after which the red organic layer was dried over MgSO<sub>4</sub>, filtered and evaporated under reduced pressure, to yield 4.51 g of a red/brown oil. After scratching the bottom with a spatula the residue turned solid, trying to purify the crude product by washing it with water the product became messy, so immediately the purification was stopped and dried with ether. A sample was taken to analyse the product by NMR, while the rest was immediately reacted with Mn(OAc)<sub>3</sub> (see complexation).

<sup>1</sup>H-NMR (400MHz) (CDCl<sub>3</sub>); δ (ppm): 1.65 (q-5, propane-A, 2H), 2.40 (t, propane-B, 4H), 3.60 (s, N-CH<sub>2</sub>-pyr, 8H), 6.95 (t, pyr-H4, 4H), 7.30 (d, pyr-H3, 4H), 7.45 (t, pyr-H5, 4H), 8.35 (d, pyr-H6, 4H).

## EP 1 109 965 B1

To a stirred solution of 4.51 g TPTN (0.0103 mol) in 40 ml methanol is added at room temperature (22°C) 2.76 g Mn(OAc)<sub>3</sub> (0.0103 mol). The colour of the reaction turned from orange to dark brown, after the addition the mixture was stirred for 30 minutes at room temperature and filtered. To the filtrate was added at room temperature 1.44 g NaClO<sub>4</sub> (0.0103 mmol) and the reaction mixture was stirred for another hour, filtered and nitrogen dried, yielding 0.73 g bright brown crystals (8%).

<sup>1</sup>H-NMR (400MHz) (CD<sub>3</sub>CN); δ (ppm): -42.66 (s), -15.43 (s), -4.8 (s, br.), 0-10 (m, br.), 13.81 (s), 45.82 (s), 49.28 (s), 60 (s, br.), 79 (s, br.), 96 (s, br.)

IR/ (cm<sup>-1</sup>): 3426, 1608 (C=C), 1563 (C=N), 1487, 1430 (C-H), 1090 (ClO<sub>4</sub>), 1030, 767, 623.

UV/Vis (λ, nm(ε, l·mol<sup>-1</sup>·cm<sup>-1</sup>): 260 (2.4 x 10<sup>4</sup>), 290 (sh), 370 (sh), 490 (5.1 x 10<sup>2</sup>), 530 (sh; 3.4 x 10<sup>2</sup>), 567 (sh), 715 (1.4 x 10<sup>2</sup>).

Mass spectrum: (ESP+) m/z 782 [TPTN Mn(II)Mn(III) (μ-OH) (μ-OAc)<sub>2</sub> (ClO<sub>4</sub>)<sub>2</sub>]<sup>+</sup>

ESR (CH<sub>3</sub>CN): The complex is ESR silent supporting the presence of a Mn(III)Mn(III) species.

Elemental analysis: found (expected for Mn<sub>2</sub>C<sub>31</sub>H<sub>38</sub>N<sub>6</sub>O<sub>14</sub>Cl<sub>2</sub> (MW=899): C 41.14 (41.4), H 4.1 (4.2), N 9.23 (9.34), O 24.8 (24.9), Cl 7.72 (7.9), Mn 12.1 (12.2).

xiv) [Mn<sub>2</sub>(tpa)<sub>2</sub>(μ-O)<sub>2</sub>](PF<sub>6</sub>)<sub>3</sub> (**14**)

*Synthesised according to D.K. Towle, C.A. Botsford, D.J. Hodgson, ICA, 141, 167 (1988);*

xv) [Fe(N4Py)(CH<sub>3</sub>CN)](ClO<sub>4</sub>)<sub>2</sub> (**15**)

*Synthesised according to WO-A-9534628;*

xvi) [Fe(MeN4Py)(CH<sub>3</sub>CN)](ClO<sub>4</sub>)<sub>2</sub> (**16**)

*Synthesised according to EP-A-0909809.*

xvii) [Mn<sub>2</sub>(2,6-bis{(bis(2-pyridylmethyl)amino)methyl}-4-methylphenol))(μ-OAc)<sub>2</sub>](ClO<sub>4</sub>)<sub>2</sub> (**17**)

*Synthesised according to H. Diril et al., J.Am.Chem.Soc., 111, 5102 (1989);*

xviii) [Mn<sub>2</sub>(N,N,N',N'-tetrakis(benzimidazol-2-ylmethyl)-propan-2-olate-1,3-diamine)] (μ-OAc)<sub>2</sub>(ClO<sub>4</sub>)<sub>2</sub> (**18**)

*Synthesised according to P. Marthur et al., J.Am.Chem.Soc., 109, 5227 (1987).*

Results:

**[0167]**

Table:

bleach activity on Tomato Oil stains expressed in $\Delta E$ values obtained for various metal complexes measured after 24 h.																				
	<b>BL *</b>	<b>1</b>	<b>2</b>	<b>3</b>	<b>4</b>	<b>5</b>	<b>6</b>	<b>7</b>	<b>8</b>	<b>9</b>	<b>10</b>	<b>11</b>	<b>12</b>	<b>13</b>	<b>14</b>	<b>15</b>	<b>16</b>	<b>17</b>	<b>18</b>	<b>16 **</b>
pH 8 -LAS	1	1	2	2	4	9	2	9	4	2	7	2	2	2	6	20	19	2	1	21
pH 8 +LAS	2	6	7	7	2	6	19	19	2	18	10	18	5	7	6	20	20	3	2	23
pH10 -LAS	1	1	3	3	9	17	2	4	9	2	2	4	2	5	4	19	18	3	8	21
pH10 +LAS	3	16	14	14	4	11	6	11	4	4	4	16	17	7	7	19	20	11	5	21

\* BL: Reference: no catalyst added, only buffer with and without LAS

\*\* Compound **16** with 10 mM hydrogen peroxide

[0168] Performing the experiment under argon (compound 16 at pH 10 with LAS) showed that the bleaching effect upon storage is absent, thus showing that dioxygen is involved in the bleaching process.

5 **Claims**

- 10 1. A method of treating a textile by contacting the textile with an organic substance which forms a complex with a transition metal, the treatment providing the organic substance in a form that is substantially devoid of peroxygen bleach or a peroxy-based or -generating bleach system whereby the complex catalyses bleaching of the textile by atmospheric oxygen after the treatment, wherein the treated textile is dried and bleaching is catalysed on the dry textile.
- 15 2. A method according to claim 1, wherein the treatment comprises contacting the textile with the organic substance in dry form.
3. A method according to claim 1, wherein the treatment comprises contacting the textile with a liquor containing the organic substance and then drying.
- 20 4. A method according to claim 3, wherein the liquor is an aqueous liquor.
5. A method according to claim 4, wherein the liquor is a spray-on fabric treatment fluid.
6. A method according to claim 4, wherein the liquor is a wash liquor for laundry cleaning.
- 25 7. A method according to claim 3, wherein the liquor is a non-aqueous liquor.
8. A method according to claim 7, wherein the liquor is a dry cleaning fluid,
9. A method according to claim 7, wherein the liquor is a spray-on aerosol fluid.
- 30 10. A method according to claim 3, wherein the medium has a pH value in the range from pH 6 to 11.
11. A method according to claim 10, wherein the liquor has a pH value in the range from pH 8 to 10.
- 35 12. A method according to any of claims 3 to 11, wherein the liquor is substantially devoid of a transition metal sequestrant.
13. A method according to any of claims 3 to 12, wherein the liquor further comprises a surfactant.
- 40 14. A method according to any of claims 3 to 13, wherein the liquor further comprises a builder.
15. A method according to any of claims 1 to 14, wherein the organic substance comprises a preformed complex of a ligand and a transition metal.
- 45 16. A method according to any of claims 3 to 14, wherein the organic substance comprises a free ligand that complexes with a transition metal present in the liquor.
17. A method according to any of claims 1 to 14, wherein the organic substance comprises a free ligand that complexes with a transition metal present in the textile.
- 50 18. A method according to any of claims 1 to 14, wherein the organic substance comprises a composition of a free ligand or a transition metal-substitutable metal-ligand complex, and a source of transition metal.
- 55 19. A method according to any of claims 1 to 18, wherein the organic substance forms a complex of the general formula (A1):



in which:

M represents a metal selected from Mn(II)-(III)-(IV)-(V), Cu(I)-(II)-(III), Fe(I)-(II)-(III)-(IV), Co(I)-(II)-(III), Ni(I)-(II)-(III), Cr(II)-(III)-(IV)-(V)-(VI)-(VII), Ti(II)-(III)-(IV), V(II)-(III)-(IV)-(V), Mo(II)-(III)-(IV)-(V)-(VI), W(IV)-(V)-(VI), Pd(II), Ru(II)-(III)-(IV)-(V) and Ag(I)-(II);

L represents a ligand, or its protonated or deprotonated analogue;

X represents a coordinating species selected from any mono, bi or tri charged anions and any neutral molecules able to coordinate the metal in a mono, bi or tridentate manner; .

Y represents any non-coordinated counter ion;

a represents an integer from 1 to 10;

k represents an integer from 1 to 10;

n represents zero or an integer from 1 to 10; and

m represents zero or an integer from 1 to 20.

**20.** A method according to claim 19, wherein in formula (A1):

X represents a coordinating species selected from  $O^{2-}$ ,  $RBO_2^{2-}$ ,  $RCOO^-$ ,  $RCONR^-$ ,  $OH^-$ ,  $NO_3^-$ ,  $NO_2^-$ ,  $NO$ ,  $CO$ ,  $S^{2-}$ ,  $RS^-$ ,  $PO_3^{4-}$ , STP-derived anions,  $PO_3OR^{3-}$ ,  $H_2O$ ,  $CO_3^{2-}$ ,  $HCO_3^-$ ,  $ROH$ ,  $NRR'R''$ ,  $RCN$ ,  $Cl^-$ ,  $Br^-$ ,  $OCN^-$ ,  $SCN^-$ ,  $CN^-$ ,  $N_3^-$ ,  $F^-$ ,  $I^-$ ,  $RO^-$ ,  $ClO_4^-$ ,  $SO_4^{2-}$ ,  $HSO_4^-$ ,  $SO_3^{2-}$  and  $RSO_3^-$ ; and

Y represents a counter ion selected from  $ClO_4^-$ ,  $BR_4^-$ ,  $[FeCl_4]^-$ ,  $PF_6^-$ ,  $RCOO^-$ ,  $NO_3^-$ ,  $NO_2^-$ ,  $RO^-$ ,  $N^+RR'R''R'''$ ,  $Cl^-$ ,  $Br^-$ ,  $F^-$ ,  $I^-$ ,  $RSO_3^-$ ,  $S_2O_6^{2-}$ ,  $OCN^-$ ,  $SCN^-$ ,  $Li^+$ ,  $Ba^{2+}$ ,  $Na^+$ ,  $Mg^{2+}$ ,  $K^+$ ,  $Ca^{2+}$ ,  $Cs^+$ ,  $PR_4^+$ ,  $RBO_2^{2-}$ ,  $SO_4^{2-}$ ,  $HSO_4^-$ ,  $SO_3^{2-}$ ,  $SbCl_6^-$ ,  $CuCl_4^{2-}$ ,  $CN$ ,  $PO_4^{3-}$ ,  $HPO_4^{2-}$ ,  $H_2PO_4^-$ , STP-derived anions,  $CO_3^{2-}$ ,  $HCO_3^-$  and  $BF_4^-$ , wherein R, R', R'', R''' independently represent a group selected from hydrogen, hydroxyl, -OR (wherein R= alkyl, alkenyl, cycloalkyl, heterocycloalkyl, aryl, heteroaryl or carbonyl derivative group), -OAr, alkyl, alkenyl, cycloalkyl, heterocycloalkyl, aryl, heteroaryl and carbonyl derivative groups, each of R, Ar, alkyl, alkenyl, cycloalkyl, heterocycloalkyl, aryl, heteroaryl and carbonyl derivative groups being optionally substituted by one or more functional groups E, or R6 together with R7 and independently R8 together with R9 represent oxygen;

E is selected from functional groups containing oxygen, sulphur, phosphorus, nitrogen, selenium, halogens, and any electron donating and/or withdrawing groups.

**21.** A method according to claim 19 or claim 20, wherein in formula (A1) :

M represents a metal selected from Mn(II)-(III)-(IV)-(V), Cu(I)-(II), Fe(II)-(III)-(IV) and Co(I)-(II)-(III);

X represents a coordinating species selected from  $O^{2-}$ ,  $RBO_2^{2-}$ ,  $RCOO^-$ ,  $OH^-$ ,  $NO_3^-$ ,  $NO_2^-$ ,  $NO$ ,  $CO$ ,  $CN^-$ ,  $S^{2-}$ ,  $RS^-$ ,  $PO_3^{4-}$ ,  $H_2O$ ,  $CO_3^{2-}$ ,  $HCO_3^-$ ,  $ROH$ ,  $NRR'R''$ ,  $Cl^-$ ,  $Br^-$ ,  $OCN^-$ ,  $SCN^-$ ,  $RCN$ ,  $N_3^-$ ,  $F^-$ ,  $I^-$ ,  $RO^-$ ,  $ClO_4^-$ ,  $SO_4^{2-}$ ,  $HSO_4^-$ ,  $SO_3^{2-}$  and  $RSO_3^-$ ;

Y represents a counter ion selected from  $ClO_4^-$ ,  $BR_4^-$ ,  $[FeCl_4]^-$ ,  $PF_6^-$ ,  $RCOO^-$ ,  $NO_3^-$ ,  $NO_2^-$ ,  $RO^-$ ,  $N^+RR'R''R'''$ ,  $Cl^-$ ,  $Br^-$ ,  $F^-$ ,  $I^-$ ,  $RSO_3^-$ ,  $S_2O_6^{2-}$ ,  $OCN^-$ ,  $SCN^-$ ,  $Li^+$ ,  $Ba^{2+}$ ,  $Na^+$ ,  $Mg^{2+}$ ,  $K^+$ ,  $Ca^{2+}$ ,  $PR_4^+$ ,  $SO_4^{2-}$ ,  $HSO_4^-$ ,  $SO_3^{2-}$  and  $BF_4^-$ , wherein

R, R', R'', R''' represent represent hydrogen, optionally substituted alkyl or optionally substituted aryl;


a represents an integer from 1 to 4;

k represents an integer from 1 to 10;

n represents zero or an integer from 1 to 4; and

m represents zero or an integer from 1 to 8.

**22.** A method according to any of claims 19 to 21, wherein L represents a ligand of the general formula (BI):



wherein

g represents zero or an integer from 1 to 6;




r represents an integer from 1 to 6;

s represents zero or an integer from 1 to 6;

Z1 and Z2 independently represent a heteroatom or a heterocyclic or heteroaromatic ring, Z1 and/or Z2 being optionally substituted by one or more functional groups E as defined below;

Q1 and Q2 independently represent a group of the formula:



wherein

$10 > d + e + f > 1$ ;  $d = 0-9$ ;  $e = 0-9$ ;  $f = 0-9$ ;

each Y1 is independently selected from -O-, -S-, -SO-, -SO<sub>2</sub>-, -(G<sup>1</sup>)N-, -(G<sup>1</sup>)(G<sup>2</sup>)N- (wherein G<sup>1</sup> and G<sup>2</sup> are as defined below), -C(O)-, arylene, alkylene, heteroarylene, -P- and -P(O)-;

if  $s > 1$ , each  $-\text{Z1}(\text{R1})-(\text{Q1})_r-$  group is independently defined;

R1, R2, R6, R7, R8, R9 independently represent a group selected from hydrogen, hydroxyl, -OR (wherein R = alkyl, alkenyl, cycloalkyl, heterocycloalkyl, aryl, heteroaryl or carbonyl derivative group), -OAr, alkyl, alkenyl, cycloalkyl, heterocycloalkyl, aryl, heteroaryl and carbonyl derivative groups, each of R, Ar, alkyl, alkenyl, cycloalkyl, heterocycloalkyl, aryl, heteroaryl and carbonyl derivative groups being optionally substituted by one or more functional groups E, or R6 together with R7 and independently R8 together with R9 represent oxygen;


E is selected from functional groups containing oxygen, sulphur, phosphorus, nitrogen, selenium, halogens, and any electron donating and/or withdrawing groups (preferably E is selected from hydroxy, mono- or polycarboxylate derivatives, aryl, heteroaryl, sulphonate, thiol (-RSH), thioethers (-R-S-R'), disulphides (-RSSR'), dithio- lenes, mono- or polyphosphonates, mono- or polyphosphates, electron donating groups and electron withdrawing groups, and groups of formulae (G<sup>1</sup>) (G<sup>2</sup>)N-, (G<sup>1</sup>) (G<sup>2</sup>) (G<sup>3</sup>)N-, (G<sup>1</sup>) (G<sup>2</sup>)N-C(O)-, G<sup>3</sup>O- and G<sup>3</sup>C(O)-, wherein each of G<sup>1</sup>, G<sup>2</sup> and G<sup>3</sup> is independently selected from hydrogen, alkyl, electron donating groups and electron withdrawing groups (in addition to any amongst the foregoing));

or one of R1-R9 is a bridging group bound to another moiety of the same general formula;

T1 and T2 independently represent groups R4 and R5, wherein R<sub>4</sub> and R<sub>5</sub> are as defined for R1-R9, and if  $g = 0$  and  $s > 0$ , R1 together with R4, and/or R2 together with R5, may optionally independently represent =CH-R10, wherein R10 is as defined for R1-R9, or

T1 and T2 may together (-T2-T1-) represent a covalent bond linkage when  $s > 1$  and  $g > 0$ ;

if Z1 and/or Z2 represent N and T1 and T2 together represent a single bond linkage and R1 and/or R2 are absent, Q1 and/or Q2 may independently represent a group of the formula:



optionally any two or more of R1, R2, R6, R7, R8, R9 independently are linked together by a covalent bond;

if Z1 and/or Z2 represents O, then R1 and/or R2 do not exist;

if Z1 and/or Z2 represents S, N, P, B or Si then R1 and/or R2 may be absent;

if Z1 and/or Z2 represents a heteroatom substituted by a functional group E then R1 and/or R2 and/or R4 and/or R5 may be absent.


**23.** A method according to claim 22, wherein Z1 and Z2 independently represent an optionally substituted heteroatom selected from N, P, O, S, B and Si or an optionally substituted heterocyclic ring or an optionally substituted heteroaromatic ring selected from pyridine, pyrimidines, pyrazine, pyrimidine, pyrazole, pyrrole, imidazole, benzimidazole, quinoline, isoquinoline, carbazole, indole, isoindole, furane, thiophene, oxazole and thiazole.

**24.** A method according to claim 22 or claim 23, wherein R1-R9 are independently selected from -H, hydroxy-C<sub>0</sub>-C<sub>20</sub>-alkyl, halo-C<sub>0</sub>-C<sub>20</sub>-alkyl, nitroso, formyl-C<sub>0</sub>-C<sub>20</sub>-alkyl, carboxyl-C<sub>0</sub>-C<sub>20</sub>-alkyl and esters and salts thereof, carbamoyl-C<sub>0</sub>-C<sub>20</sub>-alkyl, sulphy-C<sub>0</sub>-C<sub>20</sub>-alkyl and esters and salts thereof, sulphamoyl-C<sub>0</sub>-C<sub>20</sub>-alkyl, amino-C<sub>0</sub>-C<sub>20</sub>-alkyl, aryl-C<sub>0</sub>-C<sub>20</sub>-alkyl, heteroaryl-C<sub>0</sub>-C<sub>20</sub>-alkyl, C<sub>0</sub>-C<sub>20</sub>-alkyl, alkoxy-C<sub>0</sub>-C<sub>8</sub>-alkyl, carbonyl-C<sub>0</sub>-C<sub>6</sub>-alkoxy, and aryl-C<sub>0</sub>-C<sub>6</sub>-alkyl and C<sub>0</sub>-C<sub>20</sub>-alkylamide;

EP 1 109 965 B1

or one of R1-R9 is a bridging group  $-C_{n'}(R11) (R12)-(D)_p-C_m(R11) (R12) -$  bound to another moiety of the same general formula, wherein p is zero or one, D is selected from a heteroatom or a heteroatom-containing group, or is part of an aromatic or saturated homonuclear and heteronuclear ring, n' is an integer from 1 to 4, m' is an integer from 1 to 4, with the proviso that  $n'+m' \leq 4$ , R11 and R12 are each independently preferably selected from -H, NR13 and OR14, alkyl, aryl, optionally substituted, and R13 and R14 are each independently selected from -H, alkyl, aryl, both optionally substituted.


25. A method according to any of claims 22 to 24, wherein T1 and T2 together form a single bond linkage and  $s > 1$ , according to general formula (BII):




wherein Z3 independently represents a group as defined for Z1 or Z2; R3 independently represents a group as defined for R1-R9; Q3 independently represents a group as defined for Q1, Q2; h represents zero or an integer from 1 to 6; and  $s' = s - 1$ .


26. A method according to claim 25, wherein in general formula (BII),  $s' = 1, 2$  or  $3$ ;  $r = g = h = 1$ ;  $d = 2$  or  $3$ ;  $e = f = 0$ ;  $R6 = R7 = H$ .

27. A method according to claim 26, wherein the ligand has a general formula selected from:





28. A method according to claim 27, wherein the ligand has a general formula selected from:



29. A method according to claim 28, wherein R1, R2, R3 and R4 are independently selected from -H, alkyl, heteroaryl, or represents a bridging group bound to another moiety of the same general formula with the bridging group being alkylene or hydroxy-alkylene or a heteroaryl-containing bridge.

30. A method according to claim 29, wherein R1, R2, R3 and R4 are independently selected from -H, methyl, ethyl, isopropyl, nitrogen-containing heteroaryl, or a bridging group bound to another moiety of the same general formula with the bridging group being alkylene or hydroxy-alkylene.

31. A method according to any of claims 26 to 30, wherein in the complex  $[M_a L_k X_n] Y_m$ :

M= Mn(II)-(IV), Cu(I)-(III), Fe(II)-(III), Co(II)-(III);

X=  $CH_3CN$ ,  $OH_2$ ,  $Cl^-$ ,  $Br^-$ ,  $OCN^-$ ,  $N_3^-$ ,  $SCN^-$ ,  $OH^-$ ,  $O^{2-}$ ,  $PO_4^{3-}$ ,  $C_6H_5BO_2^{2-}$ ,  $RCOO^-$ ;

Y=  $ClO_4^-$ ,  $BPh_4^-$ ,  $Br^-$ ,  $Cl^-$ ,  $[FeCl_4]^-$ ,  $PF_6^-$ ,  $NO_3^-$

a= 1, 2, 3, 4;


n= 0, 1, 2, 3, 4, 5, 6, 7, 8, 9;

m= 1, 2, 3, 4; and

k= 1, 2, 4.

32. A method according to claim 25, wherein in general formula (BII),  $s'=2$ ;  $r=g=h=1$ ;  $d=f=0$ ;  $e=1$ ; and each Y1 is independently alkylene or heteroarylene.

33. A method according to claim 32, wherein the ligand has the general formula:




wherein

A<sub>1</sub>, A<sub>2</sub>, A<sub>3</sub>, A<sub>4</sub> are independently selected from C<sub>1-9</sub>-alkylene or heteroarylene groups; and  
N<sub>1</sub> and N<sub>2</sub> independently represent a hetero atom or a heteroarylene group.

- 5 **34.** A method according to claim 33, wherein  
N<sub>1</sub> represents an aliphatic nitrogen;  
N<sub>2</sub> represents a heteroarylene group;  
R<sub>1</sub>, R<sub>2</sub>, R<sub>3</sub>, R<sub>4</sub> each independently represent -H, alkyl, aryl or heteroaryl; and  
10 A<sub>1</sub>, A<sub>2</sub>, A<sub>3</sub>, A<sub>4</sub> each represent -CH<sub>2</sub>-.

- 35.** A method according to claim 34, wherein the ligand has the general formula:




wherein R<sub>1</sub>, R<sub>2</sub> each independently represent -H, alkyl, aryl or heteroaryl.

- 36.** A method according to any of claims 32 to 35, wherein in the complex [M<sub>a</sub>L<sub>k</sub>X<sub>n</sub>]<sub>Y<sub>m</sub></sub>:

M= Fe(II)-(III), Mn(II)-(IV), Cu(II), Co(II)-(III);  
X= CH<sub>3</sub>CN, OH<sub>2</sub>, Cl<sup>-</sup>, Br<sup>-</sup>, OCN<sup>-</sup>, N<sub>3</sub><sup>-</sup>, SCN<sup>-</sup>, OH<sup>-</sup>, O<sup>2-</sup>, PO<sub>4</sub><sup>3-</sup>, C<sub>6</sub>H<sub>5</sub>BO<sub>2</sub><sup>2-</sup>, RCOO<sup>-</sup>;  
Y= ClO<sub>4</sub><sup>-</sup>, BPh<sub>4</sub><sup>-</sup>, Br<sup>-</sup>, Cl<sup>-</sup>, [FeCl<sub>4</sub>]<sup>-</sup>, PF<sub>6</sub><sup>-</sup>, NO<sub>3</sub><sup>-</sup>;  
a= 1, 2, 3, 4;  
n= 0, 1, 2, 3, 4, 5, 6, 7, 8, 9;  
35 m= 1, 2, 3, 4; and  
k= 1, 2, 4.


- 37.** A method according to claim 25, wherein in general formula (BII), s'=2 and r=g=h=1, according to the general formula:



- 38.** A method according to claim 37, wherein Z<sub>1</sub>=Z<sub>2</sub>=Z<sub>3</sub>=Z<sub>4</sub>=a heteroaromatic ring; e=f=0; d=1; and R<sub>7</sub> is absent.

- 39.** A method according to claim 37, wherein Z<sub>1</sub>-Z<sub>4</sub> each represent N; R<sub>1</sub>-R<sub>4</sub> are absent; both Q<sub>1</sub> and Q<sub>3</sub> represent =CH-[Y<sub>1</sub>]<sub>e</sub>-CH=; and both Q<sub>2</sub> and Q<sub>4</sub> represent -CH<sub>2</sub>-[Y<sub>1</sub>]<sub>n</sub>-CH<sub>2</sub>-.

40. A method according to claim 39, wherein the ligand has the general formula:



wherein A represents optionally substituted alkylene optionally interrupted by a heteroatom; and n is zero or an integer from 1 to 5.

25 41. A method according to claim 40 wherein R1-R6 represent hydrogen, n=1 and A= -CH<sub>2</sub>-, -CHOH-, -CH<sub>2</sub>N(R)CH<sub>2</sub>- or -CH<sub>2</sub>CH<sub>2</sub>N(R)CH<sub>2</sub>CH<sub>2</sub>- wherein R represents hydrogen or alkyl.

42. A method according to claim 41, wherein A= -CH<sub>2</sub>-, -CHOH- or -CH<sub>2</sub>CH<sub>2</sub>NHCH<sub>2</sub>CH<sub>2</sub>-.

30 43. A method according to any of claims 37 to 42 wherein in the complex [M<sub>a</sub>L<sub>k</sub>X<sub>n</sub>]Y<sub>m</sub>:

M= Mn(II)-(IV), Co(II)-(III), Fe(II)-(III);

X= CH<sub>3</sub>CN, OH<sub>2</sub>, Cl<sup>-</sup>, Br<sup>-</sup>, OCN<sup>-</sup>, N<sub>3</sub><sup>-</sup>, SCN<sup>-</sup>, OH<sup>-</sup>, O<sup>2-</sup>, PO<sub>4</sub><sup>3-</sup>, C<sub>6</sub>H<sub>5</sub>BO<sub>2</sub><sup>2-</sup>, RCOO<sup>-</sup>;

Y= ClO<sub>4</sub><sup>-</sup>, BPh<sub>4</sub><sup>-</sup>, Br<sup>-</sup>, Cl<sup>-</sup>, [FeCl<sub>4</sub>]<sup>-</sup>, PF<sub>6</sub><sup>-</sup>, NO<sub>3</sub><sup>-</sup>;


35 a= 1, 2, 3, 4;

n= 0, 1, 2, 3, 4, 5, 6, 7, 8, 9;

m= 1, 2, 3, 4; and

k= 1, 2, 4.


40 44. A method according to any of claims 22 to 24, wherein T1 and T2 independently represent groups R4, R5 as defined for R1-R9, according to the general formula (BIII):




50 45. A method according to claim 44, wherein in general formula (BIII), s=1; r=1; g=0; d=f=1; e=1-4; Y1= -CH<sub>2</sub>-; and R1 together with R4, and/or R2 together with R5, independently represent =CH-R10, wherein R10 is as defined for R1-R9.

46. A method according to claim 45, wherein R2 together with R5 represents =CH-R10.

55 47. A method according to claim 45 or claim 46, wherein the ligand is selected from:



48. A method according to claim 47, wherein the ligand is selected from:



wherein R1 and R2 are selected from optionally substituted phenols, heteroaryl-C<sub>0</sub>-C<sub>20</sub>-alkyls, R3 and R4 are selected from -H, alkyl, aryl, optionally substituted phenols, heteroalkyl-C<sub>0</sub>-C<sub>20</sub>-alkyls, alkylaryl, aminoalkyl, alkoxy.

49. A method according to claim 48 wherein R1 and R2 are selected from optionally substituted phenols, heteroaryl-C<sub>0</sub>-C<sub>2</sub>-alkyls, R3 and R4 are selected from -H, alkyl, aryl, optionally substituted phenols, nitrogen-heteroaryl-C<sub>0</sub>-C<sub>2</sub>-alkyls.

50. A method according to any of claims 45 to 49 wherein in the complex [M<sub>a</sub>L<sub>k</sub>X<sub>n</sub>]Y<sub>m</sub>:

M= Mn(II)-(IV), Co(II)-(III), Fe(II)-(III);

X= CH<sub>3</sub>CN, OH<sub>2</sub>, Cl<sup>-</sup>, Br<sup>-</sup>, OCN<sup>-</sup>, N<sub>3</sub><sup>-</sup>, SCN<sup>-</sup>, OH<sup>-</sup>, O<sup>2-</sup>, PO<sub>4</sub><sup>3-</sup>, C<sub>6</sub>H<sub>5</sub>BO<sub>2</sub><sup>2-</sup>, RCOO<sup>-</sup>;

Y= ClO<sub>4</sub><sup>-</sup>, BPh<sub>4</sub><sup>-</sup>, Br<sup>-</sup>, Cl<sup>-</sup>, [FeCl<sub>4</sub>], PF<sub>6</sub><sup>-</sup>, NO<sub>3</sub><sup>-</sup>;

a= 1, 2, 3, 4;


n= 0, 1, 2, 3, 4, 5, 6, 7, 8, 9;

m= 1, 2, 3, 4; and

k= 1, 2, 4.

51. A method according to claim 44, wherein in general formula (BIII), s=1; r=1; g=0; d=f=1; e=1-4; Y1= -C(R')(R''), wherein R' and R'' are independently as defined for R1-R9.

52. A method according to claim 51, wherein the ligand has the general formula:



53. A method according to claim 52, wherein R1, R2, R3, R4, R5 are -H or C<sub>0</sub>-C<sub>20</sub>-alkyl, n=0 or 1, R6 is -H, alkyl, -OH or -SH, and R7, R8, R9, R10 are each independently selected from -H, C<sub>0</sub>-C<sub>20</sub>-alkyl, heteroaryl-C<sub>0</sub>-C<sub>20</sub>-alkyl, alkoxy-C<sub>0</sub>-C<sub>8</sub>-alkyl and amino-C<sub>0</sub>-C<sub>20</sub>-alkyl.

54. A method according to any of claims 51 to 53 wherein in the complex [M<sub>a</sub>L<sub>k</sub>X<sub>n</sub>]Y<sub>m</sub>:

M= Mn(II)-(IV), Fe(II)-(III), Cu(II), Co(II)-(III);

EP 1 109 965 B1

X= CH<sub>3</sub>CN, OH<sub>2</sub>, Cl<sup>-</sup>, Br<sup>-</sup>, OCN<sup>-</sup>, N<sub>3</sub><sup>-</sup>, SCN<sup>-</sup>, OH<sup>-</sup>, O<sup>2-</sup>, PO<sub>4</sub><sup>3-</sup>, C<sub>6</sub>H<sub>5</sub>BO<sub>2</sub><sup>2-</sup>, RCOO<sup>-</sup>;

Y= ClO<sub>4</sub><sup>-</sup>, BPh<sub>4</sub><sup>-</sup>, Br<sup>-</sup>, Cl<sup>-</sup>, [FeCl<sub>4</sub>]<sup>-</sup>, PF<sub>6</sub><sup>-</sup>, NO<sub>3</sub><sup>-</sup>;

a= 1, 2, 3, 4;


n= 0, 1, 2, 3, 4;

m= 0, 1, 2, 3, 4, 5, 6, 7, 8; and

k= 1, 2, 3, 4.


55. A method according to claim 44, wherein in general formula (BIII), s=0; g=1; d=e=0; f=1-4.

56. A method according to claim 55, wherein the ligand has the general formula:



57. A method according to claim 56, with the proviso that none of R1 to R3 represents hydrogen.

58. A method according to claim 56 or claim 57, wherein the ligand has the general formula:



wherein R1, R2, R3 are as defined for R2, R4, R5.

59. A method according to any of claims 55 to 58, wherein in the complex [M<sub>a</sub>L<sub>k</sub>X<sub>n</sub>]Y<sub>m</sub>:

M= Mn(II)-(IV), Fe(II)-(III), Cu(II), Co(II)-(III);

X= CH<sub>3</sub>CN, OH<sub>2</sub>, Cl<sup>-</sup>, Br<sup>-</sup>, OCN<sup>-</sup>, N<sub>3</sub><sup>-</sup>, SCN<sup>-</sup>, OH<sup>-</sup>, O<sup>2-</sup>, PO<sub>4</sub><sup>3-</sup>, C<sub>6</sub>H<sub>5</sub>BO<sub>2</sub><sup>2-</sup>, RCOO<sup>-</sup>;

Y= ClO<sub>4</sub><sup>-</sup>, BPh<sub>4</sub><sup>-</sup>, Br<sup>-</sup>, Cl<sup>-</sup>, [FeCl<sub>4</sub>]<sup>-</sup>, PF<sub>6</sub><sup>-</sup>, NO<sub>3</sub><sup>-</sup>;


a= 1, 2, 3, 4;

n= 0, 1, 2, 3, 4;

m= 0, 1, 2, 3, 4, 5, 6, 7, 8; and

k= 1, 2, 3, 4.

60. A method according to any of claims 19 to 24, wherein L represents a pentadentate ligand of the general formula (B):



wherein

EP 1 109 965 B1

each R<sup>1</sup>, R<sup>2</sup> independently represents -R<sup>4</sup>-R<sup>5</sup>,

R<sup>3</sup> represents hydrogen, optionally substituted alkyl, aryl or arylalkyl, or -R<sup>4</sup>-R<sup>5</sup>,

each R<sup>4</sup> independently represents a single bond or optionally substituted alkylene, alkenylene, oxyalkylene, aminoalkylene, alkylene ether, carboxylic ester or carboxylic amide, and

each R<sup>5</sup> independently represents an optionally N-substituted aminoalkyl group or an optionally substituted heteroaryl group selected from pyridinyl, pyrazinyl, pyrazolyl, pyrrolyl, imidazolyl, benzimidazolyl, pyrimidinyl, triazolyl and thiazolyl.

61. A method according to claim 60, with the proviso that R<sup>3</sup> does not represent hydrogen.

62. A method according to any of claims 19 to 24, wherein L represents a pentadentate or hexadentate ligand of the general formula (C):




wherein

each R<sup>1</sup> independently represents -R<sup>3</sup>-V, in which R<sup>3</sup> represents optionally substituted alkylene, alkenylene, oxyalkylene, aminoalkylene or alkylene ether, and V represents an optionally substituted heteroaryl group selected from pyridinyl, pyrazinyl, pyrazolyl, pyrrolyl, imidazolyl, benzimidazolyl, pyrimidinyl, triazolyl and thiazolyl;

W represents an optionally substituted alkylene bridging group selected from -CH<sub>2</sub>CH<sub>2</sub>-, -CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>-, -CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>-, -CH<sub>2</sub>-C<sub>6</sub>H<sub>4</sub>-CH<sub>2</sub>-, -CH<sub>2</sub>-C<sub>6</sub>H<sub>10</sub>-CH<sub>2</sub>-, and -CH<sub>2</sub>-C<sub>10</sub>H<sub>6</sub>-CH<sub>2</sub>-; and

R<sup>2</sup> represents a group selected from R<sup>1</sup>, and alkyl, aryl and arylalkyl groups optionally substituted with a substituent selected from hydroxy, alkoxy, phenoxy, carboxylate, carboxamide, carboxylic ester, sulphonate, amine, alkylamine and N<sup>+</sup>(R<sup>4</sup>)<sub>3</sub>, wherein R<sup>4</sup> is selected from hydrogen, alkanyl, alkenyl, arylalkanyl, arylalkenyl, oxyalkanyl, oxyalkenyl, aminoalkanyl, aminoalkenyl, alkanyl ether and alkenyl ether.

63. A method according to any of claims 19 to 24, wherein L represents a macrocyclic ligand of formula (E):



wherein


Z<sup>1</sup> and Z<sup>2</sup> are independently selected from monocyclic or polycyclic aromatic ring structures optionally containing one or more heteroatoms, each aromatic ring structure being substituted by one or more substituents;

Y<sup>1</sup> and Y<sup>2</sup> are independently selected from C, N, O, Si, P and S atoms;

A<sup>1</sup> and A<sup>2</sup> are independently selected from hydrogen, alkyl, alkenyl and cycloalkyl (each of alkyl, alkenyl and cycloalkyl) being optionally substituted by one or more groups selected from hydroxy, aryl, heteroaryl, sulphonate, phosphate, electron donating groups and electron withdrawing groups, and groups of formulae (G<sup>1</sup>)(G<sup>2</sup>)N-, G<sup>3</sup>OC(O)-, G<sup>3</sup>O- and G<sup>3</sup>C(O)-, wherein each of G<sup>1</sup>, G<sup>2</sup> and G<sup>3</sup> is independently selected from hydrogen and alkyl, and electron donating and/or withdrawing groups (in addition to any amongst the foregoing);

i and j are selected from 0, 1 and 2 to complete the valency of the groups Y<sup>1</sup> and Y<sup>2</sup>;

each of Q<sup>1</sup>-Q<sup>4</sup> is independently selected from groups of formula





wherein  $10 > a + b + c + d > 2$ ;

each  $Y^3$  is independently selected from -O-, -S-, -SO-, -SO<sub>2</sub>-, -(G<sup>1</sup>)(G<sup>2</sup>)N-, -(G<sup>1</sup>)N- (wherein G<sup>1</sup> and G<sup>2</sup> are as hereinbefore defined), -C(O)-, aryl, heteroaryl, -P- and -P(O)-;

each of A<sup>3</sup>-A<sup>6</sup> is independently selected from the groups hereinbefore defined for A<sup>1</sup> and A<sup>2</sup>; and

wherein any two or more of A<sup>1</sup>-A<sup>6</sup> together form a bridging group, provided that if A<sup>1</sup> and A<sup>2</sup> are linked without simultaneous linking also to any of A<sup>3</sup>-A<sup>6</sup>, then the bridging group linking A<sup>1</sup> and A<sup>2</sup> must contain at least one carbonyl group.

## Patentansprüche

1. Verfahren zum Behandeln eines Textils durch In-Kontakt-Bringen des Textils mit einer organischen Substanz, die einen Komplex mit einem Übergangsmetall bildet, wobei die Behandlung die organische Substanz in einer Form bereitstellt, die im Wesentlichen kein Persauerstoffbleichmittel oder auf Peroxy basierendes oder erzeugendes Bleichmittelsystem aufweist, wobei der Komplex Bleichen des Textils durch atmosphärischen Sauerstoff nach der Behandlung katalysiert, wobei das behandelte Textil getrocknet wird und Bleichen auf dem trockenen Textil katalysiert wird.
2. Verfahren nach Anspruch 1, wobei die Behandlung In-Kontakt-Bringen des Textils mit der organischen Substanz in trockener Form umfasst.
3. Verfahren nach Anspruch 1, wobei die Behandlung In-Kontakt-Bringen des Textils mit einer die organische Substanz enthaltenden Flüssigkeit und dann Trocknen umfasst.
4. Verfahren nach Anspruch 3, wobei die Flüssigkeit eine wässrige Flüssigkeit ist.
5. Verfahren nach Anspruch 4, wobei die Flüssigkeit ein aufzusprühendes Textilbehandlungsfluid ist.
6. Verfahren nach Anspruch 4, wobei die Flüssigkeit eine Waschflüssigkeit zum Wäschereinigen ist.
7. Verfahren nach Anspruch 3, wobei die Flüssigkeit eine nichtwässrige Flüssigkeit ist.
8. Verfahren nach Anspruch 7, wobei die Flüssigkeit ein trockenes Reinigungsfluid ist.
9. Verfahren nach Anspruch 7, wobei die Flüssigkeit ein aufzusprühendes Aerosolfluid ist.
10. Verfahren nach Anspruch 3, wobei das Medium einen pH-Wert im Bereich von pH 6 bis 11 aufweist.
11. Verfahren nach Anspruch 10, wobei die Flüssigkeit einen pH-Wert im Bereich von pH 8 bis 10 aufweist.
12. Verfahren nach einem der Ansprüche 3 bis 11, wobei die Flüssigkeit im Wesentlichen kein Übergangsmetallmaskierungsmittel aufweist.
13. Verfahren nach einem der Ansprüche 3 bis 12, wobei die Flüssigkeit weiterhin ein Tensid umfasst.
14. Verfahren nach einem der Ansprüche 3 bis 13, wobei die Flüssigkeit weiterhin einen Builder umfasst.
15. Verfahren nach einem der Ansprüche 1 bis 14, wobei die organische Substanz einen vorgebildeten Komplex eines Liganden und eines Übergangsmetalls umfasst.
16. Verfahren nach einem der Ansprüche 3 bis 14, wobei die organische Substanz einen freien Liganden umfasst, der mit einem in der Flüssigkeit vorliegenden Übergangsmetall komplexiert.
17. Verfahren nach einem der Ansprüche 1 bis 14, wobei die organische Substanz einen freien Liganden umfasst, der mit einem in dem Textil vorliegenden Übergangsmetall komplexiert.
18. Verfahren nach einem der Ansprüche 1 bis 14, wobei die organische Substanz eine Zusammensetzung eines freien Liganden oder eines Übergangsmetall-substituierbaren Metallligandenkomplexes und eine Quelle von Über-

gangsmetall umfasst.

19. Verfahren nach einem der Ansprüche 1 bis 18, wobei die organische Substanz einen Komplex der allgemeinen Formel (A1) bildet:



worin:

M ein Metall wiedergibt, ausgewählt aus Mn(II)-(III)-(IV)-(V), Cu(I)-(II)-(III), Fe(I)-(II)-(III)-(IV), Co(I)-(II)-(III), Ni(I)-(II)-(III), Cr(II)-(III)-(IV)-(V)-(VI)-(VII), Ti(II)-(III)-(IV), V(II)-(III)-(IV)-(V), Mo(II)-(III)-(IV)-(V)-(VI), W(IV)-(V)-(VI), Pd(II), Ru(II)-(III)-(IV)-(V) und Ag(I)-(II),

L einen Liganden oder sein protoniertes oder deprotoniertes Analoges wiedergibt;

X eine koordinierende Spezies, ausgewählt aus beliebigen einfach, doppelt oder dreifach geladenen Anionen und beliebigen neutralen Molekülen, die das Metall in einer ein-, zwei- oder dreizähligen Weise koordinieren können, wiedergibt;

Y beliebiges nicht-koordiniertes Gegenion wiedergibt;

a eine ganze Zahl von 1 bis 10 wiedergibt;

k eine ganze Zahl von 1 bis 10 wiedergibt;

n null oder eine ganze Zahl von 1 bis 10 wiedergibt; und

m null oder eine ganze Zahl von 1 bis 20 wiedergibt.

20. Verfahren nach Anspruch 19, wobei in Formel (A1):

X eine koordinierende Spezies wiedergibt, ausgewählt aus  $O^{2-}$ ,  $RBO_2^{2-}$ ,  $RCOO^-$ ,  $RCONR^-$ ,  $OH^-$ ,  $NO_3^-$ ,  $NO_2^-$ ,  $NO$ ,  $CO$ ,  $S^{2-}$ ,  $RS^-$ ,  $PO_4^{3-}$ , STP-abgeleiteten Anionen,  $PO_3OR^{3-}$ ,  $H_2O$ ,  $CO_3^{2-}$ ,  $HCO_3^-$ ,  $ROH$ ,  $NRR'R''$ ,  $RCN$ ,  $Cl^-$ ,  $Br^-$ ,  $OCN^-$ ,  $SCN^-$ ,  $CN^-$ ,  $N_3^-$ ,  $F^-$ ,  $I^-$ ,  $RO^-$ ,  $ClO_4^-$ ,  $SO_4^{2-}$ ,  $HSO_4^-$ ,  $SO_3^{2-}$  und  $RSO_3^-$ ; und

Y ein Gegenion wiedergibt, ausgewählt aus  $ClO_4^-$ ,  $BR_4^-$ ,  $[FeCl_4]^-$ ,  $PF_6^-$ ,  $RCOO^-$ ,  $NO_3^-$ ,  $NO_2^-$ ,  $RO^-$ ,  $N^+RR'R''R'''$ ,  $Cl^-$ ,  $Br^-$ ,  $F^-$ ,  $I^-$ ,  $RSO_3^-$ ,  $S_2O_6^{2-}$ ,  $OCN^-$ ,  $SCN^-$ ,  $Li^+$ ,  $Ba^{2+}$ ,  $Na^+$ ,  $Mg^{2+}$ ,  $K^+$ ,  $Ca^{2+}$ ,  $Cs^+$ ,  $PR_4^+$ ,  $RBO_2^{2-}$ ,  $SO_4^{2-}$ ,  $HSO_4^-$ ,  $SO_3^{2-}$ ,  $SbCl_6^-$ ,  $CuCl_4^{2-}$ ,  $CN$ ,  $PO_4^{3-}$ ,  $HPO_4^{2-}$ ,  $H_2PO_4^-$ , STP-abgeleiteten Anionen,  $CO_3^{2-}$ ,  $HCO_3^-$  und  $BF_4^-$ ;

worin

R, R', R'', R''' unabhängig eine Gruppe wiedergeben, ausgewählt aus Wasserstoff, Hydroxyl, -OR (worin R= Alkyl-, Alkenyl-, Cycloalkyl-, Heterocycloalkyl-, Aryl-, Heteroaryl- oder Carbonylderivatgruppe), -OAr, Alkyl-, Alkenyl-, Cycloalkyl-, Heterocycloalkyl-, Aryl-, Heteroaryl- und Carbonylderivatgruppen, wobei jede von R, Ar, Alkyl-, Alkenyl-, Cycloalkyl-, Heterocycloalkyl-, Aryl-, Heteroaryl- und Carbonylderivatgruppen gegebenenfalls mit einer oder mehreren funktionellen Gruppen E substituiert ist, oder R6 zusammen mit R7 und unabhängig R8 zusammen mit R9 Sauerstoff wiedergeben;

E ausgewählt ist aus funktionellen Gruppen, die Sauerstoff, Schwefel, Phosphor, Stickstoff, Selen, Halogene, und beliebige Elektronen-abgebende und/oder -anziehende Gruppen enthalten.

21. Verfahren nach Anspruch 19 oder Anspruch 20, wobei in Formel (A1):

M ein Metall wiedergibt, ausgewählt aus Mn(II)-(III)-(IV)-(V), Cu(I)-(II), Fe(II)-(III)-(IV) und Co(I)-(II)-(III);

X eine koordinierende Spezies wiedergibt, ausgewählt aus  $O^{2-}$ ,  $RBO_2^{2-}$ ,  $RCOO^-$ ,  $OH^-$ ,  $NO_3^-$ ,  $NO_2^-$ ,  $NO$ ,  $CO$ ,  $CN^-$ ,  $S^{2-}$ ,  $RS^-$ ,  $PO_4^{3-}$ ,  $H_2O$ ,  $CO_3^{2-}$ ,  $HCO_3^-$ ,  $ROH$ ,  $NRR'R''$ ,  $Cl^-$ ,  $Br^-$ ,  $OCN^-$ ,  $SCN^-$ ,  $RCN$ ,  $N_3^-$ ,  $F^-$ ,  $I^-$ ,  $RO^-$ ,  $ClO_4^-$ ,  $SO_4^{2-}$ ,  $HSO_4^-$ ,  $SO_3^{2-}$  und  $RSO_3^-$ ;

Y ein Gegenion wiedergibt, ausgewählt aus  $ClO_4^-$ ,  $BR_4^-$ ,  $[FeCl_4]^-$ ,  $PF_6^-$ ,  $RCOO^-$ ,  $NO_3^-$ ,  $NO_2^-$ ,  $RO^-$ ,  $N^+RR'R''R'''$ ,  $Cl^-$ ,  $Br^-$ ,  $F^-$ ,  $I^-$ ,  $RSO_3^-$ ,  $S_2O_6^{2-}$ ,  $OCN^-$ ,  $SCN^-$ ,  $Li^+$ ,  $Ba^{2+}$ ,  $Na^+$ ,  $Mg^{2+}$ ,  $K^+$ ,  $Ca^{2+}$ ,  $PR_4^+$ ,  $SO_4^{2-}$ ,  $HSO_4^-$ ,  $SO_3^{2-}$  und  $BF_4^-$ ; worin

R, R', R'', R''' Wasserstoff, gegebenenfalls substituiertes Alkyl oder gegebenenfalls substituiertes Aryl wiedergeben;


a eine ganze Zahl von 1 bis 4 wiedergibt;

k eine ganze Zahl von 1 bis 10 wiedergibt;

n null oder eine ganze Zahl von 1 bis 4 wiedergibt; und

m null oder eine ganze Zahl von 1 bis 8 wiedergibt.

22. Verfahren nach einem der Ansprüche 19 bis 21, wobei L einen Liganden der allgemeinen Formel (BI) wiedergibt:



worin


g null oder eine ganze Zahl von 1 bis 6 wiedergibt;

r eine ganze Zahl von 1 bis 6 wiedergibt;

s null oder eine ganze Zahl von 1 bis 6 wiedergibt;

Z1 und Z2 unabhängig ein Heteroatom oder einen heterocyclischen oder heteroaromatischen Ring wiedergeben, Z1 und/oder Z2 gegebenenfalls mit einer oder mehreren funktionellen Gruppen E, wie nachstehend definiert, substituiert sind;

Q1 und Q2 unabhängig eine Gruppe der Formel wiedergeben:



worin :

10>d+e+f>1; d=0-9; e=0-9; f=0-9;

jedes Y1 unabhängig ausgewählt ist aus -O-, -S-, -SO-, -SO<sub>2</sub>-, -(G<sup>1</sup>)N-, -(G<sup>1</sup>)(G<sup>2</sup>)N- (worin G<sup>1</sup> und G<sup>2</sup> wie nachstehend definiert sind), -C(O)-, Arylen, Alkylen, Heteroarylen, -P- und -P(O)-;

wenn s>1, jede Gruppe -[Z1(R1)-(Q1)<sub>r</sub>]- unabhängig definiert ist;

R1, R2, R6, R7, R8, R9 unabhängig eine Gruppe wiedergeben, ausgewählt aus Wasserstoff, Hydroxyl, -OR (worin R= Alkyl-, Alkenyl-, Cycloalkyl-, Heterocycloalkyl-, Aryl-, Heteroaryl- oder Carbonylderivatgruppe), -OAr, Alkyl-, Alkenyl-, Cycloalkyl-, Heterocycloalkyl-, Aryl-, Heteroaryl- und Carbonylderivatgruppen, wobei jede von R, Ar, Alkyl-, Alkenyl-, Cycloalkyl-, Heterocycloalkyl-, Aryl-, Heteroaryl- und Carbonylderivatgruppen gegebenenfalls mit einer oder mehreren funktionellen Gruppen E substituiert ist, oder R6 zusammen mit R7 und unabhängig R8 zusammen mit R9 Sauerstoff wiedergeben;

E ausgewählt ist aus funktionellen Gruppen, enthaltend Sauerstoff, Schwefel, Phosphor, Stickstoff, Selen, Halogene, und beliebige Elektronen-abgebende und/oder -anziehende Gruppen (vorzugsweise ist E ausgewählt aus Hydroxy, Mono- oder Polycarboxylatderivaten, Aryl, Heteroaryl, Sulfonat, Thiol (-RSH), Thioethern (-R-S-R'), Disulfiden (-RSSR'), Dithiolenen, Mono- oder Polyphosphonaten, Mono- oder Polyphosphaten, Elektronen-abgebenden Gruppen und Elektronen-anziehenden Gruppen, und Gruppen der Formeln (G<sup>1</sup>)(G<sup>2</sup>)N-, (G<sup>1</sup>)(G<sup>2</sup>)(G<sup>3</sup>)N-, (G<sup>1</sup>)(G<sup>2</sup>)N-C(O)-, G<sup>3</sup>O- und G<sup>3</sup>C(O)-, worin jeder von G<sup>1</sup>, G<sup>2</sup> und G<sup>3</sup> unabhängig ausgewählt ist aus Wasserstoff, Alkyl, Elektronen-abgebenden Gruppen und Elektronen-anziehenden Gruppen, (zusätzlich zu beliebigen der vorangehenden));

oder einer von R1-R9 eine Brückengruppe darstellt, die an eine weitere Einheit der gleichen allgemeinen Formel gebunden ist;

T1 und T2 unabhängig Gruppen R4 und R5 wiedergeben, worin R4 und R5 wie für R1-R9 definiert sind, und, wenn g=0 und s>0, R1 zusammen mit R4 und/oder R2 zusammen mit R5 gegebenenfalls unabhängig =CH-R10 wiedergeben können, worin R10 wie für R1-R9 definiert ist, oder

T1 und T2 zusammen (-T2-T1-) eine kovalente Bindung wiedergeben, wenn s>1 und g>0;

wenn Z1 und/oder Z2 N wiedergibt und T1 und T2 zusammen eine Einfachbindungsverknüpfung wiedergeben, und R1 und/oder R2 nicht vorliegen, Q1 und/oder Q2 unabhängig eine Gruppe der Formel: =CH-[-Y1-]<sub>e</sub>-CH= wiedergeben,

gegebenenfalls zwei oder mehrere von R1, R2, R6, R7, R8, R9 unabhängig zusammen durch eine kovalente Bindung gebunden sind;

wenn Z1 und/oder Z2 0 wiedergeben, dann R1 und/oder R2 nicht vorliegen;

wenn Z1 und/oder Z2 S, N, P, B oder Si wiedergeben, dann R1 und/oder R2 nicht vorliegen können;

wenn Z1 und/oder Z2 ein Heteroatom, substituiert mit einer funktionellen Gruppe E, wiedergeben, dann R1


und/oder R2 und/oder R4 und/oder R5 nicht vorliegen können.

23. Verfahren nach Anspruch 22, wobei Z1 und Z2 unabhängig ein gegebenenfalls substituiertes Heteroatom, ausgewählt aus N, P, O, S, B und Si, oder einen gegebenenfalls substituierten heterocyclischen Ring oder einen gegebenenfalls substituierten heteroaromatischen Ring, ausgewählt aus Pyridin, Pyrimidinen, Pyrazin, Pyrimidin, Pyrazol, Pyrrol, Imidazol, Benzimidazol, Chinolein, Isochinolin, Carbazol, Indol, Isoindol, Furan, Thiophen, Oxazol und Thiazol wiedergeben.

24. Verfahren nach Anspruch 22 oder Anspruch 23, worin R1 bis R9 unabhängig ausgewählt sind aus -H, Hydroxy-C<sub>0</sub>-C<sub>20</sub>-alkyl, Halogen-C<sub>0</sub>-C<sub>20</sub>-alkyl, Nitroso, Formyl-C<sub>0</sub>-C<sub>20</sub>-alkyl, Carboxyl-C<sub>0</sub>-C<sub>20</sub>-alkyl und Estern und Salzen davon, Carbamoyl-C<sub>0</sub>-C<sub>20</sub>-alkyl, Sulfo-C<sub>0</sub>-C<sub>20</sub>-alkyl und Estern und Salzen davon, Sulfamoyl-C<sub>0</sub>-C<sub>20</sub>-alkyl, Amino-C<sub>0</sub>-C<sub>20</sub>-alkyl, Aryl-C<sub>0</sub>-C<sub>20</sub>-alkyl, Heteroaryl-C<sub>0</sub>-C<sub>20</sub>-alkyl, C<sub>0</sub>-C<sub>20</sub>-Alkyl, Alkoxy-C<sub>0</sub>-C<sub>8</sub>-alkyl, Carbonyl-C<sub>0</sub>-C<sub>6</sub>-alkoxy und Aryl-C<sub>0</sub>-C<sub>6</sub>-alkyl und C<sub>0</sub>-C<sub>20</sub>-Alkylamid;

oder einer von R1-R9 eine Brückengruppe -C<sub>n</sub>(R11)(R12)-(D)<sub>p</sub>-C<sub>m</sub>(R11)(R12)- darstellt, die an eine weitere Einheit der gleichen allgemeinen Formel gebunden ist, wobei p null oder eins ist, D ausgewählt ist aus einem Heteroatom oder einer ein Heteroatom enthaltenden Gruppe, oder Teil eines aromatischen oder gesättigten homokernigen und heterokernigen Rings ist, n' eine ganze Zahl von 1 bis 4 ist, m' eine ganze Zahl von 1 bis 4 ist, mit der Maßgabe, dass n'+m'<4, R11 und R12 jeweils unabhängig vorzugsweise ausgewählt sind aus -H, NR13 und OR14, Alkyl, Aryl, gegebenenfalls substituiert, und R13 und R14 unabhängig ausgewählt sind aus -H, Alkyl, Aryl, beide gegebenenfalls substituiert.


25. Verfahren nach einem der Ansprüche 22 bis 24, wobei T1 und T2 zusammen eine Einfachbindungsverknüpfung bilden und s>1 gemäß der allgemeinen Formel (BII):




worin Z3 unabhängig eine Gruppe, wie für Z1 oder Z2 definiert, wiedergibt; R3 unabhängig eine Gruppe, wie für R1-R9 definiert, wiedergibt; Q3 unabhängig eine Gruppe, wie für Q1, Q2 definiert, wiedergibt; h null oder eine ganze Zahl von 1 bis 6 wiedergibt; und s'=s-1.

26. Verfahren nach Anspruch 25, wobei in allgemeiner Formel (BII) s'=1, 2 oder 3; r=g=h=1; d=2 oder 3; e=f=0; R6=R7=H.


27. Verfahren nach Anspruch 26, wobei der Ligand eine allgemeine Formel aufweist, ausgewählt aus:



5




10



15

28. Verfahren nach Anspruch 27, wobei der Ligand eine allgemeine Formel aufweist, ausgewählt aus:

20



25

29. Verfahren nach Anspruch 28, wobei R1, R2, R3 und R4 unabhängig ausgewählt sind aus -H, Alkyl, Heteroaryl oder eine Brückengruppe wiedergeben, gebunden an eine weitere Einheit der gleichen allgemeinen Formel, wobei die Brückengruppe eine Alkylen- oder Hydroxy-alkylen- oder eine Heteroarylenthaltende Brücke darstellt.

30

30. Verfahren nach Anspruch 29, wobei R1, R2, R3 und R4 unabhängig ausgewählt sind aus -H, Methyl, Ethyl, Isopropyl, Stickstoff-enthaltendem Heteroaryl oder einer Brückengruppe, die an eine weitere Einheit der gleichen allgemeinen Formel gebunden ist, wobei die Brückengruppe Alkylen oder Hydroxyalkylen darstellt.

35

31. Verfahren nach einem der Ansprüche 26 bis 30, wobei in dem Komplex  $[M_a L_k X_n] Y_m$ :

40

- M= Mn(II)-(IV), Cu(I)-(III), Fe(II)-(III), Co(II)-(III);
- X= CH<sub>3</sub>CN, OH<sub>2</sub>, Cl<sup>-</sup>, Br<sup>-</sup>, OCN<sup>-</sup>, N<sub>3</sub><sup>-</sup>, SCN<sup>-</sup>, OH<sup>-</sup>, O<sup>2-</sup>, PO<sub>4</sub><sup>3-</sup>, C<sub>6</sub>H<sub>5</sub>BO<sub>2</sub><sup>2-</sup>, RCOO<sup>-</sup>;
- Y= ClO<sub>4</sub><sup>-</sup>, BPh<sub>4</sub><sup>-</sup>, Br<sup>-</sup>, Cl<sup>-</sup>, [FeCl<sub>4</sub>]<sup>-</sup>, PF<sub>6</sub><sup>-</sup>, NO<sub>3</sub><sup>-</sup>;
- a= 1, 2, 3, 4;
- n= 0, 1, 2, 3, 4, 5, 6, 7, 8, 9;
- m= 1, 2, 3, 4; und
- k= 1, 2, 4.


45

32. Verfahren nach Anspruch 25, wobei in allgemeiner Formel (BII) s'=2; r=g=h=1; d=f=0; e=1; und jedes Y1 unabhängig Alkylen oder Heteroarylen darstellt.

50

33. Verfahren nach Anspruch 32, wobei der Ligand die allgemeine Formel aufweist:

55




worin

$A_1, A_2, A_3, A_4$  unabhängig ausgewählt sind aus  $C_{1-9}$ -Alkylen- oder Heteroarylgruppen; und  $N_1$  und  $N_2$  unabhängig ein Heteroatom oder eine Heteroarylgruppe wiedergeben.

- 5 **34.** Verfahren nach Anspruch 33, wobei  
 $N_1$  einen aliphatischen Stickstoff wiedergibt;  
 $N_2$  eine Heteroarylgruppe wiedergibt;  
 $R_1, R_2, R_3, R_4$  jeweils unabhängig -H, Alkyl, Aryl oder Heteroaryl wiedergeben; und  
 $A_1, A_2, A_3, A_4$  jeweils  $-CH_2-$  wiedergeben.

- 10 **35.** Verfahren nach Anspruch 34, wobei der Ligand die allgemeine Formel aufweist:



worin  $R_1, R_2$  jeweils unabhängig -H, Alkyl, Aryl oder Heteroaryl wiedergeben.

- 25 **36.** Verfahren nach einem der Ansprüche 32 bis 35, wobei in dem Komplex  $[M_a L_k X_n] Y_m$ :

$M = Fe(II)-(III), Mn(II)-(IV), Cu(II), Co(II)-(III)$ ;

$X = CH_3CN, OH_2, Cl^-, Br^-, OCN^-, N_3^-, SCN^-, OH^-, O^{2-}, PO_4^{3-}, C_6H_5BO_2^{2-}, RCOO^-$ ;

$Y = ClO_4^-, BPh_4^-, Br^-, Cl^-, [FeCl_4]^-, PF_6^-, NO_3^-$ ;


30  $a = 1, 2, 3, 4$ ;

$n = 0, 1, 2, 3, 4, 5, 6, 7, 8, 9$ ;

$m = 1, 2, 3, 4$ ; und

$k = 1, 2, 4$ .

- 35 **37.** Verfahren nach Anspruch 25, wobei in allgemeiner Formel (BII)  $s'=2$  und  $r=g=h=1$ , gemäß der allgemeinen Formel:




- 38.** Verfahren nach Anspruch 37, wobei  $Z_1=Z_2=Z_3=Z_4$  ein heteroaromatischer Ring;  $e=f=0$ ;  $d=1$  und  $R_7$  nicht vorliegt.

- 50 **39.** Verfahren nach Anspruch 37, wobei  $Z_1-Z_4$  jeweils N wiedergeben;  $R_1-R_4$  nicht vorliegen; sowohl  $Q_1$  als auch  $Q_3 = CH-[-Y_1]_e-CH=$  wiedergeben; und sowohl  $Q_2$  als auch  $Q_4 -CH_2-[-Y_1]_n-CH_2-$  wiedergeben.

- 40.** Verfahren nach Anspruch 39, wobei der Ligand die allgemeine Formel aufweist:

55



worin A gegebenenfalls substituiertes Alkylen, gegebenenfalls unterbrochen durch ein Heteroatom, wiedergibt; und n null oder eine ganze Zahl von 1 bis 5 ist.

41. Verfahren nach Anspruch 40, wobei R1-R6 Wasserstoff wiedergeben, n=1 und A= -CH<sub>2</sub>-, -CHOH-, -CH<sub>2</sub>N(R)CH<sub>2</sub>- oder -CH<sub>2</sub>CH<sub>2</sub>N(R)CH<sub>2</sub>CH<sub>2</sub>-, worin R Wasserstoff oder Alkyl wiedergibt.

42. Verfahren nach Anspruch 41, wobei A= -CH<sub>2</sub>-, -CHOH- oder -CH<sub>2</sub>CH<sub>2</sub>NHCH<sub>2</sub>CH<sub>2</sub>-.

43. Verfahren nach einem der Ansprüche 37 bis 42, wobei in dem Komplex [M<sub>a</sub>L<sub>k</sub>X<sub>n</sub>]Y<sub>m</sub>:

M= Mn(II)-(IV), Co(II)-(III), Fe(II)-(III);

X= CH<sub>3</sub>CN, OH<sub>2</sub>, Cl<sup>-</sup>, Br<sup>-</sup>, OCN<sup>-</sup>, N<sub>3</sub><sup>-</sup>, SCN<sup>-</sup>, OH<sup>-</sup>, O<sup>2-</sup>, PO<sub>4</sub><sup>3-</sup>, C<sub>6</sub>H<sub>5</sub>BO<sub>2</sub><sup>2-</sup>, RCOO<sup>-</sup>;

Y= ClO<sub>4</sub><sup>-</sup>, BPh<sub>4</sub><sup>-</sup>, Br<sup>-</sup>, Cl<sup>-</sup>, [FeCl<sub>4</sub>]<sup>-</sup>, PF<sub>6</sub><sup>-</sup>, NO<sub>3</sub><sup>-</sup>;


a= 1, 2, 3, 4;

n= 0, 1, 2, 3, 4, 5, 6, 7, 8, 9;

m= 1, 2, 3, 4; und

k= 1, 2, 4.


44. Verfahren nach einem der Ansprüche 22 bis 24, wobei T1 und T2 unabhängig Gruppen R4 wiedergeben, R5 wie für R1-R9 definiert ist, gemäß der allgemeinen Formel (BIII):




45. Verfahren nach Anspruch 44, wobei in allgemeiner Formel (BIII) s=1, r=1, g=0, d=f=1, e=1-4, Y1= -CH<sub>2</sub>-; und R1 zusammen mit R4 und/oder R2 zusammen mit R5 unabhängig =CH-R10 wiedergeben, worin R10 wie für R1-R9 definiert ist.

46. Verfahren nach Anspruch 45, wobei R2 zusammen mit R5 =CH-R10 wiedergibt.

47. Verfahren nach Anspruch 45 oder Anspruch 46, wobei der Ligand ausgewählt ist aus:



48. Verfahren nach Anspruch 47, wobei der Ligand ausgewählt ist aus:



worin R1 und R2 ausgewählt sind aus gegebenenfalls substituierten Phenolen, Heteroaryl-C<sub>0</sub>-C<sub>20</sub>-alkylgruppen, R3 und R4 ausgewählt sind aus -H, Alkyl, Aryl, gegebenenfalls substituierten Phenolen, Heteroaryl-C<sub>0</sub>-C<sub>20</sub>-alkylgruppen, Alkylaryl, Aminoalkyl, Alkoxy.

49. Verfahren nach Anspruch 48, wobei R1 und R2 ausgewählt sind aus gegebenenfalls substituierten Phenolen, Heteroaryl-C<sub>0</sub>-C<sub>2</sub>-alkylgruppen, R3 und R4 ausgewählt sind aus -H, Alkyl, Aryl, gegebenenfalls substituierten Phenolen, Stickstoffheteroaryl-C<sub>0</sub>-C<sub>2</sub>-alkylgruppen.

50. Verfahren nach einem der Ansprüche 45 bis 49, wobei in dem Komplex  $[M_a L_k X_n] Y_m$ :

M= Mn(II)-(IV), Co(II)-(III), Fe(II)-(III);

X= CH<sub>3</sub>CN, OH<sub>2</sub>, Cl<sup>-</sup>, Br<sup>-</sup>, OCN<sup>-</sup>, N<sub>3</sub><sup>-</sup>, SCN<sup>-</sup>, OH<sup>-</sup>, O<sup>2-</sup>, PO<sub>4</sub><sup>3-</sup>, C<sub>6</sub>H<sub>5</sub>BO<sub>2</sub><sup>2-</sup>, RCOO<sup>-</sup>;

Y= ClO<sub>4</sub><sup>-</sup>, BPh<sub>4</sub><sup>-</sup>, Br<sup>-</sup>, Cl<sup>-</sup>, [FeCl<sub>4</sub>]<sup>-</sup>, PF<sub>6</sub><sup>-</sup>, NO<sub>3</sub><sup>-</sup>;

a= 1, 2, 3, 4;


n= 0, 1, 2, 3, 4, 5, 6, 7, 8, 9;

m= 1, 2, 3, 4; und

k= 1, 2, 4.

51. Verfahren nach Anspruch 44, wobei in allgemeiner Formel (BIII) s=1, r=1, g=0, d=f=1, e=1-4, Y1= -C(R') (R''), worin R' und R'' unabhängig wie für R1-R9 definiert sind.

52. Verfahren nach Anspruch 51, wobei der Ligand die allgemeine Formel aufweist:



53. Verfahren nach Anspruch 52, wobei R1, R2, R3, R4, R5 -H oder C<sub>0</sub>-C<sub>20</sub>-Alkyl darstellen, n=0 oder 1, R6 -H, Alkyl, -OH oder -SH darstellt, und R7, R8, R9, R10 jeweils unabhängig ausgewählt sind aus -H, C<sub>0</sub>-C<sub>20</sub>-Alkyl, Heteroaryl-C<sub>0</sub>-C<sub>20</sub>-alkyl, Alkoxy-C<sub>0</sub>-C<sub>5</sub>-alkyl und Amino-C<sub>0</sub>-C<sub>20</sub>-alkyl.

54. Verfahren nach einem der Ansprüche 51 bis 53, wobei in dem Komplex  $[M_a L_k X_n] Y_m$ :

M= Mn(II)-(IV), Fe(II)-(III), Cu(II), Co(II)-(III);

X= CH<sub>3</sub>CN, OH<sub>2</sub>, Cl<sup>-</sup>, Br<sup>-</sup>, OCN<sup>-</sup>, N<sub>3</sub><sup>-</sup>, SCN<sup>-</sup>, OH<sup>-</sup>, O<sup>2-</sup>, PO<sub>4</sub><sup>3-</sup>, C<sub>6</sub>H<sub>5</sub>BO<sub>2</sub><sup>2-</sup>, RCOO<sup>-</sup>;

Y= ClO<sub>4</sub><sup>-</sup>, BPh<sub>4</sub><sup>-</sup>, Br<sup>-</sup>, Cl<sup>-</sup>, [FeCl<sub>4</sub>]<sup>-</sup>, PF<sub>6</sub><sup>-</sup>, NO<sub>3</sub><sup>-</sup>;

a= 1, 2, 3, 4;

n= 0, 1, 2, 3, 4;


m= 0, 1, 2, 3, 4, 5, 6, 7, 8; und

k= 1, 2, 3, 4.

55. Verfahren nach Anspruch 44, wobei in allgemeiner Formel (BIII) s=0, g=1, d=e=0, f=1-4.

56. Verfahren nach Anspruch 55, wobei der Ligand die allgemeine Formel aufweist:






5

57. Verfahren nach Anspruch 56, mit der Maßgabe, dass keiner von R1 bis R3 Wasserstoff wiedergibt.

10

58. Verfahren nach Anspruch 56 oder Anspruch 57, wobei der Ligand die allgemeine Formel aufweist:



15

20

worin R1, R2, R3 wie für R2, R4, R5 definiert sind.

59. Verfahren nach einem der Ansprüche 55 bis 58, wobei in dem Komplex  $[M_a L_k X_n] Y_m$ :

25

M= Mn(II)-(IV), Fe(II)-(III), Cu(II), Co(II)-(III);

X= CH<sub>3</sub>CN, OH<sub>2</sub>, Cl<sup>-</sup>, Br<sup>-</sup>, OCN<sup>-</sup>, N<sub>3</sub><sup>-</sup>, SCN<sup>-</sup>, OH<sup>-</sup>, O<sup>2-</sup>, PO<sub>4</sub><sup>3-</sup>, C<sub>6</sub>H<sub>5</sub>BO<sub>2</sub><sup>2-</sup>, RCOO<sup>-</sup>;

Y= ClO<sub>4</sub><sup>-</sup>, BPh<sub>4</sub><sup>-</sup>, Br<sup>-</sup>, Cl<sup>-</sup>, [FeCl<sub>4</sub>]<sup>-</sup>, PF<sub>6</sub><sup>-</sup>, NO<sub>3</sub><sup>-</sup>;

a= 1, 2, 3, 4;

n= 0, 1, 2, 3, 4;


30

m= 0, 1, 2, 3, 4, 5, 6, 7, 8; und

k= 1, 2, 3, 4.

60. Verfahren nach einem der Ansprüche 19 bis 24, wobei L einen fünfzähligen Liganden der allgemeinen Formel (B) wiedergibt:

35



40

worin

45

jedes R<sup>1</sup>, R<sup>2</sup> unabhängig -R<sup>4</sup>-R<sup>5</sup> wiedergibt,

R<sup>3</sup> Wasserstoff, gegebenenfalls substituiertes Alkyl, Aryl oder Arylalkyl, oder -R<sup>4</sup>-R<sup>5</sup> wiedergibt,

jedes R<sup>4</sup> unabhängig eine Einfachbindung oder gegebenenfalls substituiertes Alkyl, Alkenyl, Oxyalkyl, Aminoalkyl, Alkylener, Carbonsäureester oder Carbonsäureamid wiedergibt, und


50

jedes R<sup>5</sup> unabhängig eine gegebenenfalls N-substituierte Aminoalkylgruppe oder eine gegebenenfalls substituierte Heteroarylgruppe, ausgewählt aus Pyridinyl, Pyrazinyl, Pyrazolyl, Pyrrolyl, Imidazolyl, Benzimidazolyl, Pyrimidinyl, Triazolyl und Thiazolyl, wiedergibt.

61. Verfahren nach Anspruch 60, mit der Maßgabe, dass R<sup>3</sup> keinen Wasserstoff wiedergibt.

55

62. Verfahren nach einem der Ansprüche 19 bis 24, wobei L einen fünfzähligen oder sechszähligen Liganden der allgemeinen Formel (C) wiedergibt:



worin

5 jedes R<sup>1</sup> unabhängig -R<sup>3</sup>-V wiedergibt, worin R<sup>3</sup> gegebenenfalls substituiertes Alkylen, Alkenylen, Oxyalkylen, Aminoalkylen oder Alkylenether wiedergibt und V eine gegebenenfalls substituierte Heteroarylgruppe, ausgewählt aus Pyridinyl, Pyrazinyl, Pyrazolyl, Pyrrolyl, Imidazolyl, Benzimidazolyl, Pyrimidinyl, Triazolyl und Thiazolyl, wiedergibt;

10 W eine gegebenenfalls substituierte Alkylenbrückengruppe wiedergibt, ausgewählt aus -CH<sub>2</sub>CH<sub>2</sub>-, -CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>-, -CH<sub>2</sub>-C<sub>6</sub>H<sub>4</sub>-CH<sub>2</sub>-, -CH<sub>2</sub>-C<sub>6</sub>H<sub>10</sub>-CH<sub>2</sub>- und -CH<sub>2</sub>-C<sub>10</sub>H<sub>6</sub>-CH<sub>2</sub>wiedergibt; und

15 R<sup>2</sup> eine Gruppe wiedergibt, ausgewählt aus R<sup>1</sup> und Alkyl-, Aryl- und Arylalkylgruppen, gegebenenfalls substituiert mit einem Substituenten, ausgewählt aus Hydroxy, Alkoxy, Phenoxy, Carboxylat, Carboxamid, Carbonsäureester, Sulfonat, Amin, Alkylamin und N<sup>+</sup>(R<sup>4</sup>)<sub>3</sub>, worin R<sup>4</sup> ausgewählt ist aus Wasserstoff, Alkanyl, Alkenyl, Arylalkanyl, Arylalkenyl, Oxyalkanyl, Oxyalkenyl, Aminoalkanyl, Aminoalkenyl, Alkanylether und Alkenylether.

63. Verfahren nach einem der Ansprüche 19 bis 24, wobei L einen makrocyclischen Liganden der Formel (E) wiedergibt:



25

worin

Z<sup>1</sup> und Z<sup>2</sup> unabhängig ausgewählt sind aus monocyclischen oder polycyclischen aromatischen Ringstrukturen, die gegebenenfalls ein oder mehrere Heteroatome enthalten, wobei jede aromatische Ringstruktur mit einem oder mehreren Substituenten substituiert ist;


30 Y<sup>1</sup> und Y<sup>2</sup> unabhängig ausgewählt sind aus Atomen C, N, O, Si, P und S;

A<sup>1</sup> und A<sup>2</sup> unabhängig ausgewählt sind aus Wasserstoff, Alkyl, Alkenyl und Cycloalkyl (wobei jedes Alkyl, Alkenyl und Cycloalkyl gegebenenfalls substituiert ist mit einer oder mehreren Gruppen, ausgewählt aus Hydroxy, Aryl, Heteroaryl, Sulfonat, Phosphat, Elektronen-abgebenden Gruppen und Elektronen-anziehenden Gruppen, und Gruppen der Formeln (G<sup>1</sup>)(G<sup>2</sup>)N-, G<sup>3</sup>OC(O)-, G<sup>3</sup>O- und G<sup>3</sup>C(O)-, worin jede von G<sup>1</sup>, G<sup>2</sup> und G<sup>3</sup> unabhängig ausgewählt ist aus Wasserstoff und Alkyl, und Elektronen-abgebenden Gruppen und/oder -anziehenden Gruppen (zusätzlich zu beliebigen unter den vorangehenden));

35

i und j ausgewählt sind aus 0, 1 und 2, um die Wertigkeit der Gruppen Y<sup>1</sup> und Y<sup>2</sup> zu vervollständigen; jeder von Q<sup>1</sup>-Q<sup>4</sup> unabhängig ausgewählt ist aus Gruppen der Formel

40



45

worin 10 > a + b + c + d > 2;

50 jedes Y<sup>3</sup> unabhängig ausgewählt ist aus -O-, -S-, -SO-, -SO<sub>2</sub>-, -(G<sup>1</sup>)(G<sup>2</sup>)N-, -(G<sup>1</sup>)N-, (worin G<sup>1</sup> und G<sup>2</sup> wie vorstehend hierin definiert sind), -C(O)-, Aryl, Heteroaryl, -P- und -P(O)-;

jedes von A<sup>3</sup>-A<sup>6</sup> unabhängig ausgewählt ist aus den vorstehend hierin für A<sup>1</sup> und A<sup>2</sup> definierten Gruppen; und worin beliebige zwei oder mehrere von A<sup>1</sup>-A<sup>6</sup> zusammen eine Brückengruppe bilden, vorausgesetzt, dass, wenn A<sup>1</sup> und A<sup>2</sup> verbunden sind, ohne auch gleichzeitig an beliebige von A<sup>3</sup>-A<sup>6</sup> zu binden, dann die Brückengruppe, die A<sup>1</sup> und A<sup>2</sup> verbindet, mindestens eine Carbonylgruppe enthalten muss.

55

**Revendications**

- 5 1. Procédé de traitement d'un textile par mise en contact du textile avec une substance organique qui forme un complexe avec un métal de transition, le traitement fournissant la substance organique sous une forme qui est substantiellement exempte d'agent azurant à base de peroxygène ou d'un système générant du peroxyde ou à base de peroxyde, moyennant quoi le complexe catalyse le blanchiment du textile par l'oxygène atmosphérique après le traitement, le textile traité étant séché et le blanchiment étant catalysé sur le textile sec.
- 10 2. Procédé selon la revendication 1, dans lequel le traitement comprend la mise en contact du textile avec la substance organique sous forme sèche.
3. Procédé selon la revendication 1, dans lequel le traitement comprend la mise en contact du textile avec un bain de teinture contenant la substance organique et, ensuite, le séchage.
- 15 4. Procédé selon la revendication 3, dans lequel le bain de teinture est un bain de teinture aqueux.
5. Procédé selon la revendication 4, dans lequel le bain de teinture est un fluide de traitement du tissu appliqué au pistolet.
- 20 6. Procédé selon la revendication 4, dans lequel le bain de teinture est un bain de teinture de lavage pour un nettoyage en blanchisserie.
7. Procédé selon la revendication 3, dans lequel le bain de teinture est un bain de teinture non aqueux.
- 25 8. Procédé selon la revendication 7, dans lequel le bain de teinture est un fluide pour nettoyage à sec.
9. Procédé selon la revendication 7, dans lequel la liqueur est un fluide aérosol appliqué au pistolet.
- 30 10. Procédé selon la revendication 3, dans lequel le milieu présente une valeur pH s'inscrivant dans une plage de pH 6 à 11.
11. Procédé selon la revendication 10, dans lequel la liqueur présente une valeur de pH s'inscrivant dans une plage allant de pH 8 à 10.
- 35 12. Procédé selon l'une quelconque des revendications 3 à 11, dans lequel le bain de teinture est substantiellement exempt d'un agent séquestrant des métaux de transition.
13. Procédé selon l'une quelconque des revendications 3 à 12, dans lequel le bain de teinture comprend en outre un agent tensioactif.
- 40 14. Procédé selon l'une quelconque des revendications 3 à 13, dans lequel le bain de teinture comprend en plus un adjuvant.
- 45 15. Procédé selon l'une quelconque des revendications 1 à 14, dans lequel la substance organique comprend un complexe préformé d'un ligand et d'un métal de transition.
16. Procédé selon l'une quelconque des revendications 3 à 14, dans lequel la substance organique comprend un ligand libre qui se complexe avec un métal de transition présent dans la liqueur.
- 50 17. Procédé selon l'une quelconque des revendications 1 à 14, dans lequel la substance organique comprend un ligand libre qui se complexe avec un métal de transition présent dans le textile.
18. Procédé selon l'une quelconque des revendications 1 à 14, dans lequel la substance organique comprend une composition d'un ligand libre ou d'un complexe de ligand-métal pouvant être substitué par un métal de transition et une source de métal de transition.
- 55 19. Procédé selon l'une quelconque des revendications 1 à 18, dans lequel la substance organique forme un complexe de formule générale (A1) :



dans laquelle :

5 M représente un métal choisi parmi Mn(II)-(III)-(IV)-(V), Cu(I)-(II)-(III), Fe(I)-(II)-(III)-(IV), Co(I)-(II)-(III), Ni(I)-(II)-(III), Cr(II)-(III)-(IV)-(V)-(VI)-(VII), Ti(II)-(III)-(IV), V(II)-(III)-(IV)-(V), Mo(II)-(III)-(IV)-(V)-(VI), W(IV)-(V)-(VI), Pd(II), Ru(II)-(III)-(IV)-(V) et Ag(I)-(II) ;

10 L représente un ligand, ou son analogue protoné ou déprotoné ;

X représente une espèce assurant la coordination choisie parmi tout anion mono, bi ou trichargé et toute molécule neutre en mesure de coordonner le métal d'une manière mono, bi ou tridentate ;

15 y représente tout contre-ion non coordonné ;

a représente un nombre entier de 1 à 10 ;

20 k représente un nombre entier de 1 à 10 ;

n vaut zéro ou représente un nombre de 1 à 10 ; et

m vaut zéro ou représente un nombre entier de 1 à 20.

25 **20.** Procédé selon la revendication 19, dans lequel dans la formule (A1) :

X représente une espèce assurant la coordination choisie parmi  $O^{2-}$ ,  $RBO_2^{2-}$ ,  $RCOO^-$ ,  $RCONR^-$ ,  $OH^-$ ,  $NO_3^-$ ,  $NO_2^-$ ,  $NO$ ,  $CO$ ,  $S^{2-}$ ,  $RS^-$ ,  $PO_3^{4-}$ , des anions dérivés de STP,  $PO_3OR^{3-}$ ,  $H_2O$ ,  $CO_3^{2-}$ ,  $HCO_3^-$ ,  $ROH$ ,  $NRR'R''$ ,  $RCN$ ,  $Cl^-$ ,  $Br^-$ ,  $OCN^-$ ,  $SCN^-$ ,  $CN^-$ ,  $N_3^-$ ,  $F^-$ ,  $I^-$ ,  $RO^-$ ,  $ClO_4^-$ ,  $SO_4^{2-}$ ,  $HSO_4^-$ ,  $SO_3^{2-}$  et  $RSO_3^-$  ; et

30 Y représente un contre-ion choisi parmi  $ClO_4^-$ ,  $BR_4^-$ ,  $[FeCl_4]^-$ ,  $PF_6^-$ ,  $RCOO^-$ ,  $NO_3^-$ ,  $NO_2^-$ ,  $RO^-$ ,  $N^+RR'R''R'''$ ,  $Cl^-$ ,  $Br^-$ ,  $F^-$ ,  $I^-$ ,  $RSO_3^-$ ,  $S_2O_6^{2-}$ ,  $OCN^-$ ,  $SCN^-$ ,  $Li^+$ ,  $Ba^{2+}$ ,  $Na^+$ ,  $Mg^{2+}$ ,  $K^+$ ,  $Ca^{2+}$ ,  $Cs^+$ ,  $PR_4^-$ ,  $RBO_2^{2-}$ ,  $SO_4^{2-}$ ,  $HSO_4^-$ ,  $SO_3^{2-}$ ,  $SbCl_6^-$ ,  $CuCl_4^{2-}$ ,  $CN^-$ ,  $PO_4^{3-}$ ,  $HPO_4^{2-}$ ,  $H_2PO_4^-$ , des anions dérivés de STP,  $CO_3^{2-}$ ,  $HCO_3^-$  et  $BF_4^-$ ,

35 dans laquelle

R, R', R'', R''' représentent indépendamment un groupe choisi parmi un atome d'hydrogène, des groupes dérivés de l'hydroxyle, -OR (où R = un groupe dérivé de l'alkyle, alcényle, cycloalkyle, hétérocycloalkyle, aryle, hétéroaryle ou carbonyle), -OAr, alkyle, alcényle, cycloalkyle, hétérocycloalkyle, aryle, hétéroaryle et carbonyle, chacun des groupes dérivés de R, Ar, alkyle, alcényle, cycloalkyle, hétérocycloalkyle, aryle, hétéroaryle et carbonyle étant éventuellement substitué par un groupe fonctionnels E ou plus, ou R6 conjointement avec R7 et indépendamment R8 conjointement avec R9 représente un atome d'oxygène ;

E est choisi parmi des groupes fonctionnels contenant un atome d'oxygène, de soufre, de phosphore, d'azote, de sélénium, des halogènes, et tout groupe donneur et/ou attracteur d'électrons.

45 **21.** Procédé selon la revendication 19 ou la revendication 20, dans lequel, dans la formule (A1) :

M représente un métal choisi parmi Mn(II)-(III)-(IV)-(V), Cu(I)-(II), Fe(II)-(III)-(IV) et Co(I)-(II)-(III) ;

50 X représente une espèce assurant la coordination choisie parmi  $O^{2-}$ ,  $RBO_2^{2-}$ ,  $RCOO^-$ ,  $OH^-$ ,  $NO_3^-$ ,  $NO_2^-$ ,  $NO$ ,  $CO$ ,  $CN^-$ ,  $S^{2-}$ ,  $RS^-$ ,  $PO_3^{4-}$ ,  $H_2O$ ,  $CO_3^{2-}$ ,  $HCO_3^-$ ,  $ROH$ ,  $NRR'R''$ ,  $Cl^-$ ,  $Br^-$ ,  $OCN^-$ ,  $SCN^-$ ,  $RCN$ ,  $N_3^-$ ,  $F^-$ ,  $I^-$ ,  $RO^-$ ,  $ClO_4^-$ ,  $SO_4^{2-}$ ,  $HSO_4^-$ ,  $SO_3^{2-}$  et  $RSO_3^-$  ;


55 Y représente un contre-ion choisi parmi  $ClO_4^-$ ,  $BR_4^-$ ,  $[FeCl_4]^-$ ,  $PF_6^-$ ,  $RCOO^-$ ,  $NO_3^-$ ,  $NO_2^-$ ,  $RO^-$ ,  $N^+RR'R''R'''$ ,  $Cl^-$ ,  $Br^-$ ,  $F^-$ ,  $I^-$ ,  $RSO_3^-$ ,  $S_2O_6^{2-}$ ,  $OCN^-$ ,  $SCN^-$ ,  $Li^+$ ,  $Ba^{2+}$ ,  $Na^+$ ,  $Mg^{2+}$ ,  $K^+$ ,  $Ca^{2+}$ ,  $PR_4^+$ ,  $SO_4^{2-}$ ,  $HSO_4^-$ ,  $SO_3^{2-}$  et  $BF_4^-$ ,

dans laquelle

R, R', R'', R''' représentent un atome d'hydrogène, un groupe alkyle éventuellement substitué ou un groupe


aryle éventuellement substitué ;  
 a représente un nombre entier de 1 à 4 ;  
 k représente un nombre entier de 1 à 10 ;  
 n vaut zéro ou représente un nombre entier de 1 à 4 ; et  
 m vaut zéro ou représente un nombre entier de 1 à 8.

22. Procédé selon l'une quelconque des revendications 19 à 21, dans lequel L représente un ligand répondant à la formule générale (BI) :



dans laquelle

g vaut zéro ou représente un nombre entier de 1 à 6 ;  
 r représente un nombre entier de 1 à 6 ;  
 s vaut zéro ou représente un nombre entier de 1 à 6 ;  
 Z1 et Z2 représentent indépendamment un hétéroatome ou un noyau hétérocyclique ou hétéroaromatique, Z1 et/ou Z2 étant éventuellement substitués par un groupe fonctionnels E ou plus tels que définis précédemment ;  
 Q1 et Q2 représentent indépendamment un groupe répondant à la formule :



dans laquelle

$10 > d + e + f > 1$  ;  $d = 0 - 9$  ;  $e = 0 - 9$  ;  $f = 0 - 9$  ;  
 chaque Y1 est choisi indépendamment parmi -O-, -S-, -SO-, -SO<sub>2</sub>-, -(G<sup>1</sup>)N-, -(G<sup>1</sup>)(G<sup>2</sup>)N- (dans lequel G<sup>1</sup> et G<sup>2</sup> sont tels que définis ci-dessous), -C(O)-, arylène, alkylène, hétéroarylène, -P- et -P(O)- ;  
 si  $s > 1$ , chaque groupe  $-\{Z1(R1) - (Q1)_r\}$  est défini de façon indépendante,  
 R1, R2, R6, R7, R8, R9 représentent indépendamment un groupe choisi parmi un atome d'hydrogène, des groupes dérivés de l'hydroxyle, -OR (dans lequel R = un groupe dérivé d'alkyle, alcényle, cycloalkyle, hétérocycloalkyle, aryle, hétéroaryle ou carbonyle), -OAr, alkyle, alcényle, cycloalkyle, hétérocycloalkyle, aryle, hétéroaryle et carbonyle, chacun des groupes dérivés de R, Ar, alkyle, alcényle, cycloalkyle, hétérocycloalkyle, aryle, hétéroaryle et carbonyle étant éventuellement substitué par un des groupes fonctionnels E ou plus, ou R6 conjointement avec R7 et indépendamment R8 conjointement avec R9 représentent un atome d'oxygène ;  
 E est choisi parmi des groupes fonctionnels contenant un atome l'oxygène, du soufre, du phosphore, de l'azote, du sélénium, des halogènes, et tout groupe donneur et/ou attracteurs d'électrons (de préférence, E est choisi parmi des dérivés d'hydroxy, de mono- ou polycarboxylate, d'un groupe aryle, d'un groupe hétéroaryle, d'un sulfonate, d'un thiol (-RSH), des thioéthers (-R-S-R'), des disulfures (-RSSR'), des dithiolènes, des mono- ou polyphosphonates, des mono- ou polyphosphates, des groupes donneurs d'électrons et des groupes attracteurs d'électrons, et des groupes de formule (G<sup>1</sup>) (G<sup>2</sup>)N-, (G<sup>1</sup>) (G<sup>2</sup>) (G<sup>3</sup>)N-, (G<sup>1</sup>) (G<sup>2</sup>)N-C(O)-, G<sup>3</sup>O- et G<sup>3</sup>C(O)-, dans lesquelles chacun de G<sup>1</sup>, G<sup>2</sup> et G<sup>3</sup> est indépendamment choisi parmi un atome d'hydrogène, un groupe alkyle, des groupes donneurs d'électrons et des groupes attracteurs d'électrons (en plus de ceux de la présente invention)) ;  
 ou un de R1 à R9 est un groupe de pontage lié à un autre groupement répondant à la même formule générale ;  
 T1 et T2 représentent indépendamment des groupes R4 et R5, dans lesquels R4 et R5 sont tels que définis pour R1 à R9, et si  $g = 0$  et  $s > 0$ , R1 conjointement avec R4, et/ou R2 conjointement avec R5, peuvent éventuellement représenter indépendamment =CH-R10, où R10 est tel que défini pour R1 à R9, ou  
 T1 et T2 peuvent conjointement (-T2-T1-) représenter une liaison covalente quand  $s > 1$  et  $g > 0$  ;  
 si Z1 et/ou Z2 représentent N et T1 et T2 représentent ensemble une liaison simple et R1 et/ou R2 sont absents, Q1 et/ou Q2 peuvent représenter indépendamment un groupe répondant à la formule : =CH-[-Y1-]<sub>c</sub>-CH= ,

EP 1 109 965 B1

éventuellement, deux quelconques de R1, R2, R6, R7, R8, R9 indépendamment, ou plus sont liés ensemble par une liaison covalente ;

si Z1 et/ou Z2 représentent O, alors R1 et/ou R2 n'existent pas ;

si Z1 et/ou Z2 représentent S, N, P, B ou Si alors R1 et/ou R2 peuvent être absents ;


si Z1 et/ou Z2 représentent un hétéroatome substitué par un groupe fonctionnel E alors R1 et/ou R2 et/ou R4 et/ou R5 peuvent être absents.

23. Procédé selon la revendication 22, dans lequel Z1 et Z2 représentent indépendamment un hétéroatome éventuellement substitué choisi parmi N, P, O, S, B et Si ou un noyau hétérocyclique optionnellement substitué ou un noyau hétéroaromatique optionnellement substitué choisi parmi la pyridine, des pyrimidines, la pyrazine, la pyrimidine, le pyrazole, le pyrrole, l'imidazole, le benzimidazole, la quinoléine, l'isoquinoléine, le carbazole, l'indole, l'isindole, le furane, le thiophène, l'oxazole et le thiazole.

24. Procédé selon la revendication 22 ou 23, dans lequel R1 à R9 sont choisis indépendamment parmi un atome H, hydroxyalkyle en C<sub>0</sub>-C<sub>20</sub>, halogéno alkyle en C<sub>0</sub>-C<sub>20</sub>, nitroso, formylalkyle en C<sub>0</sub>-C<sub>20</sub>, carboxylalkyle en C<sub>0</sub>-C<sub>20</sub>, et des esters et des sels de ceux-ci, carbamoylalkyle en C<sub>0</sub>-C<sub>20</sub>, sulfoalkyle en C<sub>0</sub>-C<sub>20</sub>, et des esters et des sels de ceux-ci, sulphamoylalkyle en C<sub>0</sub>-C<sub>20</sub>, aminoalkyle en C<sub>0</sub>-C<sub>20</sub>, arylalkyle en C<sub>0</sub>-C<sub>20</sub>, hétéroarylalkyle en C<sub>0</sub>-C<sub>20</sub>, alkyle en C<sub>0</sub>-C<sub>20</sub>, alcoxyalkyle en C<sub>0</sub>-C<sub>8</sub>, carbonylalkoxy en C<sub>0</sub>-C<sub>6</sub>, et arylalkyle en C<sub>0</sub>-C<sub>6</sub> et le amide d'alkyle en C<sub>0</sub>-C<sub>20</sub>,

ou un de R1 et R9 représente un groupe de pontage -C<sub>n</sub>(R11) (R12)-(D)<sub>p</sub>-C<sub>m</sub>(R11) (R12)- lié à un autre groupement de la même formule générale, dans laquelle p vaut zéro ou un, D est choisi parmi un hétéroatome ou un groupe contenant un hétéroatome, ou est une partie d'un noyau aromatique ou homonucléaire saturé ou hétéronucléaire, n' représente un nombre entier de 1 à 4, m' représente un nombre entier de 1 à 4, à condition que n' + m' <= 4, R11 et R12 sont chacun choisis de préférence indépendamment parmi -H, NR13 et OR14, un groupe alkyle, un groupe aryle, éventuellement substitués, et R13 et R14 sont chacun indépendamment choisis parmi un atome d'hydrogène, un groupe alkyle, un groupe aryle, les deux étant éventuellement substitués.

25. Procédé selon l'une quelconque des revendications 22 à 24, dans lequel T1 et T2 conjointement forment une liaison simple et s > 1, selon la formule générale (BII) :




dans laquelle Z3 représente indépendamment un groupe tel que défini pour Z1 ou Z2 ; R3 représente indépendamment un groupe tel que défini pour R1 à R9 ; Q3 représente indépendamment un groupe tel que défini pour Q1, Q2 ; h vaut zéro ou représente un nombre entier de 1 à 6 ; et s' = s - 1.


26. Procédé selon la revendication 25, dans lequel dans la formule générale (BII), s' = 1, 2 ou 3 ; r = g = h = 1 ; d = 2 ou 3 ; e = f = 0 ; R6 = R7 = H.

27. Procédé selon la revendication 26, dans lequel le ligand répond à une formule générale choisie parmi :

5




10



15


20



25

28. Procédé selon la revendication 27, dans lequel le ligand répond à une formule générale choisie parmi :

30



35

40

29. Procédé selon la revendication 28, dans lequel R1, R2, R3 et R4 sont indépendamment choisis parmi un atome H, un groupe alkyle, un groupe hétéroaryle, ou représente un groupe de pontage lié à un autre groupement répondant à la même formule générale, un groupe de pontage étant un alkylène ou un hydroxyalkylène ou un pont contenant un groupe hétéroaryle.

45

30. Procédé selon la revendication 29, dans lequel R1, R2, R3 et R4 sont indépendamment choisis parmi un atome H, un groupe méthyle, un groupe éthyle, un groupe isopropyle, un groupe hétéroaryle contenant un azote, ou un groupe de pontage lié à un autre groupement répondant à la même formule générale, le groupe de pontage étant l'alkylène ou l'hydroxyalkylène.

50

31. Procédé selon une quelconque des revendications 26 à 30, dans lequel dans le complexe  $[M_a L_x X_a] Y_m$  :

M = Mn(II)-(IV), Cu(I)-(III), Fe(II)-(III), Co(II)-(III) ;

X = CH<sub>3</sub>CN, OH<sub>2</sub>, Cl<sup>-</sup>, Br<sup>-</sup>, OCN<sup>-</sup>, N<sub>3</sub><sup>-</sup>, SCN<sup>-</sup>, OH<sup>-</sup>, O<sup>2-</sup>, PO<sub>6</sub><sup>3-</sup>, C<sub>6</sub>H<sub>5</sub>BO<sub>2</sub><sup>2-</sup>, RCOO<sup>-</sup> ;

55

Y = ClO<sub>4</sub><sup>-</sup>, BPh<sub>4</sub><sup>-</sup>, Br<sup>-</sup>, Cl<sup>-</sup>, [FeCl<sub>4</sub>]<sup>-</sup>, PF<sub>6</sub><sup>-</sup>, NO<sub>3</sub><sup>-</sup>

a = 1, 2, 3, 4 ;

EP 1 109 965 B1


$n = 0, 1, 2, 3, 4, 5, 6, 7, 8, 9$  ;

$m = 1, 2, 3, 4$  ; et

5  $k = 1, 2, 4$ .

32. Procédé selon la revendication 25, dans lequel dans la formule générale (BII),  $s' = 2$  ;  $r = g = h = 1$  ;  $d = f = 0$ ,  $e = 1$  ; et chaque  $Y_1$  est indépendamment un alkylène ou un hétéroarylène.

10 33. Procédé selon la revendication 32, dans lequel le ligand répond à la formule générale :



dans laquelle

$A_1, A_2, A_3, A_4$  sont indépendamment choisis parmi des groupes hétéroarylène ou alkylène en  $C_1$  à  $9$  ; et  $N_1$  et  $N_2$  représentent indépendamment un hétéroatome ou un groupe hétéroarylène.

25 34. Procédé selon la revendication 33, dans lequel


$N_1$  représente un azote aliphatique ;

$N_2$  représente un groupe hétéroarylène ;

$R_1, R_2, R_3, R_4$  représentent chacun indépendamment un atome H, un groupe alkyle, un groupe aryle ou un groupe hétéroaryle ; et

30  $A_1, A_2, A_3, A_4$  représentent chacun  $-CH_2-$ .

35 35. Procédé selon la revendication 34, dans lequel le ligand répond à la formule générale :



dans laquelle  $R_1, R_2$  représentent chacun indépendamment un atome H, un groupe alkyle, un groupe aryle ou un groupe hétéroaryle.

36. Procédé selon l'une quelconque des revendications 32 à 35, dans lequel dans le complexe  $[M_a L_k X_n] Y_m$  :

50  $M = Fe(II)-(III), Mn(II)-(IV), Cu(II), Co(II)-(III)$  ;

$X = CH_3CN, OH_2, Cl^-, Br^-, OCN^-, N_3^-, SCN^-, OH^-, O^{2-}, PO_4^{3-}, C_6H_5BO_2^{2-}, RCOO^-$  ;

55  $Y = ClO_4^-, BPh_4^-, Br^-, Cl^-, [FeCl_4]^-, PF_6^-, NO_3^-$  ;

$a = 1, 2, 3, 4$  ;

$n = 0, 1, 2, 3, 4, 5, 6, 7, 8, 9$  ;




m = 1, 2, 3, 4 ; et

k = 1, 2, 4.

- 5 37. Procédé selon la revendication 25, dans lequel dans la formule générale (BII), s' = 2 et r = g = h = 1, selon la formule générale :

10




15

- 20 38. Procédé selon la revendication 37, dans lequel Z1 = Z2 = Z3 = Z4 = un noyau hétéroaromatique ; e = f = 0 ; d = 1 ; et R7 est absent.

39. Procédé selon la revendication 37, dans lequel Z1 à Z4 représentent chacun N ; R1 à R4 sont absents, Q1 et Q3 représentent tout deux =CH-[-Y1-]<sub>e</sub>-CH= ; et Q2 et Q4 représentent tout deux -CH<sub>2</sub>-[-Y1-]<sub>n</sub>-CH<sub>2</sub>-.

- 25 40. Procédé selon la revendication 39, dans lequel le ligand répond à la formule générale :

30



35

40

dans laquelle A représente un alkylène éventuellement substitué éventuellement interrompu par un hétéroatome ; et n vaut zéro ou représente un nombre entier de 1 à 5.

- 45 41. Procédé selon la revendication 40, dans lequel R1 à R6 représentent un atome d'hydrogène, n = 1 et A = -CH<sub>2</sub>, -CHOH-, -CH<sub>2</sub>N(R)CH<sub>2</sub>- ou -CH<sub>2</sub>CH<sub>2</sub>N(R)CH<sub>2</sub>CH<sub>2</sub>- où R représente un atome d'hydrogène ou un groupe alkyle.

42. Procédé selon la revendication 41, dans lequel A = -CH<sub>2</sub>-, -CHOH- ou -CH<sub>2</sub>CH<sub>2</sub>NHCH<sub>2</sub>CH<sub>2</sub>-.

- 50 43. Procédé selon l'une quelconque des revendications 37 à 42, dans lequel dans le complexe [M<sub>a</sub>L<sub>k</sub>X<sub>n</sub>]Y<sub>m</sub> :

M- Mn(II) - (IV), Co(II) - (III), Fe(II) - (III) ;

X- CH<sub>3</sub>CN, OH<sub>2</sub>, Cl<sup>-</sup>, Br<sup>-</sup>, OCN<sup>-</sup>, N<sub>3</sub><sup>-</sup>, SCN<sup>-</sup>, OH<sup>-</sup>, O<sup>2-</sup>, PO<sub>4</sub><sup>3-</sup>, C<sub>6</sub>H<sub>5</sub>BO<sub>2</sub><sup>2-</sup>, RCOO<sup>-</sup> ;

55

Y = ClO<sub>4</sub><sup>-</sup>, BPh<sub>4</sub><sup>-</sup>, Br<sup>-</sup>, Cl<sup>-</sup>, [FeCl<sub>4</sub>]<sup>-</sup>, PF<sub>6</sub><sup>-</sup>, NO<sub>3</sub><sup>-</sup> ;

a = 1, 2, 3, 4 ;


$n = 0, 1, 2, 3, 4, 5, 6, 7, 8, 9$  ;

$m = 1, 2, 3, 4$  ; et

5  $k = 1, 2, 4$ .

44. Procédé selon l'une quelconque des revendications 22 à 24, dans lequel T1 et T2 représentent indépendamment des groupes R4, R5 tels que définis pour R1 à R9, selon la formule générale (BIII) :

10




15

45. Procédé selon la revendication 44, dans lequel dans la formule générale (BIII),  $s = 1$  ;  $r = 1$  ;  $g = 0$  ;  $d = f = 1$  ;  $e = 1-4$  ;  $Y1 = -\text{CH}_2-$  ; et R1 conjointement avec R4, et/ou R2 conjointement avec R5, représentent indépendamment  $=\text{CH-R10}$ , dans laquelle R10 est tel que défini pour R1 à R9.

20 46. Procédé selon la revendication 45, dans lequel R2 conjointement avec R5 représente  $=\text{CH-R10}$ .

47. Procédé selon la revendication 45 ou 46, dans lequel le ligand est choisi parmi :


25



30

48. Procédé selon la revendication 47, dans lequel le ligand est choisi parmi :

35



40

dans laquelle R1 et R2 sont choisis parmi des phénols éventuellement substitués, des groupes hétéroarylalkyles en  $\text{C}_0\text{-C}_{20}$ , R3 et R4 sont choisis parmi un atome H, un groupe alkyle, un groupe aryle, des phénols éventuellement substitués, des groupes hétéroarylalkyles en  $\text{C}_0\text{-C}_{20}$ , un groupe alkylaryle, un groupe aminoalkyle, un groupe alcoxy.

45

49. Procédé selon la revendication 48 dans lequel R1 et R2 sont choisis parmi des phénols éventuellement substitués, des groupes hétéroarylalkyles en  $\text{C}_0\text{-C}_2$ , R3 et R4 sont choisis parmi un atome H, un groupe alkyle, un groupe aryle, des phénols éventuellement substitués, des hétéroarylalkyles en  $\text{C}_0\text{-C}_2$ -d'azote.

50 50. Procédé selon l'une quelconque des revendications 45 à 49 dans lequel dans le complexe  $[\text{M}_a\text{L}_k\text{X}_n]\text{Y}_m$  :

$\text{M} = \text{Mn(II)} - (\text{IV}), \text{Co(II)} - (\text{III}) ; \text{Fe(II)} - (\text{III})$  ;

$\text{X} = \text{CH}_3\text{CN}, \text{OH}_2, \text{Cl}^-, \text{Br}^-, \text{OCN}^-, \text{N}_3^-, \text{SCN}^-, \text{OH}^-, \text{O}^{2-}, \text{PO}_4^{3-}, \text{C}_6\text{H}_5\text{BO}_2^{2-}, \text{RCOO}^-$  ;

55

$\text{Y} = \text{ClO}_4^-, \text{BPh}_4^-, \text{Br}^-, \text{Cl}^-, [\text{FeCl}_4]^- , \text{PF}_6^-, \text{NO}_3^-$  ;

$a = 1, 2, 3, 4$ ,

EP 1 109 965 B1

$n = 0, 1, 2, 3, 4, 5, 6, 7, 8, 9$  ;

$m = 1, 2, 3, 4$  ; et

5  $k = 1, 2, 4$ .

51. Procédé selon la revendication 44, dans lequel dans la formule générale (BIII),  $s = 1$  ;  $r = 1$  ;  $g = 0$  ;  $d = f = 1$ ,  $e = 1-4$  ;  $Y1 = -C(R')(R'')$ , dans laquelle  $R'$  et  $R''$  sont indépendamment tels que définis pour R1-R9.

10 52. Procédé selon la revendication 51, dans lequel le ligand répond à la formule générale :



20 53. Procédé selon la revendication 52, dans lequel R1, R2, R3, R4, R5 représentent un atome H et/ou un groupe alkyle en  $C_0-C_{20}$ ,  $n = 0$  ou 1, R6 représente un atome H, un groupe alkyle, -OH ou -SH, et R7, R8, R9, R10 sont chacun choisis indépendamment parmi un atome H, un groupe alkyle en  $C_0-C_{20}$ , un groupe hétéroarylalkyle en  $C_0-C_{20}$ , un groupe alcoxyalkyle en  $C_0-C_{20}$  et un groupe aminoalkyle en  $C_0-C_{20}$ .

25 54. Procédé selon l'une quelconque des revendications 51 à 53 dans lequel le complexe  $[M_a L_k X_n] Y_m$  :

$M = Mn(II) - (IV), Fe(II) - (III), Cu(II), Co(II) - (III)$  ;

30  $X = CH_3CN, OH_2, Cl^-, Br^-, OCN^-, N_3^-, SCN^-, OH^-, O^{2-}, PO_4^{3-}, C_6H_5BO_2^{2-}, RCOO^-$  ;

$Y = ClO_4^-, BPh_4^-, Br^-, Cl^-, [FeCl_4]^-, PF_6^-, NO_3^-$  ;

35  $a = 1, 2, 3, 4$  ;

$n = 0, 1, 2, 3, 4$  ;

$m = 0, 1, 2, 3, 4, 5, 6, 7, 8$  ; et

40  $k = 1, 2, 3, 4$ .


55. Procédé selon la revendication 44, dans lequel dans la formule générale (BIII),  $s = 0$  ;  $g = 1$  ;  $d = e = 0$  ;  $f = 1-4$ .

45 56. Procédé selon la revendication 55, dans lequel le ligand répond à la formule générale :



55 57. Procédé selon la revendication 56, avec la condition qu'aucun de R1 et R3 ne représente un atome d'hydrogène.

58. Procédé selon la revendication 56 ou la revendication 57, dans lequel le ligand répond à la formule générale :



dans laquelle R1, R2, R3 sont tels que définis pour R2, R4, R5.

59. Procédé selon l'une quelconque des revendications 55 à 58, dans lequel le complexe  $[M_a L_k X_n] Y_m$  :

M = Mn(II) - (IV), Fe(II) - (III), Cu(II), Co(II) - (III) ;

X = CH<sub>3</sub>CN, OH<sub>2</sub>, Cl<sup>-</sup>, Br<sup>-</sup>, OCN<sup>-</sup>, N<sub>3</sub><sup>-</sup>, SCN<sup>-</sup>, OH<sup>-</sup>, O<sup>2-</sup>, PO<sub>4</sub><sup>3-</sup>, C<sub>6</sub>H<sub>5</sub>BO<sub>2</sub><sup>2-</sup>, RCOO<sup>-</sup> ;

Y = ClO<sub>4</sub><sup>-</sup>, BPh<sub>4</sub><sup>-</sup>, Br<sup>-</sup>, Cl<sup>-</sup>, [FeCl<sub>4</sub>]<sup>-</sup>, PF<sub>6</sub><sup>-</sup>, NO<sub>3</sub><sup>-</sup> ;


a = 1, 2, 3, 4 ;

n = 0, 1, 2, 3, 4 ;

m = 0, 1, 2, 3, 4, 5, 6, 7, 8 ; et

k = 1, 2, 3, 4.

60. Procédé selon l'une quelconque des revendications 19 à 24, dans lequel L représente un ligand pentadentate répondant à la formule générale (B) :



dans laquelle :

chaque R<sup>1</sup>, R<sup>2</sup> représente indépendamment -R<sup>4</sup>-R<sup>5</sup>,


R<sup>3</sup> représente un atome d'hydrogène, un alkyle éventuellement substitué, un groupe aryle ou un groupe arylalkyle, ou -R<sup>4</sup>-R<sup>5</sup>,

chaque R<sup>4</sup> représente indépendamment une liaison simple ou un alkylène, alcénylène, un oxyalkylène, un aminoalkylène, un éther d'alkylène, un ester carboxylique ou un amide carboxylique éventuellement substitué et

chaque R<sup>5</sup> représente indépendamment un groupe aminoalkyle éventuellement N-substitué ou un groupe hétéroaryle éventuellement substitué choisi parmi le groupe pyridinyle, le groupe pyrazinyle, le groupe pyrazolyle, le groupe pyrrolyle, le groupe imidazolyle, le groupe benzimidazolyle, le groupe pyrimidinyle, le groupe triazolyle et le groupe thiazolyle.

61. Procédé selon la revendication 60, à condition que R<sup>3</sup> ne représente pas un atome d'hydrogène.

62. Procédé selon l'une quelconque des revendications 19 à 24, dans lequel L représente un ligand pentadentate ou hexadentate répondant à la formule générale (C) :



dans laquelle

5 chaque R<sup>1</sup> représente indépendamment -R<sup>3</sup>-V, dans laquelle R<sup>3</sup> représente un alkylène, un alcénylène, un oxyalkylène, un aminoalkylène ou un éther d'alkylène éventuellement substitué, et V représente un groupe hétéroaryle éventuellement substitué choisi parmi le groupe pyridinyle, le groupe pyrazinyle, le groupe pyrazolyle, le groupe pyrrolyle, le groupe imidazolyle, le groupe benzimidazolyle, le groupe pyrimidinyle, le groupe triazolyle et le groupe thiazolyle ;

10 W représente un groupe de pontage d'alkylène éventuellement substitué choisis parmi -CH<sub>2</sub>CH<sub>2</sub>-, -CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>-, -CH<sub>2</sub>-CH<sub>2</sub>-C<sub>6</sub>H<sub>4</sub>-CH<sub>2</sub>-, -CH<sub>2</sub>-C<sub>6</sub>H<sub>10</sub>-CH<sub>2</sub>-, et -CH<sub>2</sub>-C<sub>10</sub>H<sub>6</sub>-CH<sub>2</sub>- ; et

15 R<sup>2</sup> représente un groupe choisi parmi R<sup>1</sup>, et des groupes alkyles, aryles et arylalkyles éventuellement substitués par un substituant choisi parmi un hydroxy, un alcoxy, un phénoxy, un carboxylate, un carboxamide, un ester carboxylique, un sulfonate, une amine, une alkylamine, et N<sup>4</sup>(R<sup>4</sup>)<sub>3</sub>, où R<sup>4</sup> est choisi parmi un atome d'hydrogène, un groupe alcanyle, un groupe alcényle, un groupe arylalcanyle, un groupe arylalcényle, un groupe oxyalcanyle, un groupe oxyalcényle, un groupe aminoalcanyle, un groupe aminoalcényle, un éther d'alcanyle et un éther d'alcényle.

20 63. Procédé selon l'une quelconque des revendications 19 à 24, dans lequel L représente un ligand macrocyclique répondant à la formule (E) :



dans laquelle

30 Z<sup>1</sup> et Z<sup>2</sup> sont choisis indépendamment parmi des structures de noyau aromatique monocyclique ou polycyclique contenant éventuellement un hétéroatome ou plus, chaque structure de noyau aromatique étant substituée par un substituant ou plus ;

Y<sup>1</sup> et Y<sup>2</sup> sont indépendamment choisis parmi des atomes C, N, O, Si, P et S ;

35 A<sup>1</sup> et A<sup>2</sup> sont choisis indépendamment parmi un atome d'hydrogène, un groupe alkyle, un groupe alcényle et un groupe cycloalkyle (chacun des groupes alkyle, alcényle et cycloalkyle) étant éventuellement substitué par un groupe ou plus choisis parmi un groupe hydroxy, un groupe aryle, un groupe hétéroaryle, un sulfonate, un phosphate, des groupes donneurs d'électrons, des groupes attracteurs d'électrons et des groupes répondant aux formules (G<sup>1</sup>) (G<sup>2</sup>)N-, G<sup>3</sup>OC(O)-, G<sup>3</sup>O-et G<sup>3</sup>C(O)-, dans lesquelles chacun de G<sup>1</sup>, G<sup>2</sup> et G<sup>3</sup> est indépendamment choisi parmi l'atome d'hydrogène ou un groupe alkyle, des groupes donneurs d'électrons et/ou attracteurs (en plus de ceux de la présente invention) ;

40 i et j sont choisis parmi 0, 1 et 2 pour compléter la valence des groupes Y<sup>1</sup> et Y<sup>2</sup> ;

chacun de Q<sup>1</sup> à Q<sup>4</sup> est indépendamment choisi parmi les groupes répondant à la formule



dans laquelle 10 > a + b + c + d > 2 ;

55 chaque Y<sup>3</sup> est indépendamment choisi parmi -O-, -S-, -SO-, -SO<sub>2</sub>-, -(G<sup>1</sup>) (G<sup>2</sup>)N-, -(G<sup>1</sup>)N- (dans lesquelles G<sup>1</sup> et G<sup>2</sup> sont tels que définis précédemment), -C(O)-, un groupe aryle, un groupe hétéroaryle, -P- et -P(O)-,

chacun de A<sup>3</sup> à A<sup>6</sup> est choisi de façon indépendante parmi les groupes définis précédemment pour A<sup>1</sup> et A<sup>2</sup> ; et

où deux quelconques ou plus de A<sup>1</sup> à A<sup>6</sup> forment ensemble un groupe de pontage, à la condition que le groupe de pontage A<sup>1</sup> et A<sup>2</sup> contienne obligatoirement au moins un groupe carbonyle, si A<sup>1</sup> et A<sup>2</sup> sont liés sans

**EP 1 109 965 B1**

liaison simultanée également à l'un quelconque de  $A^3$  à  $A^6$ .

5

10

15

20

25

30

35

40

45

50

55