



Europäisches Patentamt  
European Patent Office  
Office européen des brevets



(11) **EP 1 358 309 B1**

(12) **EUROPEAN PATENT SPECIFICATION**

(45) Date of publication and mention  
of the grant of the patent:  
**11.10.2006 Bulletin 2006/41**

(51) Int Cl.:  
**C11D 3/39** <sup>(2006.01)</sup> **C07D 233/54** <sup>(2006.01)</sup>  
**C07D 235/14** <sup>(2006.01)</sup>

(21) Application number: **02722032.6**

(86) International application number:  
**PCT/EP2002/000688**

(22) Date of filing: **22.01.2002**

(87) International publication number:  
**WO 2002/077145 (03.10.2002 Gazette 2002/40)**

(54) **COMPOSITION AND METHOD FOR BLEACHING A SUBSTRATE**

MITTEL UND VERFAHREN ZUM BLEICHEN EINES SUBSTRATS

COMPOSITION ET PROCEDE DE BLANCHIMENT D'UN SUBSTRAT

(84) Designated Contracting States:  
**AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU  
MC NL PT SE TR**

(30) Priority: **05.02.2001 GB 0102826**

(43) Date of publication of application:  
**05.11.2003 Bulletin 2003/45**

(73) Proprietors:  
• **UNILEVER PLC**  
**London EC4P 4BQ (GB)**  
Designated Contracting States:  
**CY GB IE**  
• **UNILEVER N.V.**  
**3013 AL Rotterdam (NL)**  
Designated Contracting States:  
**AT BE CH DE DK ES FI FR GR IT LI LU MC NL PT  
SE TR**

(72) Inventors:  
• **CARINA, Riccardo F.**  
**CH-1208 Geneve (CH)**

• **DELROISSE, Michel G. J.,**  
**Unilever R&D Vlaardingen**  
**3133 AT Vlaardingen (NL)**  
• **HAGE, Ronald,**  
**Unilever R&D Vlaardingen**  
**3133 AT Vlaardingen (NL)**  
• **TETARD, David,**  
**Unilever Research Port Sunlight**  
**Wirral,**  
**Merseyside CH63 3JW (GB)**

(56) References cited:  
**EP-A- 1 001 009** **WO-A-00/12808**  
**WO-A-00/60043**

• **GOMEZ-ROMERO P ET AL: "Dissimetry effects  
in u-oxo diiron (III) species: structures and  
spectroscopic properties of [N5FeOX3]+ (X=Cl,  
Br) and Implications for oxo-bridged dinuclear  
iron proteins" JOURNAL OF THE AMERICAN  
CHEMICAL SOCIETY, vol. 111, no. 25, 1989,  
pages 9039-9047, XP002207358 American  
Chemical Society, USA**

**EP 1 358 309 B1**

Note: Within nine months from the publication of the mention of the grant of the European patent, any person may give notice to the European Patent Office of opposition to the European patent granted. Notice of opposition shall be filed in a written reasoned statement. It shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

**Description**FIELD OF INVENTION

5 **[0001]** This invention relates to compositions and methods for catalytically bleaching substrates with air or a peroxy species using a defined class of ligand or complex as catalyst.

BACKGROUND OF INVENTION

10 **[0002]** The use of bleaching catalysts for stain removal has been developed over recent years. The recent discovery that some catalysts are capable of bleaching effectively in the absence of an added peroxy source has recently become the focus of some interest, for example: W09965905; WO0012667; WO0012808; WO0029537, and, W00060045. The compounds found in the aforementioned application are, in many instances, also useful as bleach catalysts for use with peroxy species.

15 **[0003]** EP 1.001.009 A discloses a bleaching oxidation catalyst including a pentadentate nitrogen-containing ligand for bleaching and cleaning fabrics.

**[0004]** The search for new classes of compounds that are suitable as air bleaching catalyst is ongoing. In addition, the selection within known groups for classes of compounds that are surprisingly effective as air bleaching catalysts and peroxy bleaching catalysts is sought.

20


SUMMARY OF INVENTION

**[0005]** We have found that a selected class of ligand or complex is surprisingly effective in catalysing the bleaching of substrates using either air or a peroxy species.

25 **[0006]** We have selected a group of ligands that are surprisingly effective as bleach catalysts. The ligand group selected has at least two imidazol groups as described herein.

**[0007]** The present invention provides a laundry bleach composition, capable of bleaching in an air bleaching mode or peroxy bleaching mode, comprising a ligand of general formula (L):

30



wherein:

X is selected from  $-CH_2CH_2-$ ,  $-CH_2CH_2CH_2-$ ,  $-CH_2C(OH)HCH_2-$ ; and,


35

$R^1$  independently represents a group selected from: alkyl, cycloalkyl, heterocycloalkyl, heteroaryl, aryl and arylalkyl groups optionally substituted with a substituent selected from hydroxy, alkoxy, phenoxy, carboxylate, carboxamide, carboxylic ester, sulphonate, amine, alkylamine and  $N^+(R^4)_3$ , wherein  $R^4$  is selected from hydrogen, alkanyl, alkenyl, arylalkanyl, arylalkenyl, oxyalkanyl, oxyalkenyl, aminoalkanyl, aminoalkenyl, alkanyl ether, alkenyl ether, and  $-CY_2-R^2$ , in which Y is independently selected from H, CH<sub>3</sub>, C<sub>2</sub>H<sub>5</sub>, C<sub>3</sub>H<sub>7</sub> and  $R^2$  is independently selected from:

40

an optionally substituted

45



50

wherein  $R^3$  is independently selected from: H, alkyl, and benzyl, and wherein at least two of  $R^1$  are  $-CY_2-R^2$ . The optionally substituted imidazole is not so substituted that it forms a benzimidazol group per se.

**[0008]** It is essential that the ligand (L) has at least two imidazole substituents as defined above, preferably at least three imidazole substituents. The nature of the remaining  $R^1$  substituent(s) has not been found to be significant to activity of the selected ligand.

55

**[0009]** The present invention also provides a method of bleaching a substrate comprising applying to the substrate, in an aqueous medium, a bleaching composition as defined herein.

**[0010]** The present invention also provides the ligands defined in claim 15.

[0011] The present invention also extends to use of a ligand or complex as defined below in the manufacture of a bleaching composition, the bleaching composition containing effective amounts of a peroxygen bleach or a peroxy-based or peroxy-generating bleach system. An effective amount of a peroxygen bleach may be provided by a composition containing at least 1%, preferably at least 5%, of a peroxy species. In the present invention, it is preferred that the composition contains the range of about 1-35% by weight, preferably from 5-25% of a peroxy species.

[0012] The present invention also extends to a commercial package comprising a bleaching composition according to the present invention together with instructions for its use.

## DESCRIPTION OF THE INVENTION

### AIR BLEACHING MODE

[0013] In an air bleaching mode the composition of the present invention uses air to bleach a substrate. This is distinct from using pure oxygen or an enriched oxygen source. Air is different to molecular oxygen. The provision of a commercially available air bleaching product negates the requirement of an additional component, namely a peroxy source. The removal of a peroxy species, an expensive component, from a bleaching composition results in a reduction in manufacturing costs of the bleaching composition. Of significant importance is that an increased retention of textile strength and less dye damage is found when a bleaching composition without an added peroxy species is used to clean fabrics. In addition, the provision of a commercially available air bleaching composition that may function without the requirement of saturated oxygen solutions and/or pressure vessels in an aqueous environment is important. Oxygen is relatively insoluble in water when compared to organic solvents. Nitrogen makes up approximately 80% of the volume of air whilst molecular oxygen makes up only approximately 20% of the volume of air.

[0014] In the present invention at least 10 %, preferably at least 50 % and optimally at least 90 % of any bleaching of the substrate is effected by oxygen sourced from the air.

[0015] In any composition containing organic matter it is difficult to avoid the presence of hydroperoxides which are readily formed from the oxygen in the air. In this regard, the air bleaching composition of the present invention has less than 1%, preferably less than 0.1%, most preferably less than 0.01%, of a peroxy species present.

### PEROXYL BLEACHING MODE

#### THE PEROXY SPECIES OR PRECURSOR THEREOF


[0016] In a peroxy bleaching mode the composition of the present invention uses a peroxy species to bleach a substrate. The peroxy bleaching species may be a compound which is capable of yielding hydrogen peroxide in aqueous solution. Hydrogen peroxide sources are well known in the art. They include the alkali metal peroxides, organic peroxides such as urea peroxide, and inorganic persalts, such as the alkali metal perborates, percarbonates, perphosphates persulfates and persulfates. Mixtures of two or more such compounds may also be suitable.

[0017] Particularly preferred are sodium perborate tetrahydrate and, especially, sodium perborate monohydrate. Sodium perborate monohydrate is preferred because of its high active oxygen content. Sodium percarbonate may also be preferred for environmental reasons. The amount thereof in the composition of the invention usually will be within the range of about 1-35% by weight, preferably from 5-25% by weight. One skilled in the art will appreciate that these amounts may be reduced in the presence of a bleach precursor e.g., N,N,N',N'-tetraacetyl ethylene diamine (TAED).

[0018] Another suitable hydrogen peroxide generating system is a combination of a C1-C4 alkanol oxidase and a C1-C4 alkanol, especially a combination of methanol oxidase (MOX) and ethanol. Such combinations are disclosed in International Application WO 9 507 972 A (PCT/EP 94/03003) (Unilever).

[0019] Alkylhydroxy peroxides are another class of peroxy bleaching compounds. Examples of these materials include cumene hydroperoxide and t-butyl hydroperoxide.


[0020] Organic peroxyacids may also be suitable as the peroxy bleaching compound. Such materials normally have the general formula:



wherein R is an alkylene or substituted alkylene group containing from 1 to about 20 carbon atoms, optionally having

an internal amide linkage; or a phenylene or substituted phenylene group; and Y is hydrogen, halogen, alkyl, aryl, an imido-aromatic or non-aromatic group, a COOH or

5



10

group or a quaternary ammonium group.

**[0021]** Typical monoperoxy acids useful herein include, for example:

15

(i) peroxybenzoic acid and ring-substituted peroxybenzoic acids, e.g. peroxy- $\alpha$ -naphthoic acid;

(ii) aliphatic, substituted aliphatic and arylalkyl monoperoxyacids, e.g. peroxyauric acid, peroxysearic acid and N, N-phthaloylaminoperoxy caproic acid (PAP); and

20

(iii) 6-octylamino-6-oxo-peroxyhexanoic acid.

Typical diperoxyacids useful herein include, for example:

(iv) 1,12-diperoxydodecanedioic acid (DPDA);

25

(v) 1,9-diperoxyazelaic acid;

(vi) diperoxybrassicic acid; diperoxysebasic acid and diperoxyisophthalic acid;

(vii) 2-decyldiperoxybutane-1,4-dioic acid; and

30

(viii) 4,4'-sulphonylbisepoxybenzoic acid.

**[0022]** Also inorganic peroxyacid compounds are suitable, such as for example potassium monopersulphate (MPS). If organic or inorganic peroxyacids are used as the peroxygen compound, the amount thereof will normally be within the range of about 2-10% by weight, preferably from 4-8% by weight.

35

**[0023]** Peroxyacid bleach precursors are known and amply described in literature, such as in the British Patents 836988; 864,798; 907,356; 1,003,310 and 1,519,351; German Patent 3,337,921; EP-A-0185522; EP-A-0174132; EP-A-0120591; and U.S. Pat. Nos. 1,246,339; 3,332,882; 4,128,494; 4,412,934 and 4,675,393.

**[0024]** Another useful class of peroxyacid bleach precursors is that of the cationic i.e. quaternary ammonium substituted peroxyacid precursors as disclosed in US Pat. Nos. 4,751,015 and 4,397,757, in EP-A0284292 and EP-A-331,229. Examples of peroxyacid bleach precursors of this class are:

40

2-(N,N,N-trimethyl ammonium) ethyl sodium-4-sulphonphenyl carbonate chloride (SPCC);

N-octyl-N,N-dimethyl-N10-carbophenoxy decyl ammonium chloride (ODC);

45

3-(N,N,N-trimethyl ammonium) propyl sodium-4-sulphophenyl carboxylate; and

N,N,N-trimethyl ammonium toluoyloxy benzene sulphonate.

50

**[0025]** A further special class of bleach precursors is formed by the cationic nitriles as disclosed in EP-A-303,520 and in European Patent Specification No.'s 458,396 and 464,880.

**[0026]** Any one of these peroxyacid bleach precursors can be used in the present invention, though some may be more preferred than others.

55

**[0027]** Of the above classes of bleach precursors, the preferred classes are the esters, including acyl phenol sulphonates and acyl alkyl phenol sulphonates; the acyl-amides; and the quaternary ammonium substituted peroxyacid precursors including the cationic nitriles.

**[0028]** Examples of said preferred peroxyacid bleach precursors or activators are sodium-4-benzoyloxy benzene sulphonate (SBOBS); N,N,N'-tetraacetyl ethylene diamine (TAED); sodium-1-methyl-2-benzoyloxy benzene-4-sul-

phonate; sodium-4-methyl-3-benzoyloxy benzoate; SPCC; trimethyl ammonium toluoyloxy-benzene sulphonate; sodium nonanoyloxybenzene sulphonate (SNOBS); sodium 3,5,5-trimethyl hexanoyloxybenzene sulphonate (STHOBS); and the substituted cationic nitriles.

**[0029]** Other classes of bleach precursors for use with the present invention are found in WO0015750, for example 6-(nonanamidocaproyl)oxybenzene sulphonate.

**[0030]** The precursors may be used in an amount of up to 12%, preferably from 2-10% by weight, of the composition.

**[0031]** The bleaching composition of the present invention has particular application in detergent formulations, especially for laundry cleaning. Accordingly, in another preferred embodiment, the present invention provides a detergent bleach composition comprising a bleaching composition as defined above and additionally a surface-active material, optionally together with detergency builder.

**[0032]** The bleach composition according to the present invention may for example contain a surface-active material in an amount of from 10 to 50% by weight. The surface-active material may be naturally derived, such as soap, or a synthetic material selected from anionic, nonionic, amphoteric, zwitterionic, cationic actives and mixtures thereof. Many suitable actives are commercially available and are fully described in the literature, for example in "Surface Active Agents and Detergents", Volumes I and II, by Schwartz, Perry and Berch.

**[0033]** Typical synthetic anionic surface-actives are usually watersoluble alkali metal salts of organic sulphates and sulphonates having alkyl groups containing from about 8 to about 22 carbon atoms, the term "alkyl" being used to include the alkyl portion of higher aryl groups. Examples of suitable synthetic anionic detergent compounds are sodium and ammonium alkyl sulphates, especially those obtained by sulphating higher ( $C_8-C_{18}$ ) alcohols produced, for example, from tallow or coconut oil; sodium and ammonium alkyl ( $C_9-C_{20}$ ) benzene sulphonates, particularly sodium linear secondary alkyl ( $C_{10}-C_{15}$ ) benzene sulphonates; sodium alkyl glyceryl ether sulphates, especially those ethers of the higher alcohols derived from tallow or coconut oil fatty acid monoglyceride sulphates and sulphonates; sodium and ammonium salts of sulphuric acid esters of higher ( $C_9-C_{18}$ ) fatty alcohol alkylene oxide, particularly ethylene oxide, reaction products; the reaction products of fatty acids such as coconut fatty acids esterified with isethionic acid and neutralised with sodium hydroxide; sodium and ammonium salts of fatty acid amides of methyl taurine; alkane monosulphonates such as those derived by reacting alpha-olefins ( $C_8-C_{20}$ ) with sodium bisulphite and those derived by reacting paraffins with  $SO_2$  and  $Cl_2$  and then hydrolysing with a base to produce a random sulphonate; sodium and ammonium ( $C_7-C_{12}$ ) dialkyl sulphosuccinates; and olefin sulphonates, which term is used to describe material made by reacting olefins, particularly ( $C_{10}-C_{20}$ ) alpha-olefins, with  $SO_3$  and then neutralising and hydrolysing the reaction product. The preferred anionic detergent compounds are sodium ( $C_{10}-C_{15}$ ) alkylbenzene sulphonates, and sodium ( $C_{16}-C_{18}$ ) alkyl ether sulphates.

**[0034]** Examples of suitable nonionic surface-active compounds which may be used, preferably together with the anionic surface-active compounds, include, in particular, the reaction products of alkylene oxides, usually ethylene oxide, with alkyl ( $C_6-C_{22}$ ) phenols, generally 5-25 EO, i.e. 5-25 units of ethylene oxides per molecule; and the condensation products of aliphatic ( $C_8-C_{18}$ ) primary or secondary linear or branched alcohols with ethylene oxide, generally 2-30 EO. Other so-called nonionic surface-actives include alkyl polyglycosides, sugar esters, long-chain tertiary amine oxides, long-chain tertiary phosphine oxides and dialkyl sulphoxides.

**[0035]** Amphoteric or zwitterionic surface-active compounds can also be used in the compositions of the invention but this is not normally desired owing to their relatively high cost. If any amphoteric or zwitterionic detergent compounds are used, it is generally in small amounts in compositions based on the much more commonly used synthetic anionic and nonionic actives.

**[0036]** The detergent bleach composition of the invention will preferably comprise from 1 to 15% wt of anionic surfactant and from 10 to 40% by weight of nonionic surfactant. In a further preferred embodiment, the detergent active system is free from  $C_{16}-C_{12}$  fatty acid soaps.

**[0037]** The bleach composition of the present invention may also contain a detergency builder, for example in an amount of from about 5 to 80% by weight, preferably from about 10 to 60% by weight.

**[0038]** Builder materials may be selected from 1) calcium sequestrant materials, 2) precipitating materials, 3) calcium ion-exchange materials and 4) mixtures thereof.

**[0039]** Examples of calcium sequestrant builder materials include alkali metal polyphosphates, such as sodium tripolyphosphate; nitrilotriacetic acid and its watersoluble salts; the alkali metal salts of carboxymethyloxy succinic acid, ethylene diamine tetraacetic acid, oxydisuccinic acid, mellitic acid, benzene polycarboxylic acids, citric acid; and polyacetal carboxylates as disclosed in US-A-4,144,226 and US-A-4,146,495.

**[0040]** Examples of precipitating builder materials include sodium orthophosphate and sodium carbonate.

**[0041]** Examples of calcium ion-exchange builder materials include the various types of water-insoluble crystalline or amorphous aluminosilicates, of which zeolites are the best known representatives, e.g. zeolite A, zeolite B (also known as zeolite P), zeolite C, zeolite X, zeolite Y and also the zeolite P-type as described in EP-A-0,384,070.

**[0042]** In particular, the compositions of the invention may contain any one of the organic and inorganic builder materials, though, for environmental reasons, phosphate builders are preferably omitted or only used in very small amounts. Typical builders usable in the present invention are, for example, sodium carbonate, calcite/carbonate, the sodium salt

of nitrilotriacetic acid, sodium citrate, carboxymethyloxy malonate, carboxymethyloxy succinate and water-insoluble crystalline or amorphous aluminosilicate builder materials, each of which can be used as the main builder, either alone or in admixture with minor amounts of other builders or polymers as co-builder.

**[0043]** It is preferred that the composition contains not more than 5% by weight of a carbonate builder, expressed as sodium carbonate, more preferably not more than 2.5% by weight to substantially nil, if the composition pH lies in the lower alkaline region of up to 10.

**[0044]** Apart from the components already mentioned, the bleach composition of the present invention can contain any of the conventional additives in amounts of which such materials are normally employed in fabric washing detergent compositions. Examples of these additives include buffers such as carbonates, lather boosters, such as alkanolamides, particularly the monoethanol amides derived from palmkernel fatty acids and coconut fatty acids; lather depressants, such as alkyl phosphates and silicones; anti-redeposition agents, such as sodium carboxymethyl cellulose and alkyl or substituted alkyl cellulose ethers; stabilisers, such as phosphonic acid derivatives (i.e. Dequest® types); fabric softening agents; inorganic salts and alkaline buffering agents, such as sodium sulphate and sodium silicate; and, usually in very small amounts, fluorescent agents; perfumes; enzymes, such as proteases, cellulases, lipases, amylases and oxidases; germicides and colourants.

**[0045]** Transition metal sequestrants such as EDTA, and phosphonic acid derivatives such as EDTMP (ethylene diamine tetra(methylene phosphonate) (same as dequest™ above) may also be included, in addition to the ligand specified, for example to improve the stability sensitive ingredients such as enzymes, fluorescent agents and perfumes, but provided the composition remains bleaching effective.

#### Additional Enzymes

**[0046]** The detergent compositions of the present invention may additionally comprise one or more enzymes, which provide cleaning performance, fabric care and/or sanitation benefits.

**[0047]** Said enzymes include oxidoreductases, transferases, hydrolases, lyases, isomerases and ligases. Suitable members of these enzyme classes are described in Enzyme nomenclature 1992: recommendations of the Nomenclature Committee of the International Union of Biochemistry and Molecular Biology on the nomenclature and classification of enzymes, 1992, ISBN 0-12-227165-3, Academic Press. The most recent information on the nomenclature of enzymes is available on the Internet through the ExpASy WWW server (<http://www.expasy.ch/>)

**[0048]** Examples of the hydrolases are carboxylic ester hydrolase, thioester hydrolase, phosphoric monoester hydrolase, and phosphoric diester hydrolase which act on the ester bond; glycosidase which acts on O-glycosyl compounds; glycosylase hydrolysing N-glycosyl compounds; thioether hydrolase which acts on the ether bond; and exopeptidases and endopeptidases which act on the peptide bond. Preferable among them are carboxylic ester hydrolase, glycosidase and exo- and endopeptidases. Specific examples of suitable hydrolases include (1) exopeptidases such as aminopeptidase and carboxypeptidase A and B and endopeptidases such as pepsin, pepsin B, chymosin, trypsin, chymotrypsin, elastase, enteropeptidase, cathepsin B, papain, chymopapain, ficain, thrombin, plasmin, renin, subtilisin, aspergillopepsin, collagenase, clostripain, kallikrein, gastricsin, cathepsin D, bromelain, chymotrypsin C, urokinase, cucumisin, oryzin, proteinase K, thermomycolin, thermitase, lactocepin, thermolysin, bacillolysin. Preferred among them is subtilisin; (2) glycosidases such as  $\alpha$ -amylase,  $\beta$ -amylase, glucoamylase, isoamylase, cellulase, endo-1,3(4)- $\beta$ -glucanase ( $\beta$ -glucanase), xylanase, dextranase, polygalacturonase (pectinase), lysozyme, invertase, hyaluronidase, pullulanase, neopullulanase, chitinase, arabinosidase, exocellobiohydrolase, hexosaminidase, mycodextranase, endo-1,4- $\beta$ -mannanase (hemicellulase), xyloglucanase, endo- $\beta$ -galactosidase (keratanase), mannanase and other saccharide gum degrading enzymes as described in WO-A-99/09127. Preferred among them are  $\alpha$ -amylase and cellulase; (3) carboxylic ester hydrolase including carboxylesterase, lipase, phospholipase, pectinesterase, cholesterol esterase, chlorophyllase, tannase and wax-ester hydrolase. Preferred among them is lipase.

**[0049]** Examples of transferases and ligases are glutathione S-transferase and acid-thiol ligase as described in WO-A-98/59028 and xyloglycan endotransglycosylase as described in WO-A-98/38288.

**[0050]** Examples of lyases are hyaluronate lyase, pectate lyase, chondroitinase, pectin lyase, alginase II. Especially preferred is pectolyase, which is a mixture of pectinase and pectin lyase.

**[0051]** Examples of the oxidoreductases are oxidases such as glucose oxidase, methanol oxidase, bilirubin oxidase, catechol oxidase, laccase, peroxidases such as ligninase and those described in WO-A-97/31090, monooxygenase, dioxygenase such as lipoxigenase and other oxygenases as described in WO-A-99/02632, WO-A-99/02638, WO-A-99/02639 and the cytochrome based enzymatic bleaching systems described in WO-A-99/02641.

**[0052]** Peroxidases are used in combination with hydrogen peroxide, which can be formulated into a detergent composition as percarbonate or perborate. The hydrogen peroxide may also be generated during the washing and/or rinsing process by an enzymatic system as e.g. described in EP-A-537381.

**[0053]** The activity of oxidoreductases, in particular the phenol oxidising enzymes in a process for bleaching stains on fabrics and/or dyes in solution and/or antimicrobial treatment can be enhanced by adding certain organic compounds,

called enhancers. Examples of enhancers are 2,2'-azo-bis-(3-ethylbenzo-thiazoline-6-sulphonate (ABTS) and Pheno-thiazine-10-propionate (PTP). More enhancers are described in WO-A-94/12619, WO-A-94/12620, WO-A-94/12621, WO-A-97/11217, WO-A-99/23887. Enhancers are generally added at a level of 0.01% to 5% by weight of detergent composition.

**[0054]** A different process for enhancing the efficacy of the bleaching action of oxidoreductases is by targeting them to stains by using antibodies or antibody fragments as described in WO-A-98/56885.

**[0055]** Antibodies can also be added to control enzyme activity as described in WO-A-98/06812.

**[0056]** A preferred combination is a detergent composition comprising of a mixture of conventional detergent enzymes such as protease, amylase, lipase, cutinase and/or cellulase together with one or more plant cell wall degrading enzymes.

**[0057]** Endopeptidases (proteolytic enzymes or proteases) of various qualities and origins and having activity in various pH ranges of from 4-12 are available and can be used in the instant invention. Examples of suitable proteolytic enzymes are the subtilisins, which can be obtained from particular strains of *B. subtilis*, *B. lentus*, *B. amyloliquefaciens* and *B. licheniformis*, such as the commercially available subtilisins Savinase™, Alcalase™, Relase™, Kannase™ and Everlase™ as supplied by Novo Industri A/S, Copenhagen, Denmark or Purafect™, PurafectOxP™ and Properase™ as supplied by Genencor International. Chemically or genetically modified variants of these enzymes are included such as described in WO-A-99/02632 pages 12 to 16 and in WO-A-99/20727 and also variants with reduced allergenicity as described in WO-A-99/00489 and WO-A-99/49056.

**[0058]** Suitable lipases include those of bacterial or fungal origin as described in WO-A-99/11770 pages 33,34, such as the commercially available Lipolase™, Lipolase Ultra™, LipoPrime™, from Novo Nordisk, or Lipomax™ from Genencor. Chemically or genetically modified variants of these enzymes are included.

**[0059]** Suitable amylases include those of bacterial or fungal origin. Chemically or genetically modified variants of these enzymes are included as described in WO-A-99/02632 pages 18,19. Commercial cellulase are sold under the tradename Purastar™, Purastar OxAm™ (formerly Purafact Ox Am™) by Genencor; Termamyl™, Fungamyl™ and Duramyl™, all available from Novo Nordisk A/S.

**[0060]** Suitable cellulases include those of bacterial or fungal origin. Chemically or genetically modified variants of these enzymes are included as described in WO-A-99/02632 page 17. Particularly useful cellulases are the endoglucanases such as the EGIII from *Trichoderma longibrachiatum* as described in WO-A-94/21801 and the E5 from *Thermomonospora fusca* as described in WO-A-97/20025. Endoglucanases may consist of a catalytic domain and a cellulose binding domain or a catalytic domain only. Preferred cellulolytic enzymes are sold under the tradename Carezyme™, Celluzyme™ and Endolase™ by Novo Nordisk A/S; Puradax™ is sold by Genencor and KAC™ is sold by Kao corporation, Japan.

**[0061]** Detergent enzymes are usually incorporated in an amount of 0.00001% to 2%, and more preferably 0.001% to 0.5%, and even more preferably 0.01% to 0.2% in terms of pure enzyme protein by weight of the composition. Detergent enzymes are commonly employed in the form of granules made of crude enzyme alone or in combination with other components in the detergent composition. Granules of crude enzyme are used in such an amount that the pure enzyme is 0.001 to 50 weight percent in the granules. The granules are used in an amount of 0.002 to 20 and preferably 0.1 to 3 weight percent. Granular forms of detergent enzymes are known as Enzoguard™ granules, prills, marumes or T-granules. Granules can be formulated so as to contain an enzyme protecting agent (e.g. oxidation scavengers) and/or a dissolution retardant material. Other suitable forms of enzymes are liquid forms such as the "L" type liquids from Novo Nordisk, slurries of enzymes in nonionic surfactants such as the "SL" type sold by Novo Nordisk and microencapsulated enzymes marketed by Novo Nordisk under the tradename "LDP" and "CC".

**[0062]** The enzymes can be added as separate single ingredients (prills, granulates, stabilised liquids, etc. containing one enzyme) or as mixtures of two or more enzymes (e.g. cogranulates). Enzymes in liquid detergents can be stabilised by various techniques as for example disclosed in US patents 4,261,868 and 4,318,818.

**[0063]** The detergent compositions of the present invention may additionally comprise one or more biologically active peptides such as swollenin proteins, expansins, bacteriocins and peptides capable of binding to stains.

**[0064]** In a particularly preferred embodiment the method of the present invention is carried out on a laundry fabric using aqueous treatment liquor. In particular the treatment may be effected in, or as an adjunct to, an essentially conventional wash cycle for cleaning laundry. More preferably, the treatment is carried out in an aqueous detergent wash liquor. The bleaching composition can be delivered into the wash liquor from a powder, granule, pellet, tablet, block, bar or other such solid form. The solid form can comprise a carrier, which can be particulate, sheet-like or comprise a three-dimensional object. The carrier can be dispersible or soluble in the wash liquor or may remain substantially intact. In other embodiments, the bleaching composition can be delivered into the wash liquor from a paste, gel or liquid concentrate. Other means for ensuring that the bleaching composition is present in the wash liquor may be envisaged.

**[0065]** For example, it is envisaged that the bleaching composition can be presented in the form of a body from which it is slowly released during the whole or part of the laundry process. Such release can occur over the course of a single wash or over the course of a plurality of washes. In the latter case it is envisaged that the bleaching composition can be released from a carrier substrate used in association with the wash process, e.g. from a body placed in the dispenser

## EP 1 358 309 B1

drawer of a washing machine, elsewhere in the delivery system or in the drum of the washing machine. When used in the drum of the washing machine the carrier can be freely moving or fixed relative to the drum. Such fixing can be achieved by mechanical means, for example by barbs that interact with the drum wall, or employ other forces, for example a magnetic force. The modification of a washing machine to provide for means to hold and retain such a carrier is envisaged similar means being known from the analogous art of toilet block manufacture. Freely moving carriers such as shuttles for dosage of surfactant materials and/or other detergent ingredients into the wash can comprise means for the release of the bleaching composition into the wash.

**[0066]** The present invention is not limited to those circumstances in which a washing machine is employed, but can be applied where washing is performed in some alternative vessel. In these circumstances it is envisaged that the bleaching composition can be delivered by means of slow release from the bowl, bucket or other vessel which is being employed, or from any implement which is being employed, such as a brush, bat or dolly, or from any suitable applicator.

**[0067]** Suitable pre-treatment means for application of the bleaching composition to the textile material prior to the main wash include sprays, pens, roller-ball devices, bars, soft solid applicator sticks and impregnated cloths or cloths containing microcapsules. Such means are well known in the analogous art of deodorant application and/or in spot treatment of textiles. Similar means for application are employed in those embodiments where the bleaching composition is applied after the main washing and/or conditioning steps have been performed, e.g. prior to or after ironing or drying of the cloth. For example, the bleaching composition may be applied using tapes, sheets or sticking plasters coated or impregnated with the substance, or containing microcapsules of the substance. The bleaching composition may for example be incorporated into a drier sheet so as to be activated or released during a tumble-drier cycle, or the substance can be provided in an impregnated or microcapsule-containing sheet so as to be delivered to the textile when ironed.

**[0068]** Throughout the description and claims generic groups have been used, for example alkyl, alkoxy, aryl. Unless otherwise specified the following are preferred group restrictions that may be applied to generic groups found within compounds disclosed herein:

alkyl: linear and branched C1-C8-alkyl,

alkenyl: C2-C6-alkenyl,

cycloalkyl: C3-C8-cycloalkyl,

aryl: selected from homoaromatic compounds having a molecular weight under 300,

heteroaryl: selected from the group consisting of: pyridinyl; pyrimidinyl; pyrazinyl; triazolyl; pyridazinyl; 1,3,5-triazinyl; quinolinyl; isoquinolinyl; quinoxalinyl; imidazolyl; pyrazolyl; benzimidazolyl; thiazolyl; oxazolidinyl; pyrrolyl; carbazolyl; indolyl; and isoindolyl, wherein the heteroaryl may be connected to the compound via any atom in the ring of the selected heteroaryl,

heterocycloalkyl: selected from the group consisting of: pyrrolinyl; pyrrolidinyl; morpholinyl; piperidinyl; piperazinyl; hexamethylene imine; 1,4-piperazinyl; tetrahydrothiophenyl; tetrahydrofuranyl; 1,4,7-triazacyclononanyl; 1,4,8,11-tetraazacyclotetradecanyl; 1,4,7,10,13-pentaazacyclopentadecanyl; 1,4-diaza-7-thiacyclononanyl; 1,4-diaza-7-oxa-cyclononanyl; 1,4,7,10-tetraazacyclododecanyl; 1,4-dioxanyl; 1,4,7-trithiacyclononanyl; tetrahydropyranyl; and oxazolidinyl, wherein the heterocycloalkyl may be connected to the compound via any atom in the ring of the selected heterocycloalkyl,

carboxylate derivative: the group -C(O)OR, wherein R is selected from: hydrogen; C1-C6-alkyl; phenyl; C1-C6-alkyl-C6H5; Li; Na; K; Cs; Mg; and Ca,

carbonyl derivative: the group -C(O)R, wherein R is selected from: hydrogen; C1-C6-alkyl; phenyl; C1-C6-alkyl-C6H5 and amine (to give amide) selected from the group: -NR'<sub>2</sub>, wherein each R' is independently selected from:

hydrogen; C1-C6-alkyl; C1-C6-alkyl-C6H5; and phenyl, wherein when both R' are C1-C6-alkyl both R' together may form an -NC<sub>3</sub> to an -NC<sub>5</sub> heterocyclic ring with any remaining alkyl chain forming an alkyl substituent to the heterocyclic ring,

sulphonate: the group -S(O)<sub>2</sub>OR, wherein R is selected from: hydrogen; C1-C6-alkyl; phenyl; C1-C6-alkyl-C6H5; Li; Na; K; Cs; Mg; and Ca,

**[0069]** Unless otherwise specified the following are more preferred group restrictions that may be applied to groups



found within compounds disclosed herein:

alkyl: linear and branched C1-C6-alkyl,

5 alkenyl: C3-C6-alkenyl,

cycloalkyl: C6-C8-cycloalkyl,

10 aryl: selected from group consisting of: phenyl; biphenyl; naphthalenyl; anthracenyl; and phenanthrenyl,

heteroaryl: selected from the group consisting of: pyridinyl; pyrimidinyl; quinolinyl; pyrazolyl; triazolyl; isoquinolinyl; imidazolyl; and oxazolidinyl, wherein the heteroaryl may be connected to the compound via any atom in the ring of the selected heteroaryl,

15 heterocycloalkyl: selected from the group consisting of: pyrrolidinyl; morpholinyl; piperidinyl; piperidinyll; 1,4-piperazinyl; tetrahydrofuranyl; 1,4,7-triazacyclononanyl; 1,4,8,11-tetraazacyclotetradecanyl; 1,4,7,10,13-pentaazacyclopentadecanyl; 1,4,7,10-tetraazacyclododecanyl; and piperazinyl, wherein the heterocycloalkyl may be connected to the compound via any atom in the ring of the selected heterocycloalkyl,

20 carboxylate derivative: the group -C(O)OR, wherein R is selected from hydrogen; Na; K; Mg; Ca; C1-C6-alkyl; and benzyl,

**[0070]** The invention will now be further illustrated by way of the following non-limiting examples:

## 25 EXAMPLES

### **Syntheses of Compounds**

#### **[(MeN4Py)FeCl]Cl**

30 **[0071]** The ligand N,N-bis(pyridin-2-yl-methyl)-1,1-bis(pyridin-2-yl)-1-aminoethane (MeN4py) was prepared as described in EP 0 909 809 A2.

The ligand MeN4Py (33.7 g; 88.5mmoles) was dissolved in dry methanol (500ml). Small portions of FeCl<sub>2</sub>·4H<sub>2</sub>O (0.95 eq; 16.7 g; 84.0 mmoles) were added, yielding a clear red solution. After addition, the solution was stirred for 30 minutes at room temperature, after which the methanol was removed (rotary-evaporator). The dry solid was ground and 150 ml of ethylacetate was added and the mixture was stirred until a fine red powder was obtained. This powder was washed twice with ethyl acetate, dried in the air and further dried under reduced pressure vacuum at 40 °C. El. Anal. Calc. for [Fe(MeN4py)Cl]Cl·2H<sub>2</sub>O: C 53.03; H 5.16; N 12.89; Cl 13.07; Fe 10.01%. Found C 52.29/ 52.03; H 5.05/5.03; N 12.55/12.61; Cl 12.73/12.69; Fe: 10.06/10.01%.

40 **[0072]** L1 has been prepared according to literature procedures (Bernal, J.; et al. J. Chem. Soc., Dalton Trans. 1995, 3667-3675). L10 has been prepared according to literature procedures (Brennan, et al., Inorg. Chem., 30, 1937 (1991).

#### Reaction procedure for ligands L2, L3, L4, L5, L6, L7, L8, and L9.

45 **[0073]** The general procedure of these ligands is as follows. To a 25 ml glass vial was added the ethylenediamine (1 mmol), and aldehyde (4 mmol).

**[0074]** The following commercially available ethylene diamines were used for the ligand syntheses:

L2: N-methylethylenediamine;

50 L3: N-ethyl-ethylenediamine;

L4: N,N'-dimethylethylenediamine;

L5 and L6: N-(1-propan-2-ol)ethylenediamine;

L7: N,N-diethyldiethylenetriamine;

L8: N-3-propan-1-ol)ethylenediamine;

55 L9: N-hexylethylenediamine.

**[0075]** The following commercially available aldehydes were used for the ligand syntheses:

## EP 1 358 309 B1

L2, L3, L4, and L5: 2-imidazolecarboxaldehyde;  
 L6, and L8: 1-methyl-2-imidazolecarboxaldehyde;  
 L7: 5-methyl-imidazol-4-methylcarboxaldehyde  
 L9 4-formylimidazole

5  
**[0076]** The above-mentioned vial was sealed with a cap and the solution was then shaken for 2 h to allow the imine formation. The mixture was treated with NaCNBH<sub>3</sub> (3.3 mmol), adjusted to pH 6 with acetic acid and shaken for 38 h. The mixture was quenched with 3 ml of 2 M HCl solution and adjusted to pH > 13 with a 7M NaOH solution. The mixture was extracted with 3x10 ml of dichloromethane, dried over sodium sulfate and evaporated under reduced pressure. Yields are typically around 50%. Purities are greater than 90 % as established by HPLC/MS.

15  
 L2: m/z 315.5 (M+H<sup>+</sup>)  
 L3: m/z 329.4 (M+H<sup>+</sup>)  
 L4: m/z 249.3 (M+H<sup>+</sup>)  
 L5: m/z 359.5 (M+H<sup>+</sup>)  
 L6: m/z 401.5 (M+H<sup>+</sup>)  
 L7: m/z 443.5 (M+H<sup>+</sup>)  
 L8: m/z 401.5 (M+H<sup>+</sup>)  
 L9: m/z 385.5 (M+H<sup>+</sup>)

### Tomato stain bleaching

25  
**[0077]** In an aqueous solution containing 10 mM carbonate buffer (pH 10) with 0.6 g/l NaLAS (linear alkylbenzene sulfonate), tomato-soya oil stained cloths were added and kept in contact with the solution under agitation for 30 minutes at 30 °C. In comparative experiments, the same experiments were done by addition of 10 μM of compound 1, or 10 μM of transition-metal salt in combination with 20 μM ligand (L1-L10), or the ligand alone without addition of metal salts (L1-L10) referred to in the table below.

30  
**[0078]** After the wash, the cloths were rinsed with water and subsequently dried at 30 °C and the change in colour was measured immediately after drying with a Linotype-Hell scanner (ex Linotype) ("t=0") and after 1 day storage in the dark ("t=1"). The change in colour (including bleaching) is expressed as the ΔE value. The measured colour difference (ΔE) between the washed cloth and the unwashed cloth is defined as follows:

$$\Delta E = [(\Delta L)^2 + (\Delta a)^2 + (\Delta b)^2]^{1/2}$$

35  
 wherein ΔL is a measure for the difference in darkness between the washed and unwashed test cloth; Δa and Δb are measures for the difference in redness and yellowness respectively between both cloths. With regard to this colour measurement technique, reference is made to Commission Internationale de l'Eclairage (CIE); Recommendation on Uniform Colour Spaces, colour difference equations, psychometric colour terms, supplement no 2 to CIE Publication, no 15, Colorimetry, Bureau Central de la CIE, Paris 1978. The results are shown below in the Table below.

40

	t=0	t=1
Blank	16	16
FeMeN4pyCl <sub>2</sub> (comparative)	5	4
MeN4py/Fe perchlorate (comparative)	7	5
COMPARATIVE EXAMPLE (NO IMIDAZOLE GROUPS)		
N-methyl-N,N',N'-tris(pyridin-2ylmethyl)ethylenediamine (L1)	8	10
L1+ Fe perchlorate	9	9
L1+ Cu perchlorate	13	13
L1+ Co perchlorate	10	7
N-Methyl-N,N',N'-Tris(imidazol-2ylmethyl)-ethylenediamine (L2)		
L2+ Fe perchlorate	2	2
L2+ Cu perchlorate	3	3
L2+ Mn perchlorate	5	4
L2+ Co perchlorate	7	5
	15	4

55

EP 1 358 309 B1

(continued)

	t=0	t=1	
5	N-ethyl-N,N',N'-Tris(imidazol-2-ylmethyl)-ethylenediamine (L3)	3	3
	L3 Fe perchlorate	3	3
	L3 Cu perchlorate	4	5
	L3 Mn perchlorate	7	5
	L3 Co perchlorate	6	5
10	N,N'-dimethyl-N,N'-bis(imidazol-2-ylmethyl)-ethylenediamine (L4)	12	3
	N-(1-propan-2-ol)-N,N',N'-Tris(imidazol-2-ylmethyl)-ethylenediamine (L5)	-	3
	L5 + Fe perchlorate	-	5
	L5+ Cu perchlorate	-	4
15	L5+ Mn perchlorate	-	4
	L5+ Co perchlorate	-	4
	N-(1-propan-2-ol)-N,N',N'-Tris(1-methylimidazol-2-ylmethyl)-ethylenediamine (L6)	5	3
	L6 + Fe perchlorate	-	5
20	L6+ Cu perchlorate	-	4
	L6+ Mn perchlorate	-	5
	L6+ Co perchlorate	14	4
	N,N-diethyl-N',N'',N''-Tris(5-methylimidazol-4-ylmethyl)-diethylenetriamine (L7)	-	-
	L7 + Fe perchlorate	-	-
25	L7+ Cu perchlorate	13	8
	L7+ Mn perchlorate	14	8
	L7+ Co perchlorate	14	6
	N-(3-propan-1-ol)-N,N',N'-Tris(1-methylimidazol-2-ylmethyl)-ethylenediamine (L8)	4	3
30	L8 + Fe perchlorate	4	-
	L8+ Cu perchlorate	5	3
	L8+ Mn perchlorate	-	3
	L8+ Co perchlorate	11	3
35	N-hexyl-N,N',N'-Tris(imidazol-2-ylmethyl)-ethylenediamine (L9)	9	6
	L9 + Fe perchlorate	-	-
	L9+ Cu perchlorate	12	9
	L9+ Mn perchlorate	14	7
40	L9+ Co perchlorate	13	6

**[0079]** The following table illustrates the present invention in peroxy bleaching mode. The experiments were conducted in substantially the same manner as described above but in the presence of 10 mmol/l hydrogen peroxide. All data given were obtained immediately after the wash.  $\Delta E$  measurements are given with respect to white.

	$\Delta E$	
45	Blank	16
	FeMeN <sub>4</sub> pyCl <sub>2</sub> (comparative)	5
50	COMPARATIVE EXAMPLE (NO IMIDAZOLE GROUPS) N,N,N',N'-tetrakis(pyridin-2-ylmethyl)ethylenediamine (L10)	8
	N-Methyl-N,N',N'-Tris(imidazol-2-ylmethyl)-ethylenediamine (L2)	2
55	L2+ Fe perchlorate	3
	L2+ Cu perchlorate	3
	L2+ Mn perchlorate	2
	L2+ Co perchlorate	3

EP 1 358 309 B1

(continued)

	$\Delta E$
N-ethyl-N,N',N'-Tris(imidazol-2ylmethyl)-ethylenediamine (L3)	3
L3 Fe perchlorate	2
L3 Cu perchlorate	2
L3 Mn perchlorate	2
L3 Co perchlorate	3
N-(1-propan-2-ol)-N,N',N'-Tris(1-methyl-imidazol-2ylmethyl)-ethylenediamine (L6)	4
L6 + Fe perchlorate	7
L6 + Cu perchlorate	8
L6 + Mn perchlorate	7
L6 + Co perchlorate	7
N-(3-propan-1-ol)-N,N',N'-Tris(1-methyl-imidazol-2ylmethyl)-ethylenediamine (L8)	3
L8 + Fe perchlorate	4
L8+ Cu perchlorate	7
L8+ Mn perchlorate	2
L8+ Co perchlorate	3

**[0080]** The above tabulated results demonstrate a substantial advantage secured by the selected group of compounds in both air bleaching and bleaching with peroxy species.


**Claims**

1. A laundry bleach composition, capable of bleaching in an air bleaching mode or peroxy bleaching mode, comprising a ligand of general formula (L):



wherein:


X is selected from  $-CH_2CH_2-$ ,  $-CH_2CH_2CH_2-$ ,  $-CH_2C(OH)HCH_2-$ ; and, R<sup>1</sup> independently represents a group selected from: alkyl, cycloalkyl, heterocycloalkyl, heteroaryl, aryl and arylalkyl groups optionally substituted with a substituent selected from hydroxy, alkoxy, phenoxy, carboxylate, carboxamide, carboxylic ester, sulphonate, amine, alkylamine and  $N^+(R^4)_3$ , wherein R<sup>4</sup> is selected from hydrogen, alkanyl, alkenyl, arylalkanyl, arylalkenyl, oxyalkanyl, oxyalkenyl, aminoalkanyl, aminoalkenyl, alkanyl ether, alkenyl ether, and  $-CY_2-R^2$ , in which Y is independently selected from H, CH<sub>3</sub>, C<sub>2</sub>H<sub>5</sub>, C<sub>3</sub>H<sub>7</sub> and R<sup>2</sup> is independently selected from: an optionally substituted



wherein R<sub>3</sub> is independently selected from: H, alkyl, and benzyl, and wherein at least two of R<sup>1</sup> are  $-CY_2-R^2$ .

2. A bleaching composition according to claim 1, wherein R<sub>3</sub> is independently selected from: H and methyl.
3. A bleaching composition according to any preceding claim, wherein the substituent of the optionally substituted

5




10

is selected from: CH<sub>3</sub>, C<sub>2</sub>H<sub>5</sub>, C<sub>3</sub>H<sub>7</sub> and C<sub>4</sub>H<sub>9</sub>, CH<sub>2</sub>-C<sub>6</sub>H<sub>5</sub>.

4. A bleaching composition according to any preceding claim, wherein R<sup>2</sup> is independently selected from:


15



20

and


25



30

5. A bleaching composition according to claim 4, wherein R<sup>2</sup> is independently selected from:

35



40

6. A bleaching composition according to any preceding claim, wherein all -CY<sub>2</sub>-R<sup>2</sup> are identical.

45

7. A bleaching composition according to any preceding claim, wherein at least three of R<sup>1</sup> are -CY<sub>2</sub>-R<sup>2</sup>.

8. A bleaching composition according to any preceding claim, wherein X is -CH<sub>2</sub>CH<sub>2</sub>-.

50

9. A bleaching composition according to claim 1, wherein the ligand is selected from N-Methyl-N,N',N'-Tris(imidazol-2-ylmethyl)-ethylenediamine; N-ethyl-N,N',N'-Tris(imidazol-2-ylmethyl)-ethylenediamine; N,N'-dimethyl-N,N'-bis(imidazol-2-ylmethyl)-ethylenediamine; N-(1-propan-2-ol)-N,N',N'-Tris(imidazol-2-ylmethyl)-ethylenediamine; N-(1-propan-2-ol)-N,N',N'-Tris(1-methylimidazol-2-ylmethyl)-ethylenediamine; N,N-diethyl-N',N'',N''-Tris(5-methylimidazol-4-ylmethyl)-diethylenetriamine; N-(3-propan-1-ol)-N,N',N'-Tris(1-methylimidazol-2-ylmethyl)-ethylenediamine; and, N-hexyl-N,N',N'-Tris(imidazol-2-ylmethyl)-ethylenediamine.

55

10. A bleaching composition according to claim 1, wherein the ligand of general formula (L) is present in the form of a complex of the general formula (A):



in which

M represents iron in the II, III, IV or V oxidation state, manganese in the II, III, IV, VI or VII oxidation state, copper in the I, II or III or cobalt in the I, II or III oxidation state;

X represents a coordinating species;

n represents zero or an integer in the range from 0 to 3;

z represents the charge of the complex and is an integer which can be positive, zero or negative;

Y represents a counter ion, the type of which is dependent on the charge of the complex;


$$q = z / [\text{charge Y}].$$

11. A bleaching composition according to any preceding claim, wherein the composition comprises a surface-active material and a detergency builder.
12. A bleaching composition according to any preceding claim, wherein in an aqueous solution at least 10 %, preferably at least 50 % and optimally at least 90 % of any bleaching of a substrate is effected by oxygen sourced from the air.
13. A bleaching composition according to any preceding claim, further comprising a sequestrant and wherein in an aqueous solution at least 90 % of any bleaching of the substrate is effected by a peroxy species not derived directly from atmospheric oxygen, preferably 99% and most preferably 100%.
14. A method of bleaching a substrate comprising the step of applying to the substrate, in an aqueous medium, a bleaching composition as defined in any preceding claim.
15. A ligand selected from the group consisting of:

N-Methyl-N,N',N''-Tris (imidazol-2ylmethyl)-ethylenediamine; N-ethyl-N,N',N''-Tris(imidazol-2ylmethyl)-ethylenediamine; N,N' -dimethyl-N,N' -bis (imidazol-2-ylmethyl)-ethylenediamine; N-(1-propan-2-ol)-N,N',N''-Tris(imidazol-2ylmethyl)-ethylenediamine; N- (1-propan-2-ol) -N,N',N''-Tris (1-methylimidazol-2ylmethyl)-ethylenediamine; N,N-diethyl-N',N'',N'''-Tris (5-methyl-imidazol-4ylmethyl)-diethylenetriamine; N-(3-propan-1-ol) -N,N',N'' -Tris(1-methyl-imidazol-2ylmethyl)-ethylenediamine; and, N-hexyl-N,N',N''-Tris(imidazol-2ylmethyl)-ethylenediamine.

## Patentansprüche

1. Wäschebleichmittelzusammensetzung, die in einem luftbleichenden Modus oder peroxybleichenden Modus bleichen kann, umfassend einen Liganden der allgemeinen Formel (L):




worin

X ausgewählt ist aus  $-\text{CH}_2\text{CH}_2-$ ,  $-\text{CH}_2\text{CH}_2\text{CH}_2-$ ,  $-\text{CH}_2\text{C}(\text{OH})\text{HCH}_2-$  und

$\text{R}^1$  unabhängig eine Gruppe wiedergibt, ausgewählt aus: Alkyl-, Cycloalkyl-, Heterocycloalkyl-, Heteroaryl-, Aryl- und Arylalkylgruppen, gegebenenfalls substituiert mit einem Substituenten, ausgewählt aus Hydroxy, Alkoxy, Phenoxy, Carboxylat, Carboxamid, Carbonsäureester, Sulfonat, Amin, Alkylamin und  $\text{N}^+(\text{R}^4)_3$ , worin  $\text{R}^4$  ausgewählt ist aus Wasserstoff, Alkanyl, Alkenyl, Arylalkanyl, Arylalkenyl, Oxyalkanyl, Oxyalkenyl, Aminoalkanyl, Aminoalkenyl, Alkanylether, Alkenylether und  $-\text{CY}_2-\text{R}^2$ , worin Y unabhängig aus H,  $\text{CH}_3$ ,  $\text{C}_2\text{H}_5$ ,  $\text{C}_3\text{H}_7$  ausgewählt ist und  $\text{R}^2$  unabhängig ausgewählt ist aus: einem gegebenenfalls substituierten


5



worin R<sup>3</sup> unabhängig ausgewählt ist aus: H, Alkyl und Benzyl und worin mindestens zwei von R<sup>1</sup> -CY<sub>2</sub>-R<sup>2</sup> darstellen.

- 10 **2.** Bleichende Zusammensetzung nach Anspruch 1, worin R<sup>3</sup> unabhängig ausgewählt ist aus: H und Methyl.
- 15 **3.** Bleichende Zusammensetzung nach einem vorangehenden Anspruch, worin der Substituent von gegebenenfalls substituiertem

15




20

ausgewählt ist aus: CH<sub>3</sub>, C<sub>2</sub>H<sub>5</sub>, C<sub>3</sub>H<sub>7</sub> und C<sub>4</sub>H<sub>9</sub>, CH<sub>2</sub>-C<sub>6</sub>H<sub>5</sub>.


- 25 **4.** Bleichende Zusammensetzung nach einem vorangehenden Anspruch, worin R<sup>2</sup> unabhängig ausgewählt ist aus

30



35 und


40



45

- 5.** Bleichende Zusammensetzung nach Anspruch 4, worin R<sup>2</sup> unabhängig ausgewählt ist aus:

50



55

- 6.** Bleichende Zusammensetzung nach einen vorangehenden Anspruch, worin alle -CY<sub>2</sub>-R<sup>2</sup> identisch sind.
- 7.** Bleichende Zusammensetzung nach einem vorangehenden Anspruch, worin mindestens drei von R<sup>1</sup> -CY<sub>2</sub>R<sup>2</sup> darstellen.

## EP 1 358 309 B1

8. Bleichende Zusammensetzung nach einem vorangehenden Anspruch, worin X -CH<sub>2</sub>CH<sub>2</sub>- darstellt.
9. Bleichende Zusammensetzung nach Anspruch 1, worin der Ligand aus N-Methyl-N,N',N'-tris(imidazol-2-ylmethyl)ethylendiamin; N-Ethyl-N,N',N'-tris(imidazol-2-ylmethyl)ethylendiamin; N,N'-Dimethyl-N,N'-bis(imidazol-2-ylmethyl)ethylendiamin; N-(1-Propan-2-ol)-N,N',N'-tris(imidazol-2-ylmethyl)ethylendiamin; N-(1-Propan-2-ol)-N,N',N'-tris(1-methylimidazol-2-ylmethyl)ethylendiamin; N,N-Diethyl-N',N'',N''-tris(5-methylimidazol-4-ylmethyl)diethylen-triamin; N-(3-Propan-1-ol)-N,N',N'-tris(1-methylimidazol-2-ylmethyl)ethylendiamin und N-Hexyl-N,N',N'-tris(imidazol-2-ylmethyl)ethylendiamin ausgewählt ist.
10. Bleichende Zusammensetzung nach Anspruch 1, worin der Ligand der allgemeinen Formel (L) in Form eines Komplexes der allgemeinen Formel (A):



vorliegt,  
worin M Eisen im Oxidationszustand II, III, IV oder V, Mangan im Oxidationszustand II, III, IV, VI oder VII, Kupfer im I-, II- oder III- oder Kobalt I-, II- oder III-Oxidationszustand wiedergibt;  
X eine koordinierende Spezies wiedergibt;  
n null oder eine ganze Zahl im Bereich von 0 bis 3 wiedergibt;  
z die Ladung des Komplexes wiedergibt und eine ganze Zahl darstellt, die positiv, null oder negativ sein kann;  
Y ein Gegenion wiedergibt, wobei der Typ davon von der Ladung des Komplexes abhängt,

$$q = z / [\text{Ladung Y}].$$

11. Bleichende Zusammensetzung nach einem vorangehenden Anspruch, worin die Zusammensetzung ein oberflächenaktives Material und einen Waschmittelbuilder umfasst.
12. Bleichende Zusammensetzung nach einem vorangehenden Anspruch, worin in einer wässrigen Lösung mindestens 10 %, vorzugsweise mindestens 50 % und optimal mindestens 90 % von beliebigem Bleichen von einem Substrat durch aus der Luft stammendem Sauerstoff bewirkt werden.
13. Bleichende Zusammensetzung nach einem vorangehenden Anspruch, weiterhin umfassend ein Maskierungsmittel und worin in einer wässrigen Lösung mindestens 90 % von beliebigem Bleichen des Substrats durch eine Peroxylspezies, die nicht direkt von atmosphärischem Sauerstoff abgeleitet ist, vorzugsweise 99 % und besonders bevorzugt 100 % bewirkt werden.
14. Verfahren zum Bleichen eines Substrats, umfassend den Schritt des Auftragens einer bleichenden Zusammensetzung, wie in einem vorangehenden Anspruch definiert, in einem wässrigen Medium auf das Substrat.
15. Ligand, ausgewählt aus der Gruppe, bestehend aus:

N-Methyl-N,N',N'-tris(imidazol-2-ylmethyl)ethylendiamin; N-Ethyl-N,N',N'-tris(imidazol-2-ylmethyl)ethylendiamin; N,N'-Dimethyl-N,N'-bis(imidazol-2-ylmethyl)ethylendiamin; N-(1-Propan-2-ol)-N,N',N'-tris(imidazol-2-ylmethyl)ethylendiamin; N-(1-Propan-2-ol)-N,N',N'-tris(1-methylimidazol-2-ylmethyl)ethylendiamin; N,N-Diethyl-N',N'',N''-tris(5-methylimidazol-4-ylmethyl)diethylen-triamin; N-(3-Propan-1-ol)-N,N',N'-tris(1-methylimidazol-2-ylmethyl)ethylendiamin und N-Hexyl-N,N',N'-tris(imidazol-2-ylmethyl)ethylendiamin.

### Revendications

1. Composition blanchissante pour le linge capable de blanchir en mode de blanchiment par l'air ou en mode de blanchiment par le peroxyde, comprenant un ligand de formule générale (L) :




dans laquelle X est choisi parmi -CH<sub>2</sub>CH<sub>2</sub>-, -CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>-, -CH<sub>2</sub>C(OH)HCH<sub>2</sub>-, et



EP 1 358 309 B1

R<sup>1</sup> représente indépendamment un groupe choisi parmi les groupes alkyles, cycloalkyles, hétéroaryles, aryles et arylalkyles facultativement substitués avec un substituant sélectionné parmi l'hydroxy, l'alkoxy, le phénoxy, le carboxylate, le carboxamide, l'ester carboxylique, le sulfonate, l'amine, l'alkylamine et N<sup>+</sup>(R<sup>4</sup>)<sub>3</sub>, dans lequel R<sup>4</sup> est choisi parmi l'hydrogène, l'alcanyle, l'alcényle, l'arylalcanyle, l'arylalcényle, l'oxyalcanyle, l'oxyalcényle, l'aminoalcanyle, l'aminoalcényle, l'éther d'alcanyle, l'éther d'alcényle et -CY<sub>2</sub>- R<sup>2</sup> dans lequel Y est indépendamment choisi parmi H, CH<sub>3</sub>, C<sub>2</sub>H<sub>5</sub>, C<sub>3</sub>H<sub>7</sub> et R<sup>2</sup> est indépendamment sélectionné parmi :




optionnellement substitués, dans lesquels R<sup>3</sup> est indépendamment choisi parmi : H, l'alkyle et le benzyle, et dans lesquels au moins deux des groupes R<sup>1</sup> sont -CY<sub>2</sub>- R<sup>2</sup>.

20

2. Composition blanchissante selon la revendication 1, dans laquelle R<sup>3</sup> est indépendamment choisi parmi H et le méthyle.

25


3. Composition blanchissante selon l'une quelconque des revendications précédentes, dans laquelle le substituant du




optionnellement substitué est choisi parmi : CH<sub>3</sub>, C<sub>2</sub>H<sub>5</sub>, C<sub>3</sub>H<sub>7</sub> et C<sub>4</sub>H<sub>9</sub>, CH<sub>2</sub>-C<sub>6</sub>H<sub>5</sub>.

40

4. Composition blanchissante selon l'une quelconque des revendications précédentes, dans laquelle R<sup>2</sup> est indépendamment choisi parmi :




et



5. Composition blanchissante selon la revendication 4, dans laquelle R<sup>2</sup> est indépendamment choisi parmi :

5



10

6. Composition blanchissante selon l'une quelconque des revendications précédentes, dans laquelle tous les groupes -CY<sub>2</sub> - R<sup>2</sup> sont identiques.

15

7. Composition détergente selon l'une quelconque des revendications précédentes, dans laquelle au moins trois des groupes R<sup>1</sup> sont -CY<sub>2</sub> - R<sup>2</sup>.

8. Composition blanchissante selon l'une quelconque des revendications précédentes, dans laquelle X est -CH<sub>2</sub> - CH<sub>2</sub>.

20

9. Composition blanchissante selon la revendication 1, dans laquelle le ligand est choisi parmi la N - méthyl - N,N',N' - tris (imidazol-2ylméthyl) - éthylène diamine ; la N - éthyl - N,N',N' - tris (imidazol-2ylméthyl) - éthylène diamine ; la N,N' - diméthyl - N,N' - bis (imidazol - 2 - ylméthyl) - éthylène diamine ; la N - (1 - propan - 2 - ol) - N,N',N' - tris (imidazol - 2ylméthyl) - éthylène diamine ; la N - (1 - propan - 2 - ol) - N,N',N' - tris (1 - méthyl - imidazol - 2ylméthyl) - éthylène diamine ; la N,N - diéthyl - N', N'', N'' - tris (5 - méthyl - imidazol - 4ylméthyl) - diéthylène triamine ; la N - (3 - propan - 1 - ol) - N,N',N' - tris (1 - méthyl - imidazol - 2ylméthyl) - éthylène diamine ; et la N - hexyl - N,N',N' - tris (imidazol - 2ylméthyl) - éthylène diamine.

25

10. composition blanchissante selon la revendication 1, dans laquelle le ligand de formule générale (L) est présent sous la forme d'un complexe de formule générale (A) :

30



dans laquelle :

35

M représente le fer à l'état d'oxydation II, III, IV ou V, le manganèse à l'état d'oxydation II, III, IV, VI ou VII, le cuivre à l'état d'oxydation I, II ou III ou le cobalt à l'état d'oxydation I, II ou III ;

X représente une espèce de coordination ;

n représente zéro ou un nombre entier dans la plage de 0 à 3 ;

z représente la charge du complexe et est un nombre entier qui peut être positif, zéro ou négatif ;

Y représente un ion antagoniste dont le type dépend de la charge du complexe ;

40

$$q = z/[\text{charge de Y}]$$

45

11. Composition blanchissante selon l'une quelconque des revendications précédentes, dans laquelle la composition comprend un matériau tensioactif et un adjuvant de détergence.

50

12. Composition blanchissante selon l'une quelconque des revendications précédentes, dans laquelle dans une solution aqueuse, au moins 10 %, de préférence au moins 50 % et optimalement au moins 90 % de tout blanchiment d'un substrat est effectué par de l'oxygène provenant de l'air.

55

13. Composition blanchissante selon l'une quelconque des revendications précédentes, comprenant en outre un séquestrant et dans laquelle; dans une solution aqueuse, au moins 90 % de tout blanchiment du substrat est effectué par une espèce peroxyde qui n'est pas directement dérivée de l'oxygène atmosphérique, de préférence 99 % et le plus préférentiellement 100 %.

14. Procédé de blanchiment d'un substrat comprenant l'étape consistant à appliquer au substrat, dans un milieu aqueux, une composition blanchissante selon l'une quelconque des revendications précédentes.

## EP 1 358 309 B1

15. Ligand choisi dans le groupe constitué de la N - méthyl - N,N',N' - tris (imidazol-2ylméthyl) - éthylène diamine ; la N - éthyl - N,N',N' - tris (imidazol-2ylméthyl) - éthylène diamine ; la N,N' - diméthyl - N,N' - bis (imidazol - 2 - ylméthyl) - éthylène diamine ; la N - (1 - propan - 2 - ol) - N,N',N' - tris (imidazol - 2ylméthyl) - éthylène diamine ; la N - (1 - propan - 2 - ol) - N,N',N' - tris (1 - méthyl - imidazol - 2ylméthyl) - éthylène diamine ; la N,N - diéthyl - N', N'', N'' - tris (5 - méthyl - imidazol - 4ylméthyl) - diéthylène triamine ; la N - (3 - propan - 1 - ol) - N,N',N' - tris (1 - méthyl - imidazol - 2ylméthyl) - éthylène diamine ; et la N - hexyl - N,N',N' - tris (imidazol - 2ylméthyl) - éthylène diamine.

5

10

15

20

25

30

35

40

45

50

55