

1 **The Potential Nutrition-, Physical- and Health-Related Benefits of Cow's Milk for Primary-**
2 **School Aged Children**

3 Penny Rumbold^{1*}, Nicola McCulloch¹, Ruth Boldon¹, Crystal Haskell-Ramsay², Lewis James³,
4 Emma Stevenson⁴, Benjamin Green^{1,5}.

5

6 ¹Department of Sport, Exercise and Rehabilitation, Faculty of Health and Life Sciences, Northumbria
7 University, UK.

8 ²Department of Psychology, Faculty of Health and Life Sciences, Northumbria University, UK.

9 ³National Centre for Sport and Exercise Medicine East Midlands, School of Sport, Exercise and
10 Health Sciences, National Centre of Sport and Exercise Medicine, Loughborough University, UK.

11 ⁴Human Nutrition Research Centre, Population Health Sciences Institute, Faculty of Medical
12 Sciences, Newcastle University, UK.

13 ⁵Danone Specialised Nutrition, UK.

14

15 ***Corresponding Author:** Penny Rumbold. Department of Sport, Exercise and Rehabilitation,
16 Faculty of Health and Life Sciences, Northumbria University, UK. E-mail:
17 Penny.rumbold@northumbria.ac.uk

18

19

20 **Short title:** Benefits of Milk for Primary-School Children

21

22 **Keywords:** Milk; Dairy; Children; Nutrition; Health; Paediatrics

23

24

25

26

27

28 **Abstract**

29 Cow's milk is a naturally nutrient-dense foodstuff. A significant source of many essential nutrients,
30 its inclusion as a component of a healthy balanced diet has been long recommended. Beyond milk's
31 nutritional value, an increasing body of evidence illustrates cow's milk may confer numerous
32 benefits related to health. Evidence from adult populations suggests that cow's milk may have a role
33 in overall dietary quality, appetite control, hydration and cognitive function. Although evidence is
34 limited compared to the adult literature, these benefits may be echoed in recent paediatric studies.
35 This article, therefore, reviews the scientific literature to provide an evidence-based evaluation of
36 the associated health benefits of cow's milk consumption in primary-school aged children (4-11
37 years). We focus on seven key areas related to nutrition and health comprising nutritional status,
38 hydration, dental and bone health, physical stature, cognitive function, and appetite control. The
39 evidence consistently demonstrates cow's milk (plain and flavoured) improves nutritional status in
40 primary-school aged children. With some confidence, cow's milk also appears beneficial for
41 hydration, dental and bone health and beneficial to neutral concerning physical stature and appetite.
42 Due to conflicting studies, reaching a conclusion has proven difficult concerning cow's milk and
43 cognitive function therefore a level of caution should be exercised when interpreting these results.
44 All areas, however, would benefit from further robust investigation, especially in free-living school
45 settings, to verify conclusions. Nonetheless, when the nutritional-, physical- and health-related
46 impact of cow's milk avoidance is considered, the evidence highlights the importance of increasing
47 cow's milk consumption.

48
49
50
51
52
53
54
55
56
57
58
59
60
61

62 **Introduction**

63 Commonly cited as nature's perfect food⁽¹⁾, milk is the foundation of life for all mammalian
64 neonates^(2,3). Representing a complex and unique liquid, milk (which is produced by the mammary
65 glands⁽⁴⁾) is the only foodstuff designed by nature to serve as a complete food, at least during the
66 **pre**-weaning period. In this sense, milk contains numerous biological constituents and an assortment
67 of nutrients necessary for immunological protection and initial growth^(2,5). Shortly after weaning,
68 most mammals stop consuming milk⁽⁶⁾. With domestication and milking of various animals,
69 however, some humans continue to drink milk throughout life⁽⁷⁾, mostly cow's milk. The value of
70 cow's milk in the human diet has been heavily debated for many years⁽⁸⁾ and has been portrayed by
71 some as 'white poison', a health hazard and promoter of Western chronic diseases⁽⁹⁾. This position
72 is based on numerous hypotheses including that cow's milk contains blood, pus, hormones and
73 antibiotics, causes acne and cancer and has a high cholesterol and fat/saturated fat content
74 collectively contributing to cardiovascular disease risk, weight gain and obesity. Nevertheless, such
75 perceptions of cow's milk being harmful to health are not supported by evidence. Conversely, cow's
76 milk has often been signified as the white elixir⁽⁶⁾.

77 Childhood is a key stage for the development of healthy eating patterns⁽¹⁰⁾, and the school
78 environment provides a valuable setting to develop such behaviours. This is particularly true
79 considering children spend much of their time at school. Indeed, dietary habits shaped throughout
80 the childhood years might carry forth and track into adulthood. Between the ages of 4-11 years
81 (primary-school age), children grow and mature at a rapid rate preceding the onset of puberty⁽¹¹⁾.
82 Childhood is therefore a critical transition period preceding adolescence, characterised by growing
83 independence and marked physical development. Good nutrition for the childhood period is
84 therefore particularly important⁽¹²⁾. Not only may good nutrition support proper growth and
85 maturation, but it may therefore act as a base for immediate and lasting health, well-being and
86 disease prevention. This is best achieved by consuming a balanced and varied diet that provides all
87 the nutrients needed. The inclusion of cow's milk as a staple component of a healthy balanced diet
88 has been long recognised and is central in most public dietary recommendations⁽¹³⁾. Milk is naturally
89 nutrient-dense and is a significant source of many essential macro and micronutrients. Indeed, cow's
90 milk may confer nutrition-, physical- and health-related benefits beyond that of helping children to
91 simply meet nutrient targets. Based on the available literature, it appears that many of these effects
92 are a product of milk's nutritional composition. It therefore seems relevant to provide readers with
93 a brief description of the nutritional composition of cow's milk.

94
95

Nutritional Composition of Cow's Milk and Patterns of Consumption

The nutritional composition of cow's milk is influenced by factors including genes (species and breed), physical state (age and stage of lactation) and environment (available nutrition and climatic conditions)⁽¹⁴⁾. The composition of whole cow's milk is approximately 87 % water and 13 % solids⁽¹⁾. The percentage split of water and solids of cow's milk, as well as **energy**, is primarily determined by the amount of fat⁽¹⁵⁾ but is also influenced by added sugars and sweeteners. Cow's milk is approximately 4.9 % carbohydrate. The primary carbohydrate portion of milk is lactose, a disaccharide comprising glucose and galactose. The lipid component of cow's milk and other milk-based dairy foods contribute numerous properties including the provision of fat-soluble micro-nutrients and essential fatty acids, as well as influencing flavour, texture and appearance⁽¹⁵⁾. Of the lipid content within cow's milk, roughly 64 % are saturated fatty acids, with a considerable amount (~26 %) from monounsaturated fatty acids and a small contribution from trans and polyunsaturated fatty acids (both ~3 %)⁽¹⁶⁾. Additionally, cow's milk provides high-quality proteins, namely casein and whey. Casein and whey constitute approximately 82 % and 18 % of the total protein found in cow's milk and provide an abundance of essential amino acids⁽¹⁷⁾. Aside from the macronutrient content, cow's milk contains essential micronutrients that contribute to dietary quality and overall nutritional status. **As shown in Table 1**, with the exception of vitamin C (which is broken down during pasteurisation) **and vitamin D (unless fortified)**, cow's milk is a good source of all vitamins⁽¹⁸⁾. Calcium and **phosphorus**, crucial to healthy growth and maturation, as well as other biological processes, are the most prominent minerals present in cow's milk. Cow's milk also makes significant contributions to intakes of other major minerals⁽¹⁹⁾. **In this sense, cow's milk makes a substantial contribution to, and is the main dietary source of, calcium (26 %), iodine (37 %), riboflavin (25 %), magnesium (10 %) and potassium (14 %) in the diets of primary-school aged children⁽²⁰⁾. While volume specific data are limited, in the UK, the National Diet and Nutrition Survey (NDNS) provides a nationally representative assessment concerning dietary habits of individuals, aged 1.5 y and older, living in private households and remains the only surveillance programme to do so. Temporal and age-related trends of cow's milk to average daily total energy, macro and micronutrient intake from 2009/10 through to 2015-16 are presented in Table 2 for children (4-10 years) and adolescents (11-18 years). Despite the clear value of cow's milk in the everyday diet it is clear intakes steadily decline as children age. This trend is not only restricted to the UK, it is also true in the US^(21, 22), Australia⁽²³⁾ other European countries⁽²⁴⁾. This is of great concern among children, especially as dietary habits may track into adulthood, as milk avoidance may have detrimental implications over time, leaving populations vulnerable to micronutrient deficiencies (e.g. calcium, iodine and riboflavin deficiencies to name a few⁽²⁵⁾) and lasting nutrition-**

130 and health-related complications (e.g. cardiovascular disease⁽²⁶⁾, metabolic syndrome⁽²⁷⁾,
131 hypertension^(27, 28), poor weight management and bone health^(25, 29)).

132

133 **Scope and Methodology of the Narrative Review**

134 Evidence from adult populations suggests that adequate cow's milk consumption is associated with
135 a reduced risk of cardiovascular disease⁽³⁰⁾, metabolic syndrome^(28, 31) and obesity⁽³²⁻³⁴⁾. Emerging
136 data also suggests that cow's milk may have a role in overall dietary quality, appetite control,
137 hydration and cognitive function. Although the evidence to date is limited compared to the adult
138 literature, these benefits appear echoed in recent paediatric studies. There is a need to review the
139 literature to assess whether there is sufficient evidence of a beneficial or detrimental effect of cow's
140 milk in the diet and health of children. The aim of this narrative review is therefore to summarise
141 and appraise the scientific literature to form an accurate, evidence-based evaluation of the associated
142 nutrition-, physical- and health-related benefits of cow's milk consumption in primary-school aged
143 children (4-11 years). Based on recent publications, it is not the intention of this narrative review to
144 focus on body weight and body composition as with some confidence it is known cow's milk
145 consumption is inversely (or not) associated with body weight and body composition in children and
146 adolescents⁽³⁵⁾. Where possible, this review focuses solely on literature in children aged 4-11 years,
147 however, where there is no applicable literature, data from general child studies will be reviewed
148 (and possibly some adult studies). This is particularly true for mechanistic considerations as much
149 of this evidence comes from adult studies, therefore any adult-specific data should be interpreted
150 with some caution as the observations cited may not always be replicated in children. The narrative
151 review will also highlight current knowledge gaps in this field and suggest directions for future
152 research.

153 To identify research articles to facilitate this narrative review, PubMed (US National Library
154 of Medicine National Institutes of Health) and Web of Science (Thomson Reuters, UK), were
155 searched using various combinations of keywords relevant to the scope of the review up to August
156 2020. Search terms included: humans, child, milk, flavoured milk, animal milk, cow milk, infant,
157 adolescent, preschool, primary-school, paediatric, diet, energy intake, nutritional status, diet quality,
158 obesity, school-age children, food habits, dairy products, milk beverages, height, stature,
159 anthropometry, calcium, health association, health benefits, dental health, caries, decay, bone health,
160 appetite, cognition, memory, mental performance, school milk, insulin, glucose, hydration, snack.
161 Studies were considered eligible for inclusion in the review if (i) they were conducted in children 4-
162 11 years; (ii) participants received cow's milk, and not supplemental forms of dairy minerals (unless
163 a milk group was included); (iii) the study had been peer-reviewed. Results from studies that grouped

164 data over a wider age-range were also considered where this was deemed appropriate. This was
165 especially true for prospective and intervention studies, yet the mean age of participants had to fall
166 between the stipulated age range. Studies were excluded if (i) human participants were not used; (ii)
167 participants lay directly outside of the stipulated age-range; (iii) studies used supplements only (iv)
168 studies used different dairy foods (i.e. yogurt, cheese, dairy desserts) (v) the milk consumed included
169 different sources of animal milk; (vi) data reporting was poor and/or not published in English.

170 In this narrative review we have summarised the evidence base, which we identified by related
171 benefits; nutrition-, physical- and health-related and provide a summary for the studies identified.
172 As this is not a systematic review, we did not perform a quality assessment of each study (based on
173 design and implementation) or conduct a meta-analysis by study design but summarise the findings
174 for each study and sub-section in Tables 3 (nutrition-related benefits), 4 (physical-related benefits)
175 and 5 (health-related benefits). Symbols ↑ and ↓ indicate a statistically significant positive and
176 negative effect/relationship, respectively, between cow's milk and related subsections, with ↔
177 indicating no statistically significant effect/relationship or a neutral effect. This approach has
178 successfully been used in a recent narrative review in this journal⁽³⁵⁾, and therefore direct readers to
179 this paper for justifications and interpretations for this approach. In brief, however, a tabulated
180 summary allowed us to draw overall conclusions for this narrative review, although we did not
181 undertake a detailed assessment of quality of individual studies.

182

183 **Nutrition-Related Benefits of Cow's Milk**

184 *Nutritional Status*

185 The importance of regular cow's milk consumption in children has been recognized for over 40
186 years⁽³⁶⁾. As described, cow's milk is a naturally nutrient-dense foodstuff, providing a rich source of
187 many essential nutrients. While the potential of cow's milk to improve nutritional status may be
188 unsurprising, the effect of this in primary-school aged children has only been investigated by
189 numerous observational studies⁽³⁷⁻⁴²⁾ and three intervention-based study⁽⁴³⁻⁴⁵⁾ (see Table 3). All
190 reported improved nutritional status with increased cow's milk consumption, yet, due to the study
191 designs utilised, findings must be interpreted with caution. In the few intervention-based studies⁽⁴³⁻
192 ⁴⁵⁾, which ranged from 16 weeks to 6 months, primary school children were provided with flavoured
193 milk or no drink (control)⁽⁴⁴⁾, 200 mL⁽⁴³⁾ and 250 mL⁽⁴⁵⁾ of multi-micronutrient fortified milk or
194 unfortified milk to establish whether unfortified and fortified cow's milk influences micronutrient
195 status in primary-school aged children. In those consuming fortified versus unfortified cow's milk
196 significant increases in blood riboflavin and vitamin B₁₂ were observed, with all other analytes
197 (selenium, ferritin, and vitamin D) comparable between drinks⁽⁴³⁾. Additionally, significant

198 increases in serum ferritin and zinc increased have been observed after 6 months with both fortified
199 versus unfortified cow's milk compared to a control drink⁽⁴⁵⁾. With respect to nutritional status,
200 dietary patterns characterised by high cow's milk consumption (both plain and flavoured milk)
201 resulted in greater intakes of energy, protein, phosphorus, magnesium, calcium, potassium, vitamin
202 A, zinc, riboflavin (vitamin B₂), and niacin compared to children who seldom consumed cow's milk.
203 Notably, diets high in cow's milk may also limit intakes of foods and beverages high in fat and
204 sugar. In studies of Dutch⁽³⁷⁾ and American primary-school aged children⁽⁴⁶⁾, for example, cow's
205 milk intake was inversely related with the intake of sugar-sweetened beverages. In these studies,
206 children with low cow's milk intakes had lower protein, fibre, calcium, magnesium, potassium and
207 phosphorus intakes. Low cow's milk consumption has significant implications for intakes of several
208 key nutrients that are of importance in childhood. Based on the nutritional contribution to dietary
209 intakes, it is important to note that children who drink cow's milk regularly are more likely to meet
210 dietary recommendations for many nutrients, and thus have a better nutritional status⁽¹³⁾. It could
211 therefore be argued that milk intake might be a marker for healthier eating habits⁽³⁷⁾.

212 With the above in mind, in the UK, it should be noted that dietary intakes of vitamin D in
213 primary-school aged children are low. Considering UK cows' milk is generally not a good source of
214 vitamin D because it is not fortified, as it is in other countries, this narrative review may provide
215 justification for UK policy makers to reconsider widespread fortification. There is evidence (albeit
216 from a theoretical modelling perspective) from Northern Ireland that cows' milk can be used as a
217 successful vehicle for vitamin D fortification⁽⁴⁷⁾. Fortification of cows' milks with 1 µg, 1.5 µg and
218 2.0 µg/100g, theoretically increased median vitamin D intakes from 2.0 µg/day to 4.2 µg, 5.1 µg and
219 5.9 µg/day, respectively and may therefore provide strong evidence for the efficacy of widespread
220 fortification. This modelling appears to translate to human studies. In support of this, a recent review
221 comprising of 20 studies showed positive associations between the consumption of vitamin D
222 fortified milk and 25(OH)D status in different population groups⁽⁴⁸⁾. Furthermore, in Finland,
223 Canada, United States who exercise a national vitamin D fortification policy (covering various fluid
224 milk products), milk products contributed 28–63 % to vitamin D intake, while in countries without
225 a fortification policy, or when the fortification covered only some dairy products (Sweden, Norway),
226 the contribution was much lower or negligible⁽⁴⁸⁾. Based on the above, widespread fortification of
227 milk seems helpful to help bolster vitamin D intakes and could be adopted by the UK to increase
228 vitamin D intakes in primary-school children.

229 Concerns about added sugars in foods and beverages has increased recently. Accordingly,
230 many schools have limited access to foods and beverages high in added sugars, included flavoured
231 cow's milk. In some studies, removal of flavoured cow's milk from the school environment reduces

232 **energy** and sugar intake, but negatively impacts essential nutrient intake and even reduced the
233 overall intake of non-flavoured cow's milk^(49, 50). In this sense, total milk consumption (both plain
234 and flavoured) decreased by 12.3 %⁽⁵⁰⁾. Contrariwise, children who drink more flavoured cow's milk
235 generally have higher non-flavoured milk intakes and display nutritional intakes similar to children
236 who only consume non-flavoured milk⁽⁵¹⁾. While greater cow's milk consumption (both plain and
237 flavoured) is generally associated with higher daily energy intake with some confidence we know
238 this does not impact body mass or composition⁽³⁵⁾. If daily energy intake is of concern for children
239 predisposed to overweight and obesity, results of a recent study suggest that replacement of whole
240 cow's milk or semi-skimmed cow's milk with skimmed cow's milk may help reduce total energy
241 intake, without impacting on nutrient provision⁽⁵²⁾.

242 When taken together, these results suggest that cow's milk consumption (both plain and
243 flavoured) might serve as a useful strategy to boost nutritional status in primary-school age children
244 and may act as a surrogate marker of diet quality. There may be a need for widespread vitamin D
245 fortification of cow's milk, however, and based on the above seems justified and should be reviewed
246 by UK policy makers. Cow's milk is a readily available, accessible, and affordable means of
247 providing valuable essential nutrients to the diets of primary-school children. The nutritional
248 implications of cow's milk provision and/or removal in the school environment must be considered
249 and is especially relevant in children suffering with nutritional inadequacies. The data are, however,
250 primarily limited to observational investigations and require verification in more controlled
251 intervention-based studies covering differing populations.

252

253 *Hydration*

254 Whole cow's milk is approximately 87 % water⁽¹⁾ and may therefore be a beneficial choice for
255 hydration. For children, maintenance of euhydration is important for good health⁽⁵³⁾ but may also
256 increase concentration and mental performance (cognitive function)⁽⁵⁴⁾ while helping reduce
257 instances of headaches, constipation and other disorders⁽⁵³⁾. This is important as children are at a
258 greater risk of dehydration compared to adults, having relatively greater fluid losses at rest and
259 during exercise and being less able to recognise thirst⁽⁵⁴⁾. Given that water is continually being lost
260 from the body water pool, it is important to constantly replenish fluid losses to prevent dehydration.
261 Establishing beverages that encourage longer-term fluid retention and maintenance of euhydration
262 has real clinical and practical implications⁽⁵⁵⁾. This is particularly true in situations where free access
263 to fluids is limited or when frequent breaks for urination are not desirable⁽⁵⁵⁾, such as in in the school
264 setting where many children arrive at school in an already hypo-hydrated state⁽⁵⁶⁾. There is little
265 research concerning the impact of cow's milk on hydration in primary-school aged children (see

266 Table 3). To date, three studies have been conducted⁽⁵⁷⁻⁵⁹⁾. Two of these were exercise-based
267 intervention studies conducted by the same research group in Canada^(58, 59), and the remaining study
268 was cross-sectional⁽⁵⁷⁾. Nonetheless, all reported improved hydration status with cow's milk in
269 comparison to water or alternative beverages. While these findings show promise, additional
270 controlled-intervention studies should be performed in different settings to examine repeatability of
271 these effects.

272 In the two exercise-based intervention studies, researchers examined the influence of post-
273 exercise cow's milk consumption on rehydration in 7-11-year-old⁽⁵⁹⁾ and 10-12-year-old children⁽⁵⁸⁾.
274 The rehydration potential of cow's milk (or milk-protein) was compared to water or a carbohydrate-
275 electrolyte drink, zero or a low cow's milk protein beverage and were given at 100 % and 150 % of
276 the children's body mass losses following exercise, respectively. In both studies, children exercised
277 in the heat (~34.5 °C) and consumed the test beverages immediately following exercise. The children
278 were subsequently observed for a period of 2 hours⁽⁵⁹⁾ and 4 hours⁽⁵⁸⁾, respectively, during which
279 measures of hydration status were collected (urine output and fluid balance). Findings over 2 hours
280 showed cow's milk was more effective than both water and the carbohydrate-electrolyte drink at
281 replacing fluid loss during exercise⁽⁵⁹⁾. This was similar over 4 hours. The authors suggested that the
282 protein in cow's milk may be a factor responsible for the rehydrating properties⁽⁵⁸⁾.

283 Montenegro-Bethancourt and colleagues⁽⁵⁷⁾ conducted a cross-sectional study designed
284 initially to establish the contribution of fruit and vegetable intake on hydration status. Children (4-
285 10-year-old) recorded all food and beverages consumed over 3 days using a weighed food diary
286 alongside 24 hour urine samples. Regular intake of fruit and vegetables made a substantial
287 contribution to hydration status, but notably, cow's milk consumption was also a strong dietary
288 predictor of hydration status. In particular, cow's milk increased free water reserve by 25 mL in boys
289 and 33 mL in girls per 100 g of intake. These findings corroborate studies in adult populations⁽⁶⁰⁻⁶²⁾.

290 Based solely from adult studies, there are a number of potential mechanisms explaining the
291 greater fluid retention (and thus hydration potential) with cow's milk⁽⁶³⁾. Firstly, milk contains
292 modest amounts of sodium (~20 mmol/L, similar to most commercial sports drinks) and large
293 amounts of potassium (~40 mmol/L). Sodium, as the main cation in the extracellular fluid plays a
294 major role in fluid retention⁽⁶⁴⁾, whilst potassium may exert some beneficial effects⁽⁶⁵⁾, although this
295 is not a consistent finding. Secondly, the protein content of cow's milk may help facilitate greater
296 fluid retention^(66, 67). While water and alternative beverages meet the basic intentions of rehydration,
297 they do not offer the abundance of nutrients present in cow's milk that will also aid in normal growth
298 and maturation. Nevertheless, it should be considered that while these findings suggest that cow's
299 milk helps improve the hydration status of children, there remains room for further studies to clarify

300 the role of cow's milk in hydration, especially in a free-living school setting. While the intervention-
301 based studies show promise, many primary-school children do not exercise in such conditions.

302

303 **Physical-Related Benefits of Cow's Milk**

304 *Dental Health*

305 Milk contains multiple nutrients that may offer anticariogenic properties, protecting against the
306 development of dental caries, and thus supporting dental health in children^(68, 69). The nutrients
307 principally believed to play a role in dental health include calcium, phosphorus and protein⁽⁶⁸⁻⁷⁰⁾. As
308 reported earlier, cow's milk intake has been shown to be inversely related with the intake of sugar-
309 sweetened beverages^(37, 46). Although speculative, one could argue increased cow's milk intake
310 might indirectly improve dental health. Calcium is required for bone and teeth formation, whereas
311 phosphorus ions works alongside calcium to maintain tooth strength⁽⁷⁰⁾. Cow's milk proteins
312 (particularly α_{s1} -, α_{s2} - and β -casein) may act to prevent tooth enamel erosion and demineralisation
313 of the tooth surface by producing casein phosphopeptides⁽⁷¹⁾. Based on the available evidence,
314 inverse associations between cow's milk intake and the incidence of tooth decay have frequently
315 been reported⁽⁷²⁻⁷⁵⁾ (see Table 4). The evidence, however, has been derived from cross-sectional
316 research, so causal conclusions cannot to be justified and caution must be exercised.

317 The findings of the cited studies were generated from recall methods which have obvious
318 shortcomings. Nevertheless, all studies (following visual examinations accompanied with food
319 frequency questionnaires) reported inverse associations with cow's milk and the development of
320 dental caries. Interestingly, though sugar is suggested to possess acidogenic and cariogenic
321 potential⁽⁷⁶⁾, one study⁽⁷⁴⁾ reported the association between cow's milk consumption and protection
322 against of dental caries was stronger for children with diets high in sucrose. This might suggest that
323 cow's milk offers protection against the harmful effects of sugar, though this is speculative. To this
324 end, cow's milk intake during primary-school years has been reported as a predictor of incidences
325 of caries later in childhood⁽⁷²⁾. In this sense, greater cow's milk intake is inversely associated with
326 indices of dental caries. Collectively, these studies support the suggestion that dietary habits
327 established and maintained during primary-school years could have longer-term effects on health
328 outcomes.

329 Although no controlled trials have been conducted in primary-school children, the available
330 literature appears to suggest that cow's milk could help reduce the incidence of dental caries and
331 contribute to dental health in children. To reduce the occurrence of tooth decay, it is recommended
332 primary-school children limit their consumption of sugary beverages (especially when not consumed
333 with a meal) and increase consumption of cow's milk. The exact mechanism by which cow's milk

334 reduces the incidence of dental caries remains uncertain, though calcium, phosphate and casein
335 phosphopeptides may all play a role⁽⁶⁸⁻⁷⁰⁾. Casein phosphopeptides are phosphorylated casein-
336 derived peptides produced by tryptic digestion of casein in the duodenum⁽⁷¹⁾. The anticariogenic
337 activity of casein phosphopeptides is due to their ability to stabilise high levels of amorphous
338 calcium phosphate on tooth surface, preventing demineralisation and enhancing remineralisation of
339 enamel caries⁽⁷⁷⁾. In addition, milk fat could be adsorbed onto the enamel surface and may have a
340 protective role; and thirdly, milk enzymes may have a role in reducing the growth of acidogenic
341 plaque bacteria⁽⁷⁸⁾. Where prior observational research provides a solid foundation, any future work
342 should seek to implement robust randomised clinical trials (RCT) to confirm any causal relationships
343 between cow's milk consumption and dental health in primary-school aged children.

344

345 *Bone Health and Physical Stature*

346 Childhood is a critical time for bone growth and lasting bone health^(79, 80) and the nutritional
347 composition of cow's milk has evolved to stimulate and support this^(2, 5). The scientific opinion that
348 regular cow's milk consumption is associated with greater physical stature has a long history, dating
349 back to the 1920s^(81, 82). Cow's milk contains multiple nutrients that may support childhood
350 growth⁽⁸⁰⁾ and lasting bone health⁽⁷⁹⁾. The beneficial effects of cow's milk consumption on bone
351 health in children may include increased bone mineral content (BMC) and bone mineral density
352 (BMD), characteristics necessary for the prevention of bone-related diseases later in life
353 (osteoporosis). Evidence of these benefits, however, is equivocal showing a beneficial to neutral
354 effect of cow's milk on these constructs^(25, 29, 83-88). In this sense, of the cited studies, two reported
355 greater total body BMC^(85, 88), while one reported no effect. One study reported no effect on total
356 body BMD⁽⁸⁷⁾, yet three studies reported increased regional BMC^(29, 83, 86) and another two studies
357 reported increased regional BMD^(25, 84) (see Table 4). **The mixed findings reported throughout these**
358 **studies may be due to a lack of consistency in methodological approaches, length of study, location**
359 **and measures of bone health, all of which confound comparisons and prevent a clear conclusion.**
360 **Furthermore, age and sex differences must be considered, especially as puberty and bone**
361 **mineralisation typically occurs earlier in girls compared to boys⁽⁸⁹⁾.**

362 In two recent studies^(85, 88), school milk interventions increased total body and regional
363 (forearm) BMC and BMD compared to children who seldom drank cow's milk. In these studies, the
364 beneficial effects of cow's milk on bone health were observed in $n = 435-757$ children (mean age
365 11 years) following daily school milk intake for 1-2 years.

366 On a comparative basis with other animal sources, whole cow's milk is the richest source of
367 calcium and represents the **biggest contributor** of dietary calcium during childhood⁽⁹⁰⁾. In addition,

368 considering beef meat and eggs for example, cow's milk is the cheapest source of protein, calcium,
369 phosphorus, and vitamin D⁽⁹¹⁾. During childhood, the beneficial effects of cow's milk on bone health
370 in children are commonly attributed to calcium⁽⁹²⁾. However, many other nutrients including
371 phosphorus and protein are needed for normal growth and lasting bone health⁽⁹³⁾. In 5-11-year-old
372 children (n = 99), for example, calcium supplementation for 10 months did not influence total body
373 or regional BMC⁽⁸⁷⁾. In contrast, in an earlier study where cow's milk (and dairy) was supplemented
374 daily for 1 year (distributed to deliver 1200 mg calcium daily), lumbar (lower back) BMD increased
375 compared to children who maintained their usual eating habits⁽⁸⁴⁾. This may illustrate that calcium
376 and other nutrients work together for bone growth, and thus lasting bone health⁽⁹³⁾. It is important to
377 note, however, that children in the calcium supplementation study were already consuming near
378 daily recommended amounts, which may illustrate intakes exceeding calcium recommendations
379 (from either supplemental calcium or cow's milk) offers no further benefit to bone health in children.

380 In several studies, it has been observed that children who avoid consuming cow's milk
381 characteristically exhibit low calcium intakes, short statures, increased fatness and lower BMC (and
382 thus exhibit reduced bone health) compared to their cow's milk-drinking counterparts^(25, 86). In
383 children (mean age = 8 years) who previously avoided cow's milk, the introduction of cow's milk
384 to the diet increased not only habitual cow's milk consumption but also increased total body
385 BMC⁽²⁹⁾. This may suggest it is never too late for children to introduce cow's milk into their diet for
386 bone health benefits.

387 From a stature perspective, available data appear to suggest that cow's milk consumption
388 almost certainly has a positive effect on growth in children. To date, fourteen studies in primary-
389 school aged children have evaluated this aspect of cow's milk consumption^(25, 29, 36, 44, 81-83, 85, 94-99).
390 Seven were intervention-based, four were observational, and the remaining three were prospective
391 designs. Based on these studies, the evidence suggests that cow's milk intake positively influences
392 physical stature in primary-school aged children. All six intervention studies showed increased
393 stature with increased cow's milk (one study included milk calcium). In addition, all prospective
394 studies reported increased stature with increased cow's milk. With regards to the cross-sectional
395 studies, two illustrated cow's milk avoiders displayed stunted growth compared to cow's milk
396 drinkers while the remaining study showed increased adult stature with increased cow's milk
397 consumption in childhood. Although trends are consistent across studies, there remains a need for
398 evidence from robust controlled-intervention trials in primary-school aged children to verify
399 causality.

400 Beneficial effects on physical stature were observed with both whole and reduced fat cow's
401 milk, distributed at a range of 190 mL – 568 mL daily. In several of these studies, it was reported

402 that childhood cow's milk intake was associated with higher skeletal development (BMD of the hip
403 and the forearm), bone growth and periosteal bone expansion. These were likely established earlier
404 during growth periods and maintained into late adolescence and young adulthood⁽⁹⁶⁻⁹⁹⁾, supporting
405 the notion that dietary habits established and maintained during the primary- and secondary-school
406 years may not only induce short-term effects but offer lasting benefits. Indeed, in children with
407 prolonged cow's milk avoidance, stunted growth and physical stature is observed compared to
408 children who habitually consume cow's milk, and this is maintained into adulthood^(25, 29, 86). During
409 a pubertal growth spurt, about 37 % of the entire skeletal mass is accumulated⁽¹⁰⁰⁾. Therefore,
410 inadequate calcium intake during this period may compromise volumetric bone density and overall
411 stature attained. Notably, in a study that explored both cow's milk and supplemental calcium, those
412 children in the cow's milk (and dairy) group were 3 cm taller (166 cm) compared to the supplemental
413 calcium (163 cm) and placebo (163 cm) group⁽⁹⁶⁾. This may indicate that cow's milk has more of a
414 beneficial effect on physical stature and growth than single cow's milk constituents (i.e. calcium).

415 The precise mechanisms or nutrients in cow's milk responsible for stimulating growth and
416 lasting bone health are not yet clear. Evidence suggests that maintaining adequate calcium intake
417 during childhood might be advantageous for the attainment of peak bone mass, which may be crucial
418 in reducing the risk of bone-related diseases later in life⁽¹⁰¹⁾. Interestingly, it appears that whole foods
419 may offer greater benefits than the equivalent amount of calcium in supplemental form. It has also
420 been suggested that the growth-stimulating effect of cow's milk is likely attributed to hormonal
421 effects that can be influenced by ingested cow's milk proteins (**predominantly whey protein and the**
422 **release of amino acids during digestion but also casein**), micronutrients and also energy⁽¹⁰²⁻¹⁰⁴⁾.
423 Observations from child studies show these nutrients stimulate the secretion of insulin-like growth
424 factor-1 and insulin, both of which are anabolic hormones that play an essential role in the regulation
425 of growth and accrual of bone mass during childhood⁽¹⁰²⁻¹⁰⁴⁾ though there is some controversy that
426 cow's milk intake upregulates insulin and insulin-like growth factor-1 signalling and thus promotes
427 chronic diseases **such cancer (prostate, breast and colorectal) and cardiovascular disease⁽⁹⁾, though**
428 **these perceptions are hypothetical at present and not supported by evidence.**

429 **Nonetheless, when taken together,** it appears that cow's milk consumption promotes both
430 health and may increase physical stature in primary-school aged children. This suggests that cow's
431 milk consumption during childhood might be important to ensure full growth potential is achieved.
432 While there is some suggestion about the improved bone health with increased cow's milk
433 consumption and the mechanisms responsible for the growth-stimulating properties of cow's milk,
434 more intervention studies are needed to elucidate the components responsible for these effects and
435 to prove and/or disprove the chronic diseases hypotheses. This is especially prudent when

436 considering bone health as the methodological approaches previously employed have been diverse
437 in nature.

438

439 **Health-Related Benefits of Cow's Milk**

440 *Appetite Regulation*

441 Appetite comprises numerous regulatory processes associated with the initiation and termination of
442 eating and the selection and amount of food consumed. The regulation of appetite and eating
443 behaviour depend on the detection and integration of signals relaying nutritional status, and their
444 interaction with signals associated with food palatability and gastrointestinal handling, in addition
445 to circadian, social, emotional, habitual and other situational influences⁽¹⁰⁵⁾. Consequently, appetite
446 and the regulation of eating behaviour are complex processes, regulated by homeostatic and non-
447 homeostatic influences⁽¹⁰⁶⁾. Cow's milk contains a host of nutrients that might exert a favourable
448 effect on appetite and eating behaviour⁽¹⁰⁷⁾, yet in primary-school children there is limited evidence
449 concerning the short-term effect of cow's milk on these measures. There are also no data on the
450 moderate- and longer-term effects of daily cow's milk consumption on these outcomes in primary-
451 school children.

452 From the available studies⁽¹⁰⁸⁻¹¹²⁾, cow's milk has principally been given as a mid-morning
453 snack or alongside breakfast, with the volume of cow's milk offered to children ranging from 160–
454 250 mL. Based on these studies, the evidence concerning cow's milk and appetite regulation is
455 inconclusive. Three of these studies found a decrease in energy intake after cow's milk consumption
456 yet three reported no effect. Two studies reported cow's milk consumption reduced subjective
457 appetite, one reported increased subjective appetite compared to a fruit-based snack, while two
458 studies did not measure subjective appetite. Only one study measured hormonal indicators of
459 appetite and reported cow's milk consumption stimulated the secretion of glucagon-like peptide-1.

460 In two studies of 34 overweight and obese boys (mean age = 11 years)^(109,110), when compared
461 with volume and energy matched servings of water or fruit-juice, 240 mL of low-fat cow's milk
462 with breakfast reduced energy intake at an *ad-libitum* lunchtime meal. In another study comprising
463 of 48 obese children (mean age = 11 years), girls reported higher satiety scores 4 hours after drinking
464 whole cow's milk compared to skimmed milk, and low-fat cow's milk significantly reduced appetite
465 compared to apple juice 2 hours after consumption. These differences, however, did not transpire to
466 changes in energy intake at an *ad-libitum* lunchtime meal across all conditions in girls, boys and the
467 group as a whole⁽¹⁰⁸⁾. As mentioned, only one investigation (comprising two experiments) has
468 sought to establish the effect of cow's milk (and other dairy products) on appetite and feeding
469 behaviour in normal weight and overweight/obese children (mean age = 11.5), where subjective

470 appetite and appetite-related analytes were measured⁽¹¹²⁾. In both experiments, preloads (experiment
471 1: 1 % fat (1 g/100g) chocolate cow's milk, 2 % (2 g/100g) fat cow's milk, 1.5 % (1.5 g/100g) fat
472 yogurt drink, fruit punch or a water drink; experiment 2: 2 % (2 g/100g) fat cow's milk or a fruit
473 punch) were provided 60 min preceding and during an *ad libitum* pizza meal. All preloads were
474 matched for volume (250 mL) and energy content (130 kcal, 543.9 kJ; except water in experiment
475 1). The first experiment comprised measures of subjective appetite, whereas the second experiment
476 included measures of subjective appetite together with appetite-related **analytes** (serum glucose,
477 insulin and plasma GLP-1 and peptide YY). Reduced energy intake was observed following
478 chocolate cow's milk and yogurt consumption compared to a water drink in the first experiment.
479 Consistent with a reduction in energy intake, subjective appetite (combined appetite score) was
480 significantly lower following 2 % (2 g/100g) fat cow's milk compared with the yogurt drink only.
481 No additional effects were observed concerning energy intake following the consumption of 2 % (2
482 g/100g) fat cow's milk and fruit punch or on subjective measures of appetite after 1 % fat chocolate
483 cow's milk, 1.5 % (1.5 g/100g) fat yogurt drink, fruit punch or water. Compared with the fruit punch
484 preload, cow's milk consumption resulted in a significantly greater GLP-1 area under the curve.
485 Nonetheless, *ad libitum* energy intake, insulin and glucose AUC were comparable between trials.
486 Considering all aforementioned studies, it is important to consider that in these studies, energy intake
487 was principally assessed via *ad libitum* assessments which may not be reflective of free-living eating
488 behaviour.

489 The mechanism(s) by which cow's milk consumption might influence eating behaviour are
490 not fully understood, but there are several plausible suggestions from the adult literature and
491 constituents of cow's milk that may act synergistically to explain possible actions. In the studies
492 where cow's milk reduced appetite and subsequent eating behaviour, it is probably unsurprising that
493 cow's milk consumption suppressed energy intake at *ad-libitum* assessment meals, considering the
494 macronutrient composition of cow's milk compared to fruit-juice drinks and water. Cow's milk
495 contains considerably more protein than fruit-juice drinks and water. Although it is not a universal
496 finding, it is widely recognised that dietary proteins are more satiating than energetic equivalents of
497 carbohydrate and fat under most conditions, commonly suppressing eating behaviour at the next
498 meal⁽¹¹³⁻¹¹⁵⁾. Consequently, cow's milk proteins (whey and casein, and their products of digestion)
499 may act to potentiate peptides of gastrointestinal, pancreatic and adipose tissue origin, increasing
500 perceptions of satiety and acutely reducing energy intake (anorexigenic behaviours)⁽¹¹³⁾. Moreover,
501 medium-chain triglycerides, conjugated linoleic acid (**CLA**) and lactose may also be implicated in
502 the reduction of energy intake after cow's milk intake⁽¹⁰⁷⁾. Medium-chain triglycerides are absorbed
503 directly into the portal circulation and transported to the liver for rapid oxidation. A combined action

504 of increased energy expenditure, decreased fat deposition and increased satiety may reduce of energy
505 intake via pre-absorptive signals, post-absorptive changes in metabolites and appetite-related
506 analytes⁽¹⁰⁷⁾ and similar appetite-related analyte responses have been observed with lactose⁽¹¹⁶⁾.
507 When considering CLA intake and its potential implications in appetite regulation, cow's milk (and
508 dairy) and cattle meat (cows and lamb) represents the almost exclusive dietary sources⁽¹¹⁷⁾ where
509 the predominant isomer of CLA (accounting for more than 90 % of the total CLA intake) is the *cis*-
510 9, *trans*-11-CLA. It is, however, strongly proposed that other isomers, such as *trans*-10, *cis*-12-CLA
511 may influence body-weight and fat changes⁽¹¹⁸⁾. In agreeance with an earlier narrative review⁽³⁴⁾, it
512 remains unknown whether cow's milk (and dairy), when providing physiological doses of CLA, can
513 elicit any meaningful impact on appetite and body weight regulation in humans. This is especially
514 prudent when considering experimental design, age, sex, energy intake and CLA metabolism of the
515 participants, and the dose and chemical form of the CLA isomer administered as differences may
516 arise solely from these methodological differences⁽¹¹⁹⁾. From a child perspective, potential
517 underlying mechanisms of CLA on appetite regulation are poorly understood though evidence from
518 adult studies suggested that CLA can inhibit fatty acid synthase and stearyl-CoA desaturase-1⁽¹¹⁸⁾,
519 enhance fat oxidation and thermogenesis and reduce lipogenesis and preadipocyte differentiation
520 and proliferation⁽¹²⁰⁾.

521 In summary, evidence suggests that cow's milk may have a unique potential to influence
522 elements of energy balance. Macro and micronutrients and other bioactive constituents might act
523 individually, though probably synergistically, to impart beneficial effects on energy balance and
524 body mass regulation through actions related to appetite, eating behaviour and metabolism.
525 However, there is little mechanistic exploration of cow's milk consumption and appetite regulation
526 from a paediatric perspective. Understanding the relationship between cow's milk consumption and
527 appetite regulation could provide novel nutritional interventions to contribute toward the fight
528 against childhood overweight and obesity⁽¹²¹⁾, whilst bolstering nutritional status and improving
529 elements of cognitive function and hydration. Controlled intervention studies are necessary to
530 determine the best possible timings to administer cow's milk and establish whether consumption
531 delivers these benefits when administered during the school day.

532 Collectively, the effects of cow's milk intake on appetite regulation in primary-school aged
533 children are unclear and could be dependent on BMI. The studies summarised suggest that mid-
534 morning milk consumption influences eating behaviour at the next meal in overweight and obese
535 children, showing that cow's milk could be beneficial for reducing body mass. In lean children, the
536 evidence suggests there is no effect of cow's milk consumption on appetite and eating behaviour,
537 but cow's milk may boost the nutritional quality of the diet. There is much scope for further studies

538 to clarify the role of cow's milk on appetite and eating behaviour, especially in a free-living school
539 environment where methodological approaches are more reflective of habitual eating behaviours. In
540 addition, it will be important to fully distinguish the effects of cow's milk on appetite- and on
541 metabolism-related peptides and subsequent eating behaviour.

542

543 *Cognitive Function*

544 Compared to children with better dietary quality, those with nutritional inadequacies demonstrate
545 decreased attention and academic performance⁽¹²²⁾. Aside from improving nutritional status and
546 dietary quality, emerging evidence illustrates that cow's milk may positively influence cognitive
547 function in primary-school children^(43, 123, 124). Varying in duration from 2 hours to 9 months, studies
548 of an intervention-based nature generally report that cow's milk consumption increases elements of
549 cognitive function, though some of the specific outcome measures demonstrate no effect or in some
550 cases a negative effect of cow's milk. These studies highlight that consideration should be given to
551 potential moderators such as sex and to the use of cow's milk as a non-stigmatised method for
552 providing nutrients through fortification⁽¹²⁵⁾. One non-interventional study considered the adherence
553 of n = 1595 children to Canadian nutrition recommendations in Grade 5 (10-11 years) and in relation
554 to academic achievement in the provincial achievement tests taken approximately one year later⁽¹²⁶⁾.
555 A positive association was identified for boys, with those who met the nutrition recommendations
556 for milk and alternatives (at the time of the study: 3-4 servings per day) scoring 3.45 % better on an
557 average measure of Math and Language Arts tests than those who did not meet the
558 recommendations⁽¹²⁶⁾. An account of the intervention studies follows, but overall, the varying
559 methodological approaches used suggests caution is needed in making firm conclusions about
560 potential links between cow's milk consumption and improved cognitive function.

561 In a study⁽¹²⁴⁾ involving n = 469 boys and girls (mean age = 8 years), evaluating the effects of
562 daily mid-morning cow's milk consumption on physical, mental and school performance,
563 researchers found that a school feeding scheme focusing on daily cow's milk intake had beneficial
564 effects on school performance for girls. In this study, children received a serving of cow's milk (250
565 mL) daily for 12 weeks. Assessments of physical, mental and school performance were conducted
566 prior to and at the end of the 12-week supplementation period. Similarly, in a smaller study⁽¹²³⁾
567 involving 40 children (mean age 11 years), the effects of a carbohydrate drink, a cow's milk drink
568 or a combination of both on subsequent cognitive function (processing speed, memory, attention
569 and perceptual speed) was assessed over a 3 hour period⁽¹²³⁾. Findings showed cow's milk
570 consumption improved short-term memory. Children were able to recall 0.7–0.8 more words
571 compared with 0.5 fewer words after the carbohydrate drink. However, this effect was only observed

572 in girls and not boys^(123, 124). There were slightly more mixed findings from a crossover study in
573 which 84 children (mean age 10 years) were given 237 mL of cow's milk or apple juice⁽¹²⁷⁾. While
574 the beverages were not standardised for temperature, participants were asked to complete
575 computerised tasks of inhibitory control, speeded working memory and sustained attention at
576 baseline and 30-, 90- and 120-min following beverage consumption. Although the significant results
577 following cow's milk compared to juice consumption were, again, only apparent in girls, there was
578 a negative effect (decreased working memory accuracy) alongside the positive one (improved
579 reaction time on the sustained attention task). There were non-significant trends in the opposite
580 direction for boys. No significant effects of the beverages were observed for on-task behaviour
581 during the testing.

582 The mechanisms responsible for the beneficial effects of cow's milk on improved cognitive
583 function are unclear. One suggestion is a sustained blood glucose response following
584 consumption^(128, 129). Findings from Anderson et al.⁽¹²⁷⁾, an adult-based study, suggest that
585 glucoregulation may play a role as participants with higher fasting glucose levels demonstrated
586 faster reaction times on an inhibition task following cow's milk compared to juice. There were
587 however no sex differences in initial glucose elevation or change in glucose levels over time to
588 explain the apparent sex differences in cognition. Such findings suggest a likelihood that factors
589 other than glucoregulation will explain why cow's milk differentially affects the cognition and
590 behaviour of boys and girls. The micronutrient content of cow's milk is another potential mechanism
591 for the observed effects⁽⁴³⁾. Low intakes of vitamin B₁, folate and vitamin B₁₂ affect short-term
592 memory and impair learning, causing low cognitive scores and development in primary-school age
593 children⁽¹³⁰⁾. In addition, low iron and riboflavin intake may adversely affect motor skill
594 development and psychomotor performance⁽¹³¹⁾. All of these micronutrient are heavily present in
595 cow's milk. In one of the longer intervention-based studies available (5 months), Kuriyan and
596 colleagues⁽⁴³⁾ attempted to establish if fortification of cow's milk with multiple micronutrients
597 influenced the mental and physical performance of children (7-10 years) compared to an unfortified
598 cow's milk drink. The children were randomised into groups and were provided with cow's milk (2
599 x 200 mL) 6 days a week for 5 months, with assessments of attention and executive function
600 conducted at baseline and 5 months. The findings showed improved cognitive and physical
601 performance in both groups, illustrating that further fortification of cow's milk provided no
602 additional benefits to cognitive and physical performance but did improve some elements of
603 nutritional status⁽³⁹⁾. Finally, in a double-blind RCT (9 months), the learning potential of 7-9-year-
604 olds (31.9 % with a moderate or severe iodine deficiency at baseline) improved to a greater degree

605 following 200 mL daily of fortified cow's milk (fortified with 45 µg iodine (given as potassium
606 iodide) and other micronutrients) than following nonfortified milk (20.8 µg iodine)⁽¹³²⁾.

607 Some of the components of cow's milk may be detrimental for the cognitive performance of
608 lactase deficient children. In a study involving children of 5-6 years (85 % lactase deficient),
609 information processing efficiency was assessed after 5 days' consumption of 150mL twice daily of
610 conventional cow's milk (containing A1 and A2 β-casein) or cow's milk containing A2 β-casein
611 only⁽¹³³⁾. While post-intervention response times were significantly improved from baseline for both
612 cow's milk products, error rates reduced in the A2 β-casein only condition. Furthermore, consuming
613 conventional cow's milk in the second phase of the crossover study appeared to undo the positive
614 effects on error rate obtained from the A2 β-casein only milk in the first phase, which were
615 maintained through the 9 day washout period.

616 It is prudent to highlight that none of the studies of cognition in primary-school children have
617 compared cow's milk to a control beverage. It is therefore difficult to ascertain whether these studies
618 simply show less detrimental effects of cow's milk than comparator beverages. Furthermore, where
619 no comparators have been employed this could reflect practice effects. Nonetheless, based on the
620 published studies, and bearing in mind the varying methodological approaches employed, it is
621 difficult to ascertain the role of cow's milk in cognitive function for primary-school aged children
622 but it may at the very least be a useful conduit for nutrient fortification. This could be particularly
623 meaningful within the school environment. The intervention studies did not measure academic
624 achievement directly, but their findings have relevant scholastic implications that warrant further
625 investigation using RCTs to establish if increased milk consumption influences academic
626 achievement in primary-school aged children. This is especially so given the identification of a
627 positive association for boys between academic achievement and meeting the recommendations for
628 consumption of milk and alternatives⁽¹²⁶⁾.

629

630 **Future Directions and Conclusions**

631 The aim of this narrative review was to evaluate evidence for the potential nutritional-, physical-
632 and health-related benefits of cow's milk consumption for primary-school aged children (4-11
633 years). Cow's milk consumption (both plain and flavoured) improves nutritional status without
634 adversely impacting on body mass and body composition⁽³⁵⁾. With some confidence, cow's milk also
635 appears beneficial for hydration, dental and bone health and to have a beneficial to neutral effect
636 concerning physical stature and appetite. Due to conflicting studies, reaching a conclusion has
637 proven difficult concerning cow's milk and cognitive function therefore a level of caution should be

638 exercised when interpreting these results. All areas, however, would benefit from further robust
639 investigation, especially in free-living school settings, to verify conclusions. Improving elements of
640 cognitive function, hydration and appetite regulation could have important long-term health and
641 scholastic implications that should certainly be explored further. Indeed, recent research involving
642 adolescent populations illustrates that high intakes of cow's milk are positively associated with
643 academic performance⁽¹³⁴⁾, increased motivation for learning⁽¹³⁴⁾ and impact favourably on eating
644 behaviour following acute and chronic consumption⁽¹³⁵⁾.

645 Despite a growing body of scientific literature exploring the potential benefits of cow's milk
646 consumption, there are several pertinent knowledge gaps that would benefit from further
647 investigation. This is particularly true for cognitive function, hydration status and appetite
648 regulation, where there is little research available, especially in free-living school settings. Further
649 research, especially of a robust and methodologically sound nature (such as double-blind
650 randomised controlled trials), should seek to explore the mechanisms responsible for any effects
651 observed. At present, few studies on this population have attempted to establish mechanisms. **Most**
652 **purported mechanisms given throughout this narrative review come from adult-specific data and**
653 **should therefore be interpreted with some caution as the observations cited may not always be**
654 **replicated in children.** When working with child populations there are various considerations that
655 must be accounted for. The methodological approaches deemed most appropriate for the study of
656 cognitive function, hydration status and appetite regulation in children will differ according to the
657 objective of the study, type of data required, and resources available⁽¹³⁶⁾. In children, it is of great
658 importance to adopt approaches that are non-invasive with a low level of participant stress. Current
659 techniques available to assess cognitive function, hydration status and appetite are arguably invasive,
660 elicit a moderate level of participant **stress** and increased ethical risk. This might explain the current
661 lack of studies **and mechanistic insight from a child perspective.** In recent years, however, research
662 groups have been pursuing non-invasive techniques that offer an opportunity to conduct
663 comprehensive mechanistic work in vulnerable populations. For example, developments in appetite
664 and metabolism research have identified fingertip-capillary blood sampling as an efficacious,
665 comparable and reproducible alternative to antecubital-venous blood sampling for the quantification
666 of appetite-related peptides^(137, 138). These developments will certainly help provide a better
667 understanding of mechanisms that influence appetite and eating behaviour in younger populations,
668 where traditional methods of venous blood sampling might be contraindicated. Furthermore,
669 fingertip-capillary blood sampling offers many advantages including simplistic application,
670 cost/time effectiveness and reduced volume of blood required for analysis⁽¹³⁹⁾.

671 Considering the nutritional-, physical- and health-related impact of cow's milk avoidance, the
672 evidence begins to highlight the importance of increasing cow's milk consumption. Cow's milk is a
673 naturally nutrient-dense foodstuff, providing a significant contribution of several essential nutrients
674 and bioactive constituents that potentially impact human health favourably. Establishing and
675 shaping healthy eating behaviours during the primary-school years is vital. Dietary behaviours
676 shaped throughout the childhood years progress through to adolescence and adulthood⁽¹⁴⁰⁾, making
677 healthy eating environments crucial. For primary-school aged children, the school setting may be an
678 ideal environment to promote cow's milk consumption, and school milk schemes are a great place
679 to start developing healthy eating behaviours given 35 % to 40 % of children's daily nutritional
680 needs are met at school⁽¹⁴¹⁾. Cow's milk is a readily available, accessible and affordable means of
681 providing valuable essential nutrients to the diets of primary-school children. Based on the evidence
682 presented in this manuscript, there appears no reason for primary-school children to limit cow's milk
683 consumption and there may, in fact, be many potential benefits to milk consumption.

684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704

705 **Author Contributions**

706 Benjamin Green and Penny Rumbold conceived and planned this narrative review. Penny Rumbold,
707 Nicola McCullogh, Ruth Boldon and Benjamin Green sourced the relevant literature. All authors
708 contributed to the writing and critically reviewed and approved the final manuscript.

709

710 **Financial Support**

711 This work was supported by funding from Cool Milk Ltd. The funding sponsors had no role in the
712 design, collection, analysis or study interpretation, in the writing of the manuscript, and in the
713 decision to publish the results.

714

715 **Conflicts of Interest**

716 Penny Rumbold has previously received funding from The Dairy Council (UK), Nourishmenow and
717 Cool Milk Ltd. Benjamin Green has previously been supported by The Dairy Council (UK) in the
718 award of a PhD studentship and has received funding from Cool Milk Ltd and is presently an
719 employee of Danone Specialised Nutrition. Lewis James has previously received funding from
720 Volac International Ltd. Emma Stevenson has previously received funding from The Dairy Council
721 (UK) and Arla Food Ingredients. Collectively, the above (with the exception of Cool Milk Ltd) was
722 not related in any way to the work presented in this article. Nicola McCullogh, Ruth Boldon and
723 Crystal Haskell-Ramsay declare no conflicts of interest.

724

725

726

727

728

729

730

731

732

733

734

735

736

737

738

739 **References**

- 740 1. Nagpal R, Behare PV, Kumar M, et al. (2012) Milk, Milk Products, and Disease Free
741 Health: An Updated Overview. *Critical Reviews in Food Science and Nutrition* 52(4):321-333.
- 742 2. Ebringer L, Ferencik M, Krajcovic J. (2008) Beneficial health effects of milk and fermented
743 dairy products — Review. *Folia Microbiologica* 53(5):378-394.
- 744 3. Nongonierma AB, FitzGerald RJ. (2015) The scientific evidence for the role of milk
745 protein-derived bioactive peptides in humans: A Review. *Journal of Functional Foods* 17:640-656.
- 746 4. Ferreira AM, Bislev SL, Bendixen E, et al. (2013) The mammary gland in domestic
747 ruminants: A systems biology perspective. *Journal of Proteomics* 94:110-123.
- 748 5. Rezaei R, Wu Z, Hou Y, et al. (2016) Amino acids and mammary gland development:
749 nutritional implications for milk production and neonatal growth. *Journal of Animal Science and*
750 *Biotechnology* 7(1):20.
- 751 6. Givens DI, Livingstone KM, Pickering JE, et al. (2014) Milk: White elixir or white poison?
752 An examination of the associations between dairy consumption and disease in human subjects.
753 *Animal Frontiers* 4(2):8-15.
- 754 7. Velten H. (2010) *Milk: A global history*: Reaktion Books.
- 755 8. Ludwig DS, Willett WC. (2013) Three Daily Servings of Reduced-Fat Milk: An Evidence-
756 Based Recommendation? *JAMA Pediatrics* 167(9):788-789.
- 757 9. Melnik BC. (2009) Milk – The promoter of chronic Western diseases. *Medical Hypotheses*
758 72(6):631-639.
- 759 10. Morgan PJ, Warren JM, Lubans DR, et al. (2010) The impact of nutrition education with
760 and without a school garden on knowledge, vegetable intake and preferences and quality of school
761 life among primary-school students. *Public Health Nutrition* 13(11):1931-1940.
- 762 11. Patton GC, Viner R. (2007) Pubertal transitions in health. *The Lancet* 369(9567):1130-1139.
- 763 12. Wilson T, Temple NJ.(2016) *How Beverages Impact Health and Nutrition*. In: Wilson T,
764 Temple NJ, editors. *Beverage Impacts on Health and Nutrition: Second Edition*. Cham: Springer
765 International Publishing. p. 3-9.
- 766 13. Rozenberg S, Body J-J, Bruyère O, et al. (2016) Effects of Dairy Products Consumption on
767 Health: Benefits and Beliefs--A Commentary from the Belgian Bone Club and the European
768 Society for Clinical and Economic Aspects of Osteoporosis, Osteoarthritis and Musculoskeletal
769 Diseases. *Calcified tissue international* 98(1):1-17.
- 770 14. Clare DA, Swaisgood HE. (2000) Bioactive Milk Peptides: A Prospectus. *Journal of Dairy*
771 *Science* 83(6):1187-1195.
- 772 15. Parodi PW. (2004) Milk fat in human nutrition. *Australian Journal of Dairy Technology*
773 59(1):3.
- 774 16. MacGibbon AKH, Taylor MW.(2006) Composition and Structure of Bovine Milk Lipids.
775 In: Fox PF, McSweeney PLH, editors. *Advanced Dairy Chemistry Volume 2 Lipids*. Boston, MA:
776 Springer US. p. 1-42.
- 777 17. Wilkinson SB, Tarnopolsky MA, MacDonald MJ, et al. (2007) Consumption of fluid skim
778 milk promotes greater muscle protein accretion after resistance exercise than does consumption of
779 an isonitrogenous and isoenergetic soy-protein beverage. *The American Journal of Clinical*
780 *Nutrition* 85(4):1031-1040.
- 781 18. Tunick MH, Van Hekken DL. (2015) Dairy Products and Health: Recent Insights. *Journal of*
782 *Agricultural and Food Chemistry* 63(43):9381-9388.
- 783 19. Bates B, Lennox A, Prentice A, et al. (2014) National diet and nutrition survey: Results
784 from years 1, 2, 3 and 4 (combined) of the rolling programme (2008/2009-2011/2012): A survey
785 carried out on behalf of public health England and the food standards agency: Public Health
786 England.
- 787 20. Roberts C, Steer T, Maplethorpe N, et al. (2018) National Diet and nutrition survey: results
788 from years 7 and 8 (combined) of the rolling Programme (2014/2015–2015/2016).

- 789 21. Cavadini C, Siega-Riz AM, Popkin BM. (2000) US adolescent food intake trends from 1965
790 to 1996. *Archives of Disease in Childhood* 83(1):18-24.
- 791 22. Fiorito LM, Mitchell DC, Smiciklas-Wright H, et al. (2006) Dairy and Dairy-Related
792 Nutrient Intake During Middle Childhood. *Journal of the American Dietetic Association*
793 106(4):534-542.
- 794 23. Baird DL, Syrette J, Hendrie GA, et al. (2012) Dairy food intake of Australian children and
795 adolescents 2–16 years of age: 2007 Australian National Children's Nutrition and Physical Activity
796 Survey. *Public Health Nutrition* 15(11):2060-2073.
- 797 24. van Rossum CT, Fransen HP, Verkaik-Kloosterman J, et al. (2011) Dutch National Food
798 Consumption Survey 2007-2010: Diet of children and adults aged 7 to 69 years. RIVM report
799 350050006/2011:1-143.
- 800 25. Black RE, Williams SM, Jones IE, et al. (2002) Children who avoid drinking cow milk have
801 low dietary calcium intakes and poor bone health. *The American Journal of Clinical Nutrition*
802 76(3):675-680.
- 803 26. Appel LJ, Moore TJ, Obarzanek E, et al. (1997) A Clinical Trial of the Effects of Dietary
804 Patterns on Blood Pressure. *The New England Journal of Medicine* 336(16):1117-1124.
- 805 27. Pereira MA, Jacobs J, David R., Van Horn L, et al. (2002) Dairy Consumption, Obesity, and
806 the Insulin Resistance Syndrome in Young Adults The CARDIA Study. *JAMA* 287(16):2081-2089.
- 807 28. Elwood PC, Pickering JE, Fehily AM. (2007) Milk and dairy consumption, diabetes and the
808 metabolic syndrome: the Caerphilly prospective study. *Journal of Epidemiology and Community*
809 *Health* 61(8):695-698.
- 810 29. Rockell JEP, Williams SM, Taylor RW, et al. (2005) Two-year changes in bone and body
811 composition in young children with a history of prolonged milk avoidance. *Osteoporosis*
812 *International* 16(9):1016-1023.
- 813 30. Soedamah-Muthu SS, de Goede J. (2018) Dairy Consumption and Cardiometabolic
814 Diseases: Systematic Review and Updated Meta-Analyses of Prospective Cohort Studies. *Current*
815 *Nutrition Reports* 7(4):171-182.
- 816 31. Crichton GE, Bryan J, Buckley J, et al. (2011) Dairy consumption and metabolic syndrome:
817 a systematic review of findings and methodological issues. *Obesity Reviews* 12(5):e190-e201.
- 818 32. Abargouei AS, Janghorbani M, Salehi-Marzjarani M, et al. (2012) Effect of dairy
819 consumption on weight and body composition in adults: a systematic review and meta-analysis of
820 randomized controlled clinical trials. *International Journal of Obesity* 36(12):1485-1493.
- 821 33. Barba G, Russo P. (2006) Dairy foods, dietary calcium and obesity: A short review of the
822 evidence. *Nutrition, Metabolism and Cardiovascular Diseases* 16(6):445-451.
- 823 34. Dougkas A, Reynolds CK, Givens ID, et al. (2011) Associations between dairy consumption
824 and body weight: a review of the evidence and underlying mechanisms. *Nutrition Research*
825 *Reviews* 24(1):72-95.
- 826 35. Dougkas A, Barr S, Reddy S, et al. (2019) A critical review of the role of milk and other
827 dairy products in the development of obesity in children and adolescents. *Nutrition Research*
828 *Reviews* 32(1):106-127.
- 829 36. Cook J, Irwig LM, Chinn S, et al. (1979) The influence of availability of free school milk on
830 the height of children in England and Scotland. *Journal of Epidemiology and Community Health*
831 33(3):171-176.
- 832 37. Campmans-Kuijpers MJE, Singh-Povel C, Steijns J, et al. (2016) The association of dairy
833 intake of children and adolescents with different food and nutrient intakes in the Netherlands. *BMC*
834 *Pediatrics* 16(1):2.
- 835 38. Cook J, Altman DG, Jacoby A, et al. (1975) The contribution made by school milk to the
836 nutrition of primary schoolchildren. *British Journal of Nutrition* 34(1):91-103.

- 837 39. LaRowe TL, Moeller SM, Adams AK. (2007) Beverage Patterns, Diet Quality, and Body
838 Mass Index of US Preschool and School-Aged Children. *Journal of the American Dietetic*
839 *Association* 107(7):1124-1133.
- 840 40. Murphy MM, Douglass JS, Johnson RK, et al. (2008) Drinking Flavored or Plain Milk Is
841 Positively Associated with Nutrient Intake and Is Not Associated with Adverse Effects on Weight
842 Status in US Children and Adolescents. *Journal of the American Dietetic Association* 108(4):631-
843 639.
- 844 41. Rangan AM, Flood VM, Denyer G, et al. (2012) Dairy Consumption and Diet Quality in a
845 Sample of Australian Children. *Journal of the American College of Nutrition* 31(3):185-193.
- 846 42. Wang J, Shang L, Johnson-Down L, et al. (2012) Contribution of flavoured milk on dietary
847 intakes and its association with body mass index percentile in children at high risk of overweight.
848 *The FASEB Journal* 26(S1):lb389-lb389.
- 849 43. Kuriyan R, Thankachan P, Selvam S, et al. (2016) The effects of regular consumption of a
850 multiple micronutrient fortified milk beverage on the micronutrient status of school children and on
851 their mental and physical performance. *Clinical Nutrition* 35(1):190-198.
- 852 44. Albala C, Ebbeling CB, Cifuentes M, et al. (2008) Effects of replacing the habitual
853 consumption of sugar-sweetened beverages with milk in Chilean children. *The American Journal of*
854 *Clinical Nutrition* 88(3):605-611.
- 855 45. Lien DTK, Nhung BT, Khan NC, et al. (2009) Impact of Milk Consumption on Performance
856 and Health of Primary School Children in Rural Vietnam. *Asia Pacific Journal of Clinical Nutrition*
857 18:326–334.
- 858 46. Fiorito LM, Marini M, Mitchell DC, et al. (2010) Girls' Early Sweetened Carbonated
859 Beverage Intake Predicts Different Patterns of Beverage and Nutrient Intake across Childhood and
860 Adolescence. *Journal of the American Dietetic Association* 110(4):543-550.
- 861 47. Weir RR, Johnston M, Lewis C, et al. (2020) Vitamin D3 content of cows' milk produced in
862 Northern Ireland and its efficacy as a vehicle for vitamin D fortification: a UK model. *International*
863 *Journal of Food Sciences and Nutrition*:1-9.
- 864 48. Itkonen ST, Erkkola M, Lamberg-Allardt CJE. (2018) Vitamin D Fortification of Fluid Milk
865 Products and Their Contribution to Vitamin D Intake and Vitamin D Status in Observational
866 Studies—A Review. *Nutrients* 10(8):1054.
- 867 49. Hanks AS, Just DR, Wansink B. (2014) Chocolate Milk Consequences: A Pilot Study
868 Evaluating the Consequences of Banning Chocolate Milk in School Cafeterias. *PLOS ONE*
869 9(4):e91022.
- 870 50. Henry C, Whiting SJ, Phillips T, et al. (2015) Impact of the removal of chocolate milk from
871 school milk programs for children in Saskatoon, Canada. *Applied Physiology, Nutrition, and*
872 *Metabolism* 40(3):245-250.
- 873 51. Fayet-Moore F. (2015) Effect of flavored milk vs plain milk on total milk intake and
874 nutrient provision in children. *Nutrition Reviews* 74(1):1-17.
- 875 52. Rehm CD, Drewnowski A, Monsivais P. (2015) Potential Population-Level Nutritional
876 Impact of Replacing Whole and Reduced-Fat Milk With Low-Fat and Skim Milk Among US
877 Children Aged 2–19 Years. *Journal of Nutrition Education and Behavior* 47(1):61-68.e61.
- 878 53. Popkin BM, D'Anci KE, Rosenberg IH. (2010) Water, hydration, and health. *Nutrition*
879 *Reviews* 68(8):439-458.
- 880 54. D'Anci KE, Constant F, Rosenberg IH. (2006) Hydration and Cognitive Function in
881 Children. *Nutrition Reviews* 64(10):457-464.
- 882 55. Maughan RJ, Watson P, Cordery PA, et al. (2015) A randomized trial to assess the potential
883 of different beverages to affect hydration status: development of a beverage hydration index. *The*
884 *American Journal of Clinical Nutrition* 103(3):717-723.
- 885 56. Bonnet F, Lepicard EM, Cathrin L, et al. (2012) French Children Start Their School Day
886 with a Hydration Deficit. *Annals of Nutrition and Metabolism* 60(4):257-263.

- 887 57. Montenegro-Bethancourt G, Johner SA, Remer T. (2013) Contribution of fruit and vegetable
888 intake to hydration status in schoolchildren. *The American Journal of Clinical Nutrition* 98(4):1103-
889 1112.
- 890 58. Volterman K, Moore D, Obeid J, et al. (2016) The Effect of Postexercise Milk Protein
891 Intake on Rehydration of Children. *Pediatric Exercise Science* 28(2):286.
- 892 59. Volterman KA, Obeid J, Wilk B, et al. (2014) Effect of milk consumption on rehydration in
893 youth following exercise in the heat. *Applied Physiology, Nutrition, and Metabolism* 39(11):1257-
894 1264.
- 895 60. James L. (2012) Milk ingestion in athletes and physically active individuals. *Nutrition*
896 *Bulletin* 37(3):257-261.
- 897 61. Pegoretti C, Antunes AEC, Manchado-Gobatto FdB, et al. (2015) Milk: An Alternative
898 Beverage for Hydration? *Food and Nutrition Sciences* 6(6):8.
- 899 62. Shirreffs SM, Watson P, Maughan RJ. (2007) Milk as an effective post-exercise rehydration
900 drink. *British Journal of Nutrition* 98(1):173-180.
- 901 63. James LJ, Stevenson EJ, Rumbold PLS, et al. (2019) Cow's milk as a post-exercise recovery
902 drink: implications for performance and health. *European Journal of Sport Science* 19(1):40-48.
- 903 64. Maughan RJ, Leiper JB. (1995) Sodium intake and post-exercise rehydration in man.
904 *European Journal of Applied Physiology and Occupational Physiology* 71(4):311-319.
- 905 65. Maughan RJ, Owen JH, Shirreffs SM, et al. (1994) Post-exercise rehydration in man: effects
906 of electrolyte addition to ingested fluids. *European Journal of Applied Physiology and Occupational*
907 *Physiology* 69(3):209-215.
- 908 66. James LJ, Clayton D, Evans GH. (2011) Effect of milk protein addition to a carbohydrate-
909 electrolyte rehydration solution ingested after exercise in the heat. *British Journal of Nutrition*
910 105(3):393-399.
- 911 67. James LJ, Evans GH, Madin J, et al. (2013) Effect of varying the concentrations of
912 carbohydrate and milk protein in rehydration solutions ingested after exercise in the heat. *British*
913 *Journal of Nutrition* 110(7):1285-1291.
- 914 68. Aimutis WR. (2004) Bioactive Properties of Milk Proteins with Particular Focus on
915 Anticariogenesis. *The Journal of Nutrition* 134(4):989S-995S.
- 916 69. Levine RS. (2001) Milk, flavoured milk products and caries. *British Dental Journal*
917 191(1):20-20.
- 918 70. Moynihan P. (2000) Foods and factors that protect against dental caries. *Nutrition Bulletin*
919 25(4):281-286.
- 920 71. Ferrazzano G, Cantile T, Quarto M, et al. (2008) Protective effect of yogurt extract on dental
921 enamel demineralization in vitro. *Australian Dental Journal* 53(4):314-319.
- 922 72. Levine RS, Nugent ZJ, Rudolf MC, et al. (2007) Dietary patterns, toothbrushing habits and
923 caries experience of schoolchildren in West Yorkshire, England. *Community Dental Health*
924 24(2):82-87.
- 925 73. Llana C, Forner L. (2008) Dietary Habits in a Child Population in Relation to Caries
926 Experience. *Caries Research* 42(5):387-393.
- 927 74. Petti S, Simonetti R, D'Arca AS. (1997) The effect of milk and sucrose consumption on
928 caries in 6-to-11-year-old Italian schoolchildren. *European Journal of Epidemiology* 13(6):659-664.
- 929 75. Curtis AM, VanBuren J, Cavanaugh JE, et al. (2018) Longitudinal associations between
930 dental caries increment and risk factors in late childhood and adolescence. *Journal of Public Health*
931 *and Dentistry* 78(4):321-328.
- 932 76. Giacaman RA, Campos P, Muñoz-Sandoval C, et al. (2013) Cariogenic potential of
933 commercial sweeteners in an experimental biofilm caries model on enamel. *Archives of Oral*
934 *Biology* 58(9):1116-1122.

- 935 77. Cross KJ, Huq NL, Palamara JE, et al. (2005) Physicochemical Characterization of Casein
936 Phosphopeptide-Amorphous Calcium Phosphate Nanocomplexes. *Journal of Biological Chemistry*
937 280(15):15362-15369.
- 938 78. McDougall WA. (1977) Effect of Milk on Enamel Demineralization and Remineralization
939 in vitro. *Caries Research* 11(3):166-172.
- 940 79. Weaver CM. (2014) Milk Consumption and Bone Health. *JAMA Pediatrics* 168(1):12-13.
- 941 80. Wiley AS. (2010) Dairy and milk consumption and child growth: Is BMI involved? An
942 analysis of NHANES 1999–2004. *American Journal of Human Biology* 22(4):517-525.
- 943 81. Leighton G, Clark ML. (1929) Milk consumption and the growth of school children: Second
944 preliminary report on tests to the Scottish Board of Health. *British medical journal* 1(3548):23-25.
- 945 82. Orr JB. (1928) Influence of amount of milk consumption on the rate of growth of school
946 children. *British medical journal* 1(3499):140-141.
- 947 83. Bonjour JP, Carrie AL, Ferrari S, et al. (1997) Calcium-enriched foods and bone mass
948 growth in prepubertal girls: a randomized, double-blind, placebo-controlled trial. *The Journal of*
949 *Clinical Investigation* 99(6):1287-1294.
- 950 84. Chan GM, Hoffman K, McMurry M. (1995) Effects of dairy products on bone and body
951 composition in pubertal girls. *The Journal of Pediatrics* 126(4):551-556.
- 952 85. Du X, Zhu K, Trube A, et al. (2004) School-milk intervention trial enhances growth and
953 bone mineral accretion in Chinese girls aged 10–12 years in Beijing. *British Journal of Nutrition*
954 92(1):159-168.
- 955 86. Goulding A, Rockell JEP, Black RE, et al. (2004) Children who avoid drinking cow's milk
956 are at increased risk for prepubertal bone fractures. *Journal of the American Dietetic Association*
957 104(2):250-253.
- 958 87. Iuliano-Burns S, Wang XF, Evans A, et al. (2006) Skeletal benefits from calcium
959 supplementation are limited in children with calcium intakes near 800 mg daily. *Osteoporosis*
960 *International* 17(12):1794-1800.
- 961 88. Zhou Y, Hu Y, Ma Z, et al. (2011) Relation of long-term consumption of school milk to
962 bone density in children. *Journal of hygiene research* 40(1):65-67.
- 963 89. Feskanich D, Bischoff-Ferrari HA, Frazier AL, et al. (2014) Milk consumption during
964 teenage years and risk of hip fractures in older adults. *JAMA Pediatrics* 168(1):54-60.
- 965 90. Sunyecz JA. (2008) The use of calcium and vitamin D in the management of osteoporosis.
966 *Therapeutics and clinical risk management* 4(4):827-836.
- 967 91. Gaucheron F. (2011) Milk and Dairy Products: A Unique Micronutrient Combination.
968 *Journal of the American College of Nutrition* 30(sup5):400S-409S.
- 969 92. Caroli A, Poli A, Ricotta D, et al. (2011) Invited review: Dairy intake and bone health: A
970 viewpoint from the state of the art. *Journal of Dairy Science* 94(11):5249-5262.
- 971 93. Rizzoli R. (2014) Dairy products, yogurts, and bone health. *The American Journal of*
972 *Clinical Nutrition* 99(5):1256S-1262S.
- 973 94. Baker IA, Elwood PC, Hughes J, et al. (1980) A randomised controlled trial of the effect of
974 the provision of free school milk on the growth of children. *Journal of Epidemiology and*
975 *Community Health* 34(1):31-34.
- 976 95. Guo Q, Wang B, Cao S, et al. (2020) Association between milk intake and childhood
977 growth: results from a nationwide cross-sectional survey. *International Journal of Obesity*.
- 978 96. Matkovic V, Landoll JD, Badenhop-Stevens NE, et al. (2004) Nutrition Influences Skeletal
979 Development from Childhood to Adulthood: a Study of Hip, Spine, and Forearm in Adolescent
980 Females. *The Journal of Nutrition* 134(3):701S-705S.
- 981 97. Okada T. (2004) Effect of cow milk consumption on longitudinal height gain in children.
982 *American Journal of Clinical Nutrition* 80(4):1088-1089.

- 983 98. Rona RJ, Chinn S. (1989) School meals, school milk and height of primary school children
984 in England and Scotland in the eighties. *Journal of Epidemiology and Community Health* 43(1):66-
985 71.
- 986 99. Wiley AS. (2005) Does milk make children grow? relationships between milk consumption
987 and height in NHANES 1999–2002. *American Journal of Human Biology* 17(4):425-441.
- 988 100. Matkovic V, Jelic T, Wardlaw GM, et al. (1994) Timing of peak bone mass in Caucasian
989 females and its implication for the prevention of osteoporosis. Inference from a cross-sectional
990 model. *The Journal of Clinical Investigation* 93(2):799-808.
- 991 101. Luiza Loro M, Sayre J, Roe TF, et al. (2000) Early Identification of Children Predisposed to
992 Low Peak Bone Mass and Osteoporosis Later in Life¹. *The Journal of Clinical Endocrinology &*
993 *Metabolism* 85(10):3908-3918.
- 994 102. Hoppe C, Mølgaard C, Vaag A, et al. (2005) High intakes of milk, but not meat, increase s-
995 insulin and insulin resistance in 8-year-old boys. *European Journal of Clinical Nutrition* 59(3):393-
996 398.
- 997 103. Hoppe C, Udam TR, Lauritzen L, et al. (2004) Animal protein intake, serum insulin-like
998 growth factor I, and growth in healthy 2.5-y-old Danish children. *American Journal of Clinical*
999 *Nutrition* 80(2):447-452.
- 1000 104. Rogers I, Emmett P, Gunnell D, et al. (2006) Milk as a food for growth? The insulin-like
1001 growth factors link. *Public Health Nutrition* 9(3):359-368.
- 1002 105. Woods SC, Lutz TA, Geary N, et al. (2006) Pancreatic signals controlling food intake;
1003 insulin, glucagon and amylin. *Philosophical Transactions of the Royal Society B* 361(1471):1219-
1004 1235.
- 1005 106. Berthoud H-R. (2006) Homeostatic and Non-homeostatic Pathways Involved in the Control
1006 of Food Intake and Energy Balance. *Obesity* 14(S8):197S-200S.
- 1007 107. Aziz A, Anderson GH.(2007) The effects of dairy components on food intake and satiety.
1008 In: Saarela M, editor. *Functional Dairy Products*: Woodhead Publishing. p. 19-45.
- 1009 108. Kavezade S, Mozaffari-Khosravi H, Aflatoonian M, et al. (2018) The effects of whole milk
1010 compared to skim milk and apple juice consumption in breakfast on appetite and energy intake in
1011 obese children: a three-way randomized crossover clinical trial. *BMC Nutrition* 4(1):44.
- 1012 109. Mehrabani S, Safavi SM, Mehrabani S, et al. (2016) Effects of low-fat milk consumption at
1013 breakfast on satiety and short-term energy intake in 10- to 12-year-old obese boys. *European*
1014 *Journal of Nutrition* 55(4):1389-1396.
- 1015 110. Mehrabani S, Salehi-Abargouei A, Asemi M, et al. (2014) Effect of Low-fat Milk
1016 Consumption Compared to Apple Juice and Water on the Energy Intake Among 10-12-Year-Old
1017 Obese Boys: A Three-way Cross-over Clinical Trial. *International journal of preventive medicine*
1018 5(11):1405-1411.
- 1019 111. Rumbold PLS, Dodd-Reynolds CJ, Stevenson EJ. (2013) Informing Primary School
1020 Nutritional Policy: Effects of Mid-Morning Snacks on Appetite and Energy Control. *Food and*
1021 *Nutrition Sciences* 4(5):529-537.
- 1022 112. Vien S, Patel B, Panahi S, et al. (2014) The effect of fluid dairy products on food intake,
1023 glycemic and appetite hormone responses in children (120.3). *The FASEB Journal* 28(S1):120.123.
- 1024 113. Anderson GH, Moore SE. (2004) Dietary Proteins in the Regulation of Food Intake and
1025 Body Weight in Humans. *The Journal of Nutrition* 134(4):974S-979S.
- 1026 114. Astrup A. (2005) The satiating power of protein—a key to obesity prevention? *The*
1027 *American Journal of Clinical Nutrition* 82(1):1-2.
- 1028 115. Rolls BJ, Hetherington M, Burley VJ. (1988) The specificity of satiety: The influence of
1029 foods of different macronutrient content on the development of satiety. *Physiology & Behavior*
1030 43(2):145-153.

- 1031 116. Bowen J, Noakes M, Trenergy C, et al. (2006) Energy Intake, Ghrelin, and Cholecystokinin
1032 after Different Carbohydrate and Protein Preloads in Overweight Men. *The Journal of Clinical*
1033 *Endocrinology & Metabolism* 91(4):1477-1483.
- 1034 117. Terpstra AH. (2004) Effect of conjugated linoleic acid on body composition and plasma
1035 lipids in humans: an overview of the literature. *American Journal of Clinical Nutrition* 79(3):352-
1036 361.
- 1037 118. Li J-J, Huang CJ, Xie D. (2008) Anti-obesity effects of conjugated linoleic acid,
1038 docosahexaenoic acid, and eicosapentaenoic acid. *Molecular Nutrition and Food Research*
1039 52(6):631-645.
- 1040 119. Plourde M, Jew S, Cunnane SC, et al. (2008) Conjugated linoleic acids: why the
1041 discrepancy between animal and human studies? *Nutrition Reviews* 66(7):415-421.
- 1042 120. Wang YW, Jones PJH. (2004) Conjugated linoleic acid and obesity control: efficacy and
1043 mechanisms. *International Journal of Obesity* 28(8):941-955.
- 1044 121. Jo H. (2008) Understanding and addressing the epidemic of obesity: an energy balance
1045 perspective. *Journal of Cardiopulmonary Rehabilitation and Prevention* 28(5):345.
- 1046 122. Florence MD, Asbridge M, Veugelers PJ. (2008) Diet Quality and Academic Performance.
1047 78(4):209-215.
- 1048 123. Brindal E, Baird D, Slater A, et al. (2013) The effect of beverages varying in glycaemic load
1049 on postprandial glucose responses, appetite and cognition in 10–12-year-old school children. *British*
1050 *Journal of Nutrition* 110(3):529-537.
- 1051 124. Rahmani K, Djazayeri A, Habibi MI, et al. (2011) Effects of daily milk supplementation on
1052 improving the physical and mental function as well as school performance among children: results
1053 from a school feeding program. *Journal of Research in Medical Sciences* 16(4):469-476.
- 1054 125. Petrova D, Bernabeu Litrán MA, García-Mármol E, et al. (2019) Effects of fortified milk on
1055 cognitive abilities in school-aged children: results from a randomized-controlled trial. *European*
1056 *Journal of Nutrition* 58(5):1863-1872.
- 1057 126. Faught EL, Montemurro G, Storey KE, et al. (2017) Adherence to Dietary
1058 Recommendations Supports Canadian Children’s Academic Achievement. *Canadian Journal of*
1059 *Dietetic Practice and Research* 78(3):102-108.
- 1060 127. Anderson JR, Gunstad J, Updegraff J, et al. (2020) Biological sex and glucoregulation
1061 modulate postprandial cognition following dairy milk and fruit juice in healthy school-age children.
1062 *Nutritional Neuroscience* 23(5):374-383.
- 1063 128. Amiel SA, Pottinger RC, Archibald HR, et al. (1991) Effect of antecedent glucose control
1064 on cerebral function during hypoglycemia. *Diabetes Care* 14(2):109-118.
- 1065 129. Gold AE, MacLeod KM, Thomson KJ, et al. (1995) Cognitive function during insulin-
1066 induced hypoglycemia in humans: Short-term cerebral adaptation does not occur.
1067 *Psychopharmacology* 119(3):325-333.
- 1068 130. Vaz M, Pauline M, Unni US, et al. (2011) Micronutrient Supplementation Improves
1069 Physical Performance Measures in Asian Indian School-Age Children. *The Journal of Nutrition*
1070 141(11):2017-2023.
- 1071 131. Swaminathan S, Edward BS, Kurpad AV. (2013) Micronutrient deficiency and cognitive
1072 and physical performance in Indian children. *European Journal of Clinical Nutrition* 67(5):467-474.
- 1073 132. Zahrou FE, Azlaf M, El Menchawy I, et al. (2016) Fortified Iodine Milk Improves Iodine
1074 Status and Cognitive Abilities in Schoolchildren Aged 7–9 Years Living in a Rural Mountainous
1075 Area of Morocco. *Journal of Nutrition and Metabolism* 2016:8468594.
- 1076 133. Sheng X, Li Z, Ni J, et al. (2019) Effects of Conventional Milk Versus Milk Containing
1077 Only A2 β -Casein on Digestion in Chinese Children: A Randomized Study. *Journal of Pediatric*
1078 *Gastroenterology and Nutrition* 69(3):375-382.

- 1079 134. Kim SH, Kim WK, Kang M-H. (2016) Relationships between milk consumption and
1080 academic performance, learning motivation and strategy, and personality in Korean adolescents.
1081 *Nutrition Research and Practice* 10(2):198-205.
- 1082 135. Green BP, Stevenson EJ, Rumbold PLS. (2017) Metabolic, endocrine and appetite-related
1083 responses to acute and daily milk snack consumption in healthy, adolescent males. *Appetite* 108:93-
1084 103.
- 1085 136. Gibbons C, Finlayson G, Dalton M, et al. (2014) Metabolic phenotyping guidelines:
1086 Studying eating behaviour in humans. *Journal of Endocrinology* 222(2):G1.
- 1087 137. Allsop S, Rumbold PLS, Green BP. (2016) The between-day reproducibility of fasting,
1088 satiety-related analytes, in 8 to 11 year-old boys. *Physiology & Behavior* 164:207-213.
- 1089 138. Green BP, Gonzalez JT, Thomas K, et al. (2014) Agreement between fingertip-capillary and
1090 antecubital-venous appetite-related peptides. *Endocrine connections* 3(4):233.
- 1091 139. Dayre McNally J, Matheson LA, Sankaran K, et al. (2008) Capillary blood sampling as an
1092 alternative to venipuncture in the assessment of serum 25 hydroxyvitamin D levels. *The Journal of*
1093 *Steroid Biochemistry and Molecular Biology* 112(1):164-168.
- 1094 140. Lake AA, Mathers JC, Rugg-Gunn AJ, et al. (2006) Longitudinal change in food habits
1095 between adolescence (11–12 years) and adulthood (32–33 years): the ASH30 Study. *Journal of*
1096 *Public Health* 28(1):10-16.
- 1097 141. Gortmaker SL, Peterson K, Wiecha J, et al. (1999) Reducing Obesity via a School-Based
1098 Interdisciplinary Intervention Among Youth: Planet Health. *Archives of Pediatrics & Adolescent*
1099 *Medicine* 153(4):409-418.

1100

1101

1102

1103

1104

1105

1106

1107

1108

1109

1110

1111

1112

1113

1114

1115

1116

1117 **Table 1.0 Nutritional composition of cow's milk**

Per 100 g	Whole milk	Semi-Skimmed milk	Skimmed milk	Flavoured milk*
Energy (kJ)	274	195	144	270
Protein (g)	3.3	3.5	3.5	3.6
Carbohydrate (g)	4.6	4.7	4.8	9.6
of which sugars (g)	4.6	4.7	4.8	8.9
Fat (g)	3.9	1.7	0.3	1.5
of which saturates	2.5	1.1	0.1	1
monounsaturates	1	0.4	0.1	0.3
polyunsaturates	0.1	Trace	Trace	0.1
trans fatty acids	0.1	0.1	Trace	Trace
Thiamin (mg)	0.03	0.03	0.03	0.03
Niacin (mg)	0.2	0.1	0.1	0.1
Niacin from Tryptophan (mg)	0.6	0.6	0.7	0.8
Calcium (mg)	118	120	125	120
Riboflavin (mg)	0.23	0.24	0.22	0.17
Vitamin B6 (mg)	0.06	0.06	0.06	0.03
Vitamin B12 (µg)	0.6	0.9	0.8	0.1
Folate (µg)	8	9	9	2
Vitamin D (µg)	Trace	Trace	Trace	Trace
Biotin (µg)	2.5	3.0	2.5	2.2
Pantothenate (mg)	0.6	0.7	0.5	0.3
Vitamin C (mg)	2	2	1	Trace
Retinol (µg)	30	19	1	20
Carotene (µg)	19	9	Trace	8
Sodium (mg)	43	43	44	52
Potassium (mg)	155	156	162	168
Magnesium (mg)	11	11	11	12
Phosphorus (mg)	93	94	96	102
Zinc (mg)	0.4	0.4	0.5	0.4
Copper (mg)	Trace	Trace	Trace	Trace
Selenium (µg)	1	1	1	N/A
Manganese (mg)	Trace	Trace	Trace	Trace
Iodine (µg)	31	30	30	N/A

1118 Note: N/A = values not available for this food; Trace = nutrient is present in less than 0.1g per 100g. *Data taken from a
 1119 sample of strawberry and banana flavoured sugar sweetened milk.

1120 Adapted from the Dairy UK. Available at: <http://www.milk.co.uk/publications/default.aspx>

1121

1122

1123

1124

1125

1126

1127

1128

1129

1130 **Table 2. Percentage contribution of cow's milk to average daily total energy, macro- and micro-nutrient intake overtime by sex and age**

	NDNS Rolling Programme Years 1-2; 2009/10				NDNS Rolling Programme Years 3-4; 2011/12				NDNS Rolling Programme Years 5-6; 2013/14				NDNS Rolling Programme Years 7-8; 2015/16	
	4-10 years		11-18 years		4-10 years		11-18 years		4-10 years		11-18 years		4-10 years	11-18 years
	Boys	Girls	Boys	Girls	Boys	Girls	Boys	Girls	Boys	Girls	Boys	Girls	Combined	Combined
Energy	8	8	4	4	7	7	5	4	8	7	4	4	7	4
Protein	14	12	8	7	13	12	9	8	13	13	10	7	11	8
Carbohydrate	5	5	4	3	5	5	4	3	5	5	4	3	5	4
Fat	9	10	6	5	9	10	6	4	10	11	6	4	9	6
Saturated fat	17	16	8	9	15	14	9	7	16	17	9	8	14	9
trans fatty acids	14	12	8	8	19	19	14	11	20	21	14	12	17	13
Vitamin A	11	10	7	7	10	10	9	4	12	12	9	7	11	9
Calcium	31	29	21	22	29	27	22	19	29	29	23	20	26	22
Riboflavin	32	29	22	21	28	29	23	20	29	31	23	18	25	19
Vitamin D	0	0	0	0	1	0	0	0	0	1	0	0	1	1
Sodium	5	4	3	4	6	5	3	2	6	6	3	4	6	4
Potassium	16	15	9	9	14	13	11	10	15	16	11	8	14	9
Magnesium	12	10	8	8	11	10	8	7	11	12	8	7	10	7
Zinc	14	13	8	9	13	13	9	8	13	13	10	8	12	8
Iron	1	1	0	1	1	0	1	0	1	1	1	1	1	1
Selenium	6	5	3	3	5	5	3	3	5	5	3	3	4	3
Folate	9	8	4	5	8	7	5	5	8	9	6	4	7	5
Iodine	40	38	30	30	37	37	32	28	39	28	34	27	37	30

1131 Note: Percentage contribution information taken from National Diet and Nutrition Survey (NDNS) rolling programme. Percentage contributions for Vitamin B₆, Vitamin B₁₂,
1132 Vitamin C and Phosphorus are not reported in the NDNS. * The NDNS rolling programme for 2015/16 ([https://www.gov.uk/government/statistics/ndns-results-from-years-7-](https://www.gov.uk/government/statistics/ndns-results-from-years-7-and-8-combined)
1133 [and-8-combined](https://www.gov.uk/government/statistics/ndns-results-from-years-7-and-8-combined)) stopped providing sex specific intake data so the data are presented as boys and girls combined.

1134 **Table 3** Studies in primary-school aged children that measured nutritional status and hydration with cow's milk consumption

Reference	Details	Design	Methodological approach	Results and conclusion	Effect
<i>Nutritional Status</i>					
Cook et al. (1975) ⁽³⁸⁾	n 312 (159 boys) Age: 8-11 years United Kingdom	Observational	7 day diet record	Nutrient intake was higher in children drinking school milk every day than non-consumers. Boys: energy (P < 0.05), protein (P < 0.01), fat (P < 0.05), calcium (P < 0.001), thiamin (P < 0.01) and riboflavin (P < 0.001) intakes higher than non-consumers Girls: energy (P < 0.001), protein (P < 0.001), fat (P < 0.01), calcium (P < 0.001), thiamin (P < 0.001) and riboflavin (P < 0.001) intakes higher than non-consumers	↑ ↑ ↑
LaRowe et al. (2007) ⁽³⁹⁾	n 793 (410 boys) Age: 6-11 years United States	Observational	24 hour dietary recalls collected during NHANES 2001-2002.	General linear models showed children clustered as high-fat milk consumers had higher (P < 0.05 for all) intakes of energy, protein, riboflavin, folate, vitamin A, vitamin C and calcium compared to clusters of water, SSB, soda and mix/light drinks	↑
Albala et al. (2008) ⁽⁴⁴⁾	n 98 (56 boys) Age: 8-10 years Chile	Randomised, control trial	16 week intervention. Children drank 3 servings/d of flavoured milk or control (no drink)	Protein and calcium intakes increased (P = 0.0001) and energy intake decreased (P = 0.009) with milk compared to controls	↑
Murphy et al. (2008) ⁽⁴⁰⁾	n 2097 (1061 boys) Age: 6-11 years United States	Observational	24 hour dietary recalls collected during NHANES 1999-2002. Participants grouped by age (2-5 years; 6-11 years; 12-18 years).	Calcium, phosphorus, magnesium, potassium, and vitamin A intakes comparable between plain and flavoured milk consumers Calcium, phosphorus, magnesium, potassium, and vitamin A intakes significantly higher (P < 0.05 for all) with milk consumption (plain and flavoured) compared to non-consumers	↔ ↑

Lien et al. (2009) ⁽⁴⁵⁾	n 454 (217 boys) Age: 7-8 years Vietnam	Double-blind, intervention	Volume matched drink (250 mL) served twice daily (morning), 6 days per week for 6 months: (a) multi-micronutrient fortified milk (b) unfortified milk (c) control (no drink)	Intakes of energy, protein, fat, sugar and vitamin A increased ($P < 0.05$ for all) after 6 months in conditions (a) and (b) and compared with condition (c) Serum ferritin and zinc increased ($P < 0.05$ for all) after 6 months in conditions (a) and (b) and compared with condition (c) Zinc levels however also increased in condition (c) ($P < 0.05$)	↑ ↑
Wang et al. (2012) ⁽⁴²⁾	n 632 Age: 8-10 years Canada	Observational	3 24 hour diet recalls	FM drinkers had a higher mean intake for calcium (930 vs. 837 mg; $P = 0.010$), Vitamin D (6.9 vs. 5.9 μg ; $P = 0.021$) and total sugar (99 g vs. 90 g, $P = 0.015$) than non-consumers	↑
Rangan et al. (2013) ⁽⁴¹⁾	n 222 (121 boys) Age: 8-10 years Australia	Observational	3 non-consecutive 24 hour diet recalls	Milk (and dairy) intake significantly associated with increased intake of energy ($P < 0.001$), protein ($P = 0.02$), calcium ($P < 0.001$), phosphorus ($P < 0.001$), magnesium ($P < 0.001$), potassium ($P = 0.009$), zinc ($P = 0.019$), vitamin A ($P < 0.001$), riboflavin ($P < 0.001$), and niacin ($P = 0.03$)	↑
Campmans-Kuijpers et al. (2016) ⁽³⁷⁾	n 1007 (504 boys) Age: 7-13 years Netherlands	Observational	Two non-consecutive 24 hour dietary recalls with an interval of 2 to 6 weeks	Higher milk consumption was associated with significantly higher intakes of energy ($P = 0.003$), protein ($P < 0.0001$), fat ($P = 0.03$), fibre ($P = 0.02$), calcium ($P < 0.0001$), folate ($P < 0.0001$), iodine ($P < 0.0001$), potassium ($P < 0.0001$), magnesium ($P < 0.0001$), phosphorus ($P < 0.0001$), selenium ($P = 0.002$), zinc ($P < 0.0001$) and vitamins B ₁ ($P < 0.0001$), B ₂ ($P < 0.0001$) and B ₁₂ ($P < 0.0001$)	↑
Kuriyan et al. (2016) ⁽⁴³⁾	n 225 (52 boys) Age: 7-10 years India	Double-blind, randomised placebo-controlled	Volume matched drink (200 mL) served twice daily (morning and afternoon), 6 days per week for 5 months: (a) multi-micronutrient fortified milk	At the end of the intervention, levels of blood Vitamin B ₁₂ , and riboflavin were significantly different ($P < 0.001$) between the study groups, in favour of the fortified milk group	↑ ↔

			(b) unfortified milk	Vitamin D, selenium and body iron showed no difference with either group	
Hydration					
Montenegro-Bethancourt et al. (2013) ⁽⁵⁷⁾	n 442 (309 boys) Age: 4-10 years, Germany	Observational	DONALD Study 3 day weighed dietary records	Milk consumption significantly increased FWR (P < 0.05) by 25 mL in boys and 33 mL in girls per 100 g of intake	↑
Volterman et al. (2014) ⁽⁵⁹⁾	n 38 (19 boys) Age: 7-11 years and 14-17 years, Canada	Randomised, repeated-measures cross-over	3 exercise sessions in 34.5 °C (2 × 20-min cycling bouts at 60% peak oxygen uptake) followed by consumption of: (a) plain water (W) (b) a carbohydrate/electrolyte solution (CES) (c) skimmed milk (SM)	Fraction of ingested beverage retained at 2 hour of recovery was greater with SM (74% ± 18%) than W (47% ± 26%) and CES (59% ± 20%, p < 0.001 for both), and greater in CES than water (p < 0.001)	↑
Volterman et al (2016) ⁽⁵⁸⁾	n 15 (7 boys) Age: 10-12 Canada	Randomised, counterbalanced crossover design	45 min of alternating running and cycling exercise in 34.5 °C followed by consumption (150% of body mass loss) of isocaloric and electrolyte-matched beverages containing: (a) 0 g (control), (b) 0.76 g (Lo-PRO) (c) 1.5 g (Hi-PRO) of milk protein/100 mL	Less negative fluid balance at 2 hour of recovery after consumption of Hi-PRO vs. CONT (P = 0.01) Compared with CONT, beverage retention was enhanced by Hi-PRO at 2 h (P < 0.05)	↑ ↑

1135 Abbreviations in order of appearance: FM = flavoured milk; NHANES = National Health and Nutrition Examination Survey; SSB = sugar sweetened beverage;
1136 FWR = free water reserve; W = water; CES = carbohydrate/electrolyte solution; SM = skimmed milk; Lo-PRO = low protein; Hi-PRO = high protein.

1137

1138

1139

1140

1141

Table 4 Studies in primary-school aged children that measured dental health, bone health and physical stature with cow's milk consumption

Reference	Details	Design	Methodological approach	Results and conclusion	Effect
Dental Health					
Petti et al. (1997) ⁽⁷⁴⁾	n 439 (217 boys) Age: 6-11 years Italy	Descriptive cross-sectional	24 hour diet diary, oral examination using Plaque Index and number of decayed, extracted or filled teeth	Milk consumption significantly reduced the probability of carries (P < 0.05); greater significance in high-sucrose (<4/day) consuming children (P < 0.01)	↑
Levine et al. (2007) ⁽⁷²⁾	n 315 Age: 7-11 years England	Prospective cohort/cross-sectional	Three-day dietary diary, tooth brushing habits, dental examination using BASCD survey	Significantly (P < 0.05) less caries associated with moderate consumption (1-2/day) of milk (and dairy) products by children aged 11-15 years	↑
Llena & Forner (2008) ⁽⁷³⁾	n 369 (220 boys) Age: 6-10 years Spain	Descriptive cross-sectional	Food frequency questionnaire, one dental examination for caries and fillings	No significant impact of milk on the incidence of carries	↔
Curtis et al. (2018) ⁽⁷⁵⁾	n 392 (183 boys) Age: 0-19 years United States	Longitudinal, observational	Dietary questionnaires at six-month intervals, cavities assessed age 5, 9, 13 and 17 years	Higher milk intake associated with lower expected adjusted decayed and filled surface increments, however, was not significant	↑
Bone Health and Physical Stature					
Cook et al. (1979) ⁽³⁶⁾	n 1210 (581 boys) Age 6-7 years United Kingdom	Prospective cohort	Free school milk consumed (190 mL/d at school) vs control; height gain	Significantly greater height gain in the milk group in Scottish females (P < 0.05) No difference in males or any groups in England	↑ ↔
Rona & Chinn (1989) ⁽⁹⁸⁾	n 670 Age 5-11 years United Kingdom	Prospective cohort	Free school availability (190 mL/d at school) vs unavailable; change in standardised height score	Significantly greater increments in standardised height scores in 'milk available' group after 2 years in Scotland (P < 0.05) No difference in England	↑ ↔
Baker et al. (1980) ⁽⁹⁴⁾	n 581 (267 boys) Age 7-8 years United Kingdom	Randomised, control trial	12 month intervention; 190 mL/d of milk at school vs control; height gain	Significantly greater height gain in milk group than control (p < 0.05)	↑
Chan et al. (1995) ⁽⁸⁴⁾	n 48 females Age: 9-13 years USA	Randomised control trial	12 month intervention; ≥ 1200 mg/d calcium from dairy vs control; DXA of total body bone mineral content	Greater gain in total body bone mineral content in dairy calcium group (P < 0.001)	↑

Bonjour et al. (1997)⁽⁸³⁾	n 149 females Age: 6-10 years Switzerland	Randomised control trial	12 month intervention; 850 mg/d milk-extracted calcium vs placebo; DXA of regional bone mineral content, height gain and change in lumbar spine length	Greater increment in six-site mean bone mineral content in supplementation group ($P < 0.05$). At 1 year follow-up greater increment in femoral shaft bone mineral content ($P < 0.02$) No difference in height though significance approached ($P \leq 0.08$) favouring supplement group Significant difference in lumbar spine length following intervention and 1 year follow up ($P \leq 0.05$)	↑ ↔ ↑
Black et al. (2002)⁽²⁵⁾	n 250 (120 boys) Age 3-10 years New Zealand	Cross-sectional	Food frequency questionnaire, and DXA of total body and regional bone mineral content and height	Milk avoiders had significantly lower total body bone mineral content than age, sex-matched controls ($P < 0.01$) Milk avoiders significantly shorter than control children of the same age and sex ($P < 0.01$)	↑ ↑
Du et al. (2004)⁽⁸⁵⁾	n 757 (girls) Age 10-12 years China	Randomised control trial	24 month intervention; 330 mL fortified milk (560 mg calcium) / school day vs 300mL fortified milk (560 mg calcium) and 5 or 8 µg cholecalciferol / school day vs control group. DXA of regional and total body bone mineral content with stature assessed pre, mid and post-trial	Greater increase in total-body bone mineral content ($> 1.2\%$) and bone mineral density ($> 3.2\%$) in milk groups compared to control Significantly greater % change in height in milk supplement groups compared to control group ($P < 0.005$)	↑ ↑
Goulding et al. (2004)⁽⁸⁶⁾	n 50 (20 boys) Age: 3-10 years New Zealand	Observational longitudinal	Milk avoiders vs birth cohort population, DXA of total body and regional bone mineral content, history of bone injuries, estimation of calcium intake	Greater number of children reporting fractures and increased total fractures in milk-avoidance compared to control ($P < 0.001$)	↑
Rockell et al. (2005)⁽²⁹⁾	n 46 (18 boys) Age 5-12 years New Zealand	Cross-sectional, longitudinal	Food frequency questionnaire, four-day food diary and DXA of total body bone mineral content and stature	Increase in total body bone mineral content in prior milk avoiders when milk consumption had increased ($P < 0.05$) but remained lower than none milk avoiders	↑

Wiley (2005) ⁽⁹⁹⁾	n 2592 Age 5-11 year United States	Cross-sectional	24 hour recall, milk frequency in past 30 day; height and adult height	Milk intake not associated with height at age 5-11 years (P = 0.385)	↔
Iuliano-Burns et al. (2006) ⁽⁸⁷⁾	n 99 (58 boys) Age 5-11 years Austria	Randomised control trial	10 month intervention; 800 mg/d of calcium from CaCO ₃ vs 800 mg/d of calcium from milk vs placebo, DXA or total and regional bone mineral content	Greater gain in pelvic bone mineral content in pre-pubertal children vs. controls at 10 months in the milk mineral group (P < 0.02) but not biologically meaningful	↔
Albala et al. (2008) ⁽⁴⁴⁾	n 98 (56 boys) Age: 8-10 years Chile	Randomised, control trial	16 week intervention; Children drank 3 servings/d of flavoured milk or control (no drink)	Significantly greater increase (P = 0.01) Smaller increase in height in females in intervention group vs control (P = 0.10)	↑ ↔
Zhou et al. (2011) ⁽⁸⁸⁾	n 435 (188 boys) Age 10-12 years China	Retrospective cohort	School milk group (1 year, 4/wk or 3 year, 1-3/wk) vs. 'seldom milk' group, ulna bone mineral content	Greater ulna bone mineral content in school milk group (P < 0.001)	↑
Guo et al. (2020) ⁽⁹⁵⁾	n 41,439 (19,618 boys) Age 6-17 years China	Cross-sectional	Validated questionnaire assessing milk intake classified as low, medium and high; stature	Milk intake not associated with stature (P > 0.05)	↔

1143 Abbreviations in order of appearance: DXA = dual energy X-ray absorptiometry; CaCO₃ = calcium carbonate.

1144

1145

1146

1147

1148

1149

1150

1151

1152

Table 5 Studies in primary-school aged children that measured appetite and cognitive function with cow's milk consumption

Reference	Details	Design	Methodological approach	Results and conclusion	Effect
Appetite Regulation					
Rumbold et al. (2013) ⁽¹¹¹⁾	n 25 (11 boys) Age: 9-10 years United Kingdom	Randomised controlled cross-over	Energy matched (0.3 MJ) preloads during school mid-morning break: (a) 160 mL of semi-skimmed milk (b) 153 g of apple	Boys and girls felt less hunger and expressed a lower desire to eat when apple was consumed compared to semi-skimmed milk ($P = 0.02$) Energy intake comparable at lunch between semi-skimmed milk and apple ($P > 0.05$)	↓ ↔
Mehrabani et al. (2014) ⁽¹¹⁰⁾	n 34 (obese boys) Age: 10-12 years Iran	Three-way repeated measure randomised controlled cross-over	Volume matched preloads with breakfast (all 240 mL): (a) low fat milk (1.5 %) (b) apple juice (isoenergetic control) (c) water (control)	Energy intake lower at lunch following low fat milk consumption compared to water ($P < 0.005$) and apple juice ($P < 0.05$)	↑
Vien et al. (2014) ⁽¹¹²⁾	Normal and overweight children Mean age: 11.5 years Av age: 11.5 years Canada	Not stated	Volume (250 mL) and energy matched (130 kcal) with the exception of water given pre- and within meals: (a) 1 % fat (1 g/100 g) flavoured milk (b) 2 % fat (2 g/100 g) milk (c) 1.5 % fat (1.5 g/100 g) yogurt drink (d) fruit punch (e) water	<i>Experiment 1</i> Energy intake lower after chocolate milk and yogurt ($P < 0.01$) compared to water Post-meal SA was lower after milk than yogurt ($P < 0.01$) Post-meal SA was lower after milk than fruit punch ($P < 0.01$) <i>Experiment 2</i> GLP-1 AUC was higher after milk than fruit punch and in OWOB compared to NW children ($P < 0.03$). Post-prandial drop in ghrelin was greater after milk than fruit punch in OWOB children (-24 vs. -14%) but was not significant ($P = 0.06$)	↑ ↑ ↑ ↑ ↔ ↔

				Energy intake, insulin and glucose AUC were comparable between all preloads ($P > 0.05$)	
Mehrabani et al. (2016) ⁽¹⁰⁹⁾	n 34 obese boys Age: 10-12 years Iran	Three-way repeated measure randomised controlled cross-over	Volume matched preloads with breakfast (all 240 mL): (a) low fat milk (1.5 %) (b) apple juice (isoenergetic control) (c) water (control)	Higher satiety scores after drinking low fat milk with breakfast compared with water and apple juice ($P < 0.05$) Energy intake lower at lunch following low fat milk compared to water ($P < 0.001$) and apple juice ($P = 0.03$)	↑ ↑
Cognitive Function					
Rahmani et al. (2011) ⁽¹²⁴⁾	n 469 (230 boys) Av age: 8 years Iran	Case-control population-based intervention	Drink at morning school break: (a) tetra-pack sterilised and homogenised milk (250 mL) (b) control (no milk supplementation)	Grade-point average increased from pre- to post-test in girls with milk ($P < 0.05$) No change for the control group, nor for either of the groups of boys. Mean IQ after milk was better in boys compared to boys at post-test in the control ($P < 0.05$)	↑ ↔ ↑
Brindal et al. (2013) ⁽¹²³⁾	n 40 (19 boys) Age: 10-12 years, Australia	Double-blind, randomised three-way repeated measures crossover	Volume matched (455 mL) morning drink following ≥ 8 hours fasting: (a) glucose beverage (b) full milk beverage (c) half milk/glucose beverage	No effect of beverage type in speed of processing, working memory, short-term memory, attention switching, perceptual speed and inspection time No interactions between beverage type and timing of the cognitive testing (60, 120 and 180 minutes post-drink) In the conditions with milk, girls recalled 0.7–0.8 more words but in the glucose condition they recalled 0.5 fewer words ($P = 0.014$). For boys, no difference between beverages was found ($P > 0.09$). Sex differences identified for change in word recall after full milk ($P < 0.001$) and half	↔ ↔ ↑ ↔ ↔

				milk/glucose ($P < 0.001$) conditions: girls showed an increase and boys showed a decrease	
Kuriyan et al. (2016) ⁽⁴³⁾	n 225 (52 boys) Age: 7-10 years India	Double-blind, randomised placebo-controlled design	Volume matched drink (200 mL) served twice daily (morning and afternoon), 6 days per week for 5 months: (a) multi-micronutrient fortified milk (b) unfortified milk	No group \times time interaction for short-term memory and executive functions from baseline	\leftrightarrow
Zahrou et al. (2016) ⁽¹³²⁾	n 200 Age: 7-9 years Morocco	Double-blind, controlled design	Volume matched (200 mL) drink served daily at school for 9 months, including weekends: (a) fortified milk; added potassium iodide and other micronutrients (b) non-fortified milk	Fortified milk group performed significantly better than the non-fortified milk group ($P=0.02$) on a dynamic testing procedure designed to assess children's learning potential	\leftrightarrow
Faught et al. (2017) ⁽¹²⁶⁾	n 1595 (732 boys) Age: 10-11 years Canada	Observational	Analysis of secondary data (food frequency questionnaire) from a 2012 population-based survey using a stratified random sampling design	Boys meeting the nutrition recommendations for milk and alternatives (3-4 servings per day) scored 3.45 % better on an average measure of Math and Language Arts standardised provincial achievement tests than those not meeting the recommendations ($P=0.02$)	\uparrow
Petrova et al. (2019) ⁽¹²⁵⁾	n 103 (52 boys) Age: 8-14 years Spain	Double-blind, randomised controlled design	Volume matched (200 mL) drink served three times per day for 5 months: (a) fortified milk beverage; added micronutrients and essential fatty acids (b) regular full milk	Greater baseline to post-intervention increases in working memory capacity in the fortified milk group (32 % increase) compared to the regular milk group (13 % increase) ($P = 0.027$)	\leftrightarrow
Sheng et al. (2019) ⁽¹³³⁾	n 75 (42 boys) Age: 5-6 years China	Double-blind, multicentre randomised controlled parallel crossover	Volume matched (150 mL) drinks consumed twice daily following meals for 5 days: (a) conventional milk, containing A1 and A2 β -casein (b) milk containing only A2 β -casein	Subtle Cognitive Impairment Test improved following both conventional milk ($P < 0.014$) and milk containing only A2 β -casein ($P < 0.002$) No difference in Subtle Cognitive Impairment Test between drinks	\leftrightarrow \leftrightarrow \leftrightarrow

				<p>Consumers of conventional milk in phase 1 and milk containing only A2 β-casein in phase 2, no effect on error rates in phase 1 ($P = 0.101$) but a decrease in error rates in phase 2 ($P < 0.001$)</p> <p>Consumers of milk containing only A2 β-casein in phase 1 and conventional milk in phase 2, there was a decrease in the error rate in phase 1 which continued through the washout period ($P < 0.001$), but error rates then increased in phase 2 ($P < 0.001$)</p>	↓
Anderson et al. (2020) ⁽¹²⁷⁾	n 84 (39 boys) Age: 8-12 years, United States	Randomised counterbalanced crossover	Volume matched (237 mL) morning drink following ≥ 8 hours' fasting: (a) 1% fat milk (b) apple juice	<p><i>Inhibitory control</i> Reaction time significantly faster after milk compared to apple juice ($P < 0.05$)</p> <p>No effect of beverage on accuracy</p> <p><i>Speeded working memory</i> No difference between the beverages on reaction time ($P = 0.45$)</p> <p>Sex \times beverage interaction for accuracy ($P = 0.003$); compared to apple juice, milk decreased performance accuracy for females whereas non-significant increase in accuracy for males</p> <p><i>Sustained attention</i> Sex \times beverage interaction ($P = 0.02$); compared to apple juice, milk significantly improved reaction time for females whereas reaction time decreased in males (though not significantly)</p>	<p>↑</p> <p>↔</p> <p>↔</p> <p>↑↔</p> <p>↑↔</p>

1154 Abbreviations in order of appearance: SA = subjective appetite; GLP-1 = glucagon like peptide-1; OWOB = overweight and obese; NW = normal weight; AUC
1155 = area under the curve; IQ = intelligence quotient