

Risk Factors for Mobility Decline in Community-Dwelling Older Adults: A Systematic Literature Review

Philippa J.A. Nicolson, Maria T. Sanchez-Santos, Julie Bruce, Shona Kirtley, Lesley Ward, Esther Williamson, and Sarah E. Lamb

Mobility is essential to maintaining independence for older adults. This systematic review aimed to summarize evidence about self-reported risk factors for self-reported mobility decline; and to provide an overview of published prognostic models for self-reported mobility decline among community-dwelling older adults. Databases were searched from inception to June 2, 2020. Studies were screened by two independent reviewers who extracted data and assessed study quality. Sixty-one studies (45,187 participants) were included, providing information on 107 risk factors. High-quality evidence and moderate/large effect sizes for the association with mobility decline were found for older age beyond 75 years, the presence of widespread pain, and mobility modifications. Moderate–high quality evidence and small effect sizes were found for a further 21 factors. Three model development studies demonstrated acceptable model performance, limited by high risk of bias. These findings should be considered in intervention development, and in developing a prediction instrument for practical application.

Keywords: mobility disability, predictors, self-reported

Mobility is defined as “the ability to move oneself within community environments that expand from one’s home, to the neighborhood, and to regions beyond” (Webber, Porter, & Menec, 2010). Maintaining mobility is fundamental to aging well for older adults (Guralnik et al., 1993). Limited mobility is linked to functional decline, mortality, and increased health care costs (Guralnik et al., 1993). Changes in mobility are important as conceptually they precede disability within models of disablement and are potential targets for rehabilitation (Verbrugge & Jette, 1994).

Mobility may be assessed using objective or self-reported measures. While it is acknowledged that different aspects of mobility may be measured with each, multiple studies have found a moderate correlation between the two approaches (Alexander et al., 2000; Syddall, Westbury, Cooper, & Sayer, 2015). Self-reported mobility is important as it is the most commonly used measure clinically (Chung, Demiris, & Thompson, 2015). Objective physical performance measures do not reflect the individual’s perceptions about their own mobility, and are most often conducted in a supervised, controlled environment, thus unreflective of a real-

world situation. Potential risk factors for mobility decline may also be measured objectively or by self-report. Self-reported measures of these factors are also important, as they have low response burden, can capture both current and historical information, and can assess psychological and social factors.

Two previous reviews of risk factors for mobility decline combined objective and self-reported measures of risk factors and mobility, but these were narrative rather than systematic reviews (Rantakokko, Mänty, & Rantanen, 2013; Yeom, Fleury, & Keller, 2008). Consolidated evidence of self-reported risk factors contributing to self-reported mobility decline is needed to inform clinical decision making and to inform the development of a prognostic tool to help clinicians and researchers identify older adults at risk of mobility decline. The primary objective of this systematic review was to synthesize available evidence for self-reported factors which predict decline in self-reported mobility after 12 months to 5 years of follow-up among community-dwelling older adults. The secondary objective was to identify, describe, and synthesize the predictive accuracy of prognostic models developed to predict risk of self-reported mobility decline in community-dwelling older adults.

Methods

This systematic review is registered at the International Prospective Register of Systematic Reviews database (PROSPERO 2019 CRD42019135420. Available from: https://www.crd.york.ac.uk/prospero/display_record.php?ID=CRD42019135420). The review is reported according to the Preferred Reporting Items for Systematic Review and Meta-Analyses guidelines (Moher, Liberati, Tetzlaff, & Altman, 2009), and the CHECKlist for critical Appraisal and data extraction for systematic Reviews of prediction Modelling Studies (CHARMS) (Moons et al., 2014).

Search Strategy

The search strategy was developed jointly with an information scientist (SK). We searched the MEDLINE (via OVID), EMBASE

© 2021 The Authors. Published by Human Kinetics, Inc. This is an Open Access article distributed under the terms of the Creative Commons Attribution 4.0 International License, CC BY 4.0, which permits unrestricted noncommercial and commercial use, distribution, and reproduction in any medium, provided the original work is properly cited, the new use includes a link to the license, and any changes are indicated. See <http://creativecommons.org/licenses/by/4.0>. This license does not cover any third-party material that may appear with permission in the article.

Nicolson, Sanchez-Santos, Kirtley, Williamson, and Lamb are with the Nuffield Department of Orthopaedics, Rheumatology and Musculoskeletal Sciences (NDORMS), University of Oxford, Oxford, United Kingdom. Bruce is with the Warwick Clinical Trials Unit, Warwick Medical School, University of Warwick, Coventry, United Kingdom. Ward is with the Department of Sport, Exercise & Rehabilitation, Northumbria University, Newcastle upon Tyne, United Kingdom. Lamb is also with the College of Medicine and Health, University of Exeter, Exeter, United Kingdom. Nicolson (philippa.nicolson@ndorms.ox.ac.uk) is corresponding author.

(via OVID), PsycINFO (via OVID), and CINAHL (via EBSCOhost) databases to identify studies that assessed the predictive value of self-reported risk factors on self-reported mobility decline. Search terms included controlled vocabulary (e.g., MeSH) terms and free-text terms. No date or language limits were applied to the search. The search strategy for MEDLINE is presented in [Supplementary Table S1](#) (available online), and this was translated to the relevant controlled vocabulary headings and syntax for each database. The search was conducted on June 26, 2019, with an update run on June 2, 2020. Supplementary searches of the reference and citation lists of included studies and relevant systematic reviews were undertaken to identify additional eligible studies.

Study Selection

Inclusion criteria.

- Studies involving community-dwelling older adults with a minimum mean age of 60 years at recruitment.
- Longitudinal cohort studies evaluating the ability of self-reported factors to predict self-reported mobility decline. Randomized controlled trials were eligible if they provided appropriate data and treatment-effect modification analyses were reported.
- Prognostic model studies, with or without validation. Models had to contain at least two self-reported risk factors, and include age and sex as a minimum of adjustment factors.
- A self-reported measure of mobility, defined as difficulty with walking and/or difficulty in climbing stairs and/or life-space assessment, assessed at baseline and follow-up.
- A clear definition of mobility decline.
- Where a study collected both self-reported measures and objective measures, the study was eligible if data reported the relationship between self-reported measures and self-reported outcomes.
- Follow-up period of between 12 months and 5 years.
- Published in English, Spanish, or Portuguese (languages spoken by the research team).

Exclusion criteria.

- Studies including only patients with a specific medical condition (e.g., Parkinson's disease, dementia, stroke).
- Studies conducted in residential aged care settings.
- Studies published in abstract form only.

Search results were imported into Covidence (Veritas Health Innovation, Melbourne, Australia, www.covidence.org) for screening. Titles and abstracts of all identified citations were independently screened by two reviewers (PN and MS). Following title and abstract screening, the full text of all potentially eligible articles was retrieved and screened independently by the same two reviewers. Inclusion disagreements were resolved through discussion, with an adjudicator (PN and MS) available.

Data Extraction

A data extraction form in Microsoft Excel (Microsoft Corporation, Redmond, WA) was developed and piloted independently by two reviewers before independent data extraction from all articles (PN and MS). Disagreements were resolved by discussion.

Studies were categorized as etiological, predictor finding, or model development using the following definitions (Moons, Royston, Vergouwe, Grobbee, & Altman, 2009):

- **Etiological studies:** A study examining causal association between predefined risk factor/s and mobility decline.
- **Predictor finding studies:** A study examining the contribution of multiple risk factors to predict mobility decline, without quantification of the predictive ability of the model.
- **Model development studies:** A study that aimed to develop a multivariable prediction model, assigning weighting to each identified risk factor and developing a final prediction model. These studies may or may not include internal validation studies.

Data extracted from all studies included: country of origin, year of study conduct, recruited and analyzed sample sizes, study design, type of study, length of follow-up, and participant characteristics (population source and setting, eligibility criteria, recruitment method, and sociodemographic characteristics). Method of assessment was recorded for each risk factor and mobility outcome.

For etiological studies, we extracted data for the specific risk factor/s of interest as stated in the study. For predictor finding studies, we extracted data on all included risk factors. We extracted unadjusted and adjusted effect estimates (odds ratio [OR], relative risk [RR], or hazard ratio [HR]) and corresponding *SE* or confidence intervals for each risk factor. For model development studies, we extracted modeling method, handling of risk factors and missing data, method for selection of risk factors, model performance measures, and model evaluation. We contacted authors when insufficient information was reported.

Assessment of Risk of Bias in Included Studies

Two reviewers (PN and MS) independently assessed the risk of bias in included studies using the Quality In Prognosis Studies tool for etiological and predictor finding studies and the Prediction model Risk of Bias Assessment Tool for model development studies (Hayden, van der Windt, Cartwright, Cote, & Bombardier, 2013; Wolff et al., 2019).

The Quality In Prognosis Studies tool assesses risk of bias across six domains: study participation, study attrition, risk factor measurement, outcome measurement, study confounding, and statistical analysis and reporting (Hayden et al., 2013). Each domain was rated as low, moderate, or high risk of bias (Hayden et al., 2013). The Prediction model Risk of Bias Assessment tool includes 20 signaling questions grouped within four domains (participant selection, predictors, outcome, and analysis) to evaluate the risk of bias of prediction models (Wolff et al., 2019). Studies were classified as low risk of bias if all domains were rated as low risk of bias. Any disagreements in scoring were resolved through discussion.

Data Synthesis and Analysis

Descriptive characteristics of all included studies were summarized and assessed for suitability for meta-analysis by considering the clinical heterogeneity of included studies based on population, definition of risk factors and/or mobility decline, and methodology (Riley et al., 2019). Adjusted effect estimates (minimum adjustment for age and gender/sex) were reported in the narrative analysis. Odds ratios of 1.4–2.5 were considered as small effect, 2.5–4.25 were classified as moderate, and 4.25 or greater as large (Chinn, 2000). Odds ratios were considered to indicate no association if the 95% confidence interval crossed 1.0. Information from risk factor and prognostic model studies were evaluated separately because their objectives and model-building methods are different (van den Berg et al., 2013). Performance of prognostic models were summarized and presented narratively.

Narrative Analysis

We applied the following criteria proposed by Hayden, Tougas, Riley, Iles, and Pincus (2014) to categorize the consistency of evidence for each risk factor. Consistent evidence defined as two or more studies from two or more cohorts with $\geq 75\%$ of studies showing the same direction of effect (i.e., four studies from three cohorts, of which three studies show the same direction of effect). Limited evidence defined as: single study or multiple studies from the same cohort. Inconsistent evidence defined as: two or more studies from two or more cohorts with $< 75\%$ of studies showing the same direction of effect (i.e., four studies from three cohorts, of which two studies show the same direction of effect).

Assessment of Quality of Evidence

The quality of the body of evidence was synthesized using the Grading of Recommendations Assessment, Development and Evaluation approach modified for use in reviews of risk factor studies (Huguet et al., 2013). Overall quality of evidence was rated as high, moderate, low, or very low considering study limitations, inconsistency of results, indirectness, imprecision, publication bias, moderate/large effect size, and the presence of a dose effect.

Results

Study Selection

The process of study selection is shown in Figure 1. In total, 26,079 citations (18,201 with duplicates removed) were identified. Of these, 187 articles progressed to full-text review, and 61 studies were included in the review. Among included studies, 47 were classified as etiological studies, 11 as predictor finding studies, and three were model development studies.

Characteristics of Included Studies

Characteristics and details of the 61 included studies are presented in [Supplementary Table S2](#) (available online). Studies were published between 1993 and 2019. A total of 45,187 participants were analyzed. Included studies were all secondary analyses of data derived from 25 established cohorts. The cohorts from which the highest number of studies originated were the Nordic Research on Ageing Study (11 studies; Avlund, Damsgaard, & Osler, 2004; Avlund, Damsgaard, Sakari-Rantala, Laukkanen, & Schroll, 2002; Avlund, Davidsen, & Schultz-Larsen, 1995; Avlund, Lund, Holstein, & Due, 2004; Avlund, Osler, Damsgaard, Christensen, & Schroll, 2000; Avlund, Pedersen, & Schroll, 2003; Avlund, Vass, & Hendriksen, 2003; Jørgensen, Lund, Siersma, & Nilsson,




Figure 1 — Preferred Reporting Items for Systematic Reviews and Meta-Analyses flowchart.

2017; Nilsson, Avlund, & Lund, 2010; Nilsson, Lund, & Avlund, 2008; Schroll, Avlund, & Davidsen, 1997), the Health, Ageing and Body Composition Study (seven studies; Carbone et al., 2013; Duan-Porter et al., 2019; Koster et al., 2005; Lee et al., 2005; Mehta et al., 2007; Thorpe et al., 2011; Visser et al., 2005), and the first Women's Health and Ageing Study (four studies; Brenes et al., 2005; Leveille, Bean, Ngo, McMullen, & Guralnik, 2007; Leveille et al., 2001; Onder et al., 2003). The majority of studies were based on populations from the United States (27 studies; 11 cohorts) or Finland (16 studies; five cohorts).

Age and sex characteristics. The analyzed sample size of included studies ranged from 136 to 6,981 participants. The mean of the mean ages of participants at baseline was 74.3 years. Mean ages at baseline ranged from 64 to 80 years. The gender distribution of participants varied: nine studies (Brenes et al., 2005; Chaves, Garrett, & Fried, 2000; Fried, Bandeen-Roche, Chaves, & Johnson, 2000; Leveille et al., 2001, 2007; Manty et al., 2009; Onder et al., 2003; Viljanen et al., 2012; Weiss, Wolff, Eggleston, Seplaki, & Fried, 2012) (from four cohorts) reported data for women only, and three studies (Avlund et al., 1995; Liljas et al., 2016; van den Brink et al., 2004) (from two cohorts) reported data for men only. Among studies reporting on mixed-gender populations, the proportion of women varied from 46% to 75%.

Measures of mobility and mobility decline. Thirty-three different measures of mobility were used. The two most commonly used were the Mobility-Help (Mobility-H) scale (11 studies; Avlund, Damsgaard, & Osler, 2004; Avlund et al., 1995, 2000, 2002; Avlund, Lund, et al., 2004; Avlund, Pedersen, & Schroll, 2003; Avlund, Vass, & Hendriksen, 2003; Jørgensen et al., 2017; Nilsson et al., 2008, 2010; Schroll et al., 1997) and the single-item question "Do you have any difficulty walking 1/4 mile/400 m or climbing 1 flight of stairs/10 steps without assistance?" (10 studies; Auais et al., 2018, 2019; Chen, Covinsky, Stijacic Cenzer, Adler, & Williams, 2012; Deshpande, Metter, Guralnik, Bandinelli, & Ferrucci, 2014; Eggermont et al., 2014; Koster et al., 2005; Lee et al., 2005; Mehta et al., 2007; Thakral, Shi, Shmerling, Bean, & Leveille, 2014; Thorpe et al., 2011). Almost all included studies (59/61; 97%) used mobility outcomes that focused solely on walking and/or climbing stairs. Only two studies (Polku et al., 2015; Tsuji, Rantakokko, Portegijs, Viljanen, & Rantanen, 2018) used the Life Space Assessment which includes ability to move beyond one's neighborhood and/or town by means, such as public transport or driving. There was also variability in the definition of mobility decline. Among studies using the Mobility-H scale, mobility decline was defined in three different ways, most commonly by progression from "0 activities requiring help" at baseline to "need help in at least one activity" at follow-up (8/11 studies, 73%). Mobility decline was measured at a mean of 37 months from recruitment (range = 12–60 months).

Risk of Bias in Included Studies

Of the 61 included studies, the 58 etiological and predictor finding studies were appraised using Quality In Prognosis Studies (Supplementary Table S3 [available online]). Risk of bias varied across the studies: 11 studies (Avlund, Vass, & Hendriksen, 2003; Brenes et al., 2005; Guralnik et al., 1993; Ho, Woo, Yuen, Sham, & Chan, 1997; Jørgensen et al., 2017; Keskinen, Rantakokko, Suomi, Rantanen, & Portegijs, 2020; Mehta et al., 2007; Polku et al., 2015; Thorpe et al., 2011; Tsuji et al., 2018; Visser et al., 2005) (11/58, 19%) were rated as low risk of bias for all six domains and 30

studies (Abizanda et al., 2013; Auais et al., 2018, 2019; Avlund, Damsgaard, & Osler, 2004; Avlund et al., 1995, 2000, 2002; Ayis, Gooberman-Hill, Bowling, & Ebrahim, 2006; Carbone et al., 2013; Deshpande et al., 2014; Duan-Porter et al., 2019; Eggermont et al., 2014; Eronen, von Bonsdorff, Rantakokko, & Rantanen, 2013; Fried et al., 2000; Koster et al., 2005; LaCroix, Guralnik, Berkman, Wallace, & Satterfield, 1993; Lang, Llewellyn, Langa, Wallace, & Melzer, 2008; Leveille et al., 2001; Liljas et al., 2016; Manty et al., 2007; Manty et al., 2009; Onder et al., 2003; Rantakokko, Iwarsson, Manty, Leinonen, & Rantanen, 2012; Rantakokko, Portegijs, Viljanen, Iwarsson, & Rantanen, 2016; Schroll et al., 1997; Simonsick, Aronson, et al., 2018; Simonsick et al., 2016; Simonsick, Schrack, et al., 2018; Thakral et al., 2014; Weiss et al., 2012) (30/58, 52%) were rated as low risk of bias for five domains. Sources of bias most commonly related to study attrition, specifically lack of details provided regarding reasons for loss to follow-up or differences in key characteristics between those included and those lost to follow-up. Nine studies (Avlund et al., 2000; Ayis, Bowling, Gooberman-Hill, & Ebrahim, 2007; Ayis et al., 2006; Eronen et al., 2013; Manty et al., 2009; Onder et al., 2003; Rantakokko et al., 2012; van den Brink et al., 2004; Weiss et al., 2012) (9/58, 16%) were rated as high risk of bias for the study attrition domain, and 33 studies (Abizanda et al., 2013; An & Lu, 2016; Auais et al., 2018, 2019; Avlund, Damsgaard, & Osler, 2004; Avlund et al., 2002; Avlund, Lund, et al., 2004; Avlund, Pedersen, & Schroll, 2003; Carbone et al., 2013; Chen et al., 2012; Crimmins & Saito, 1993; Deshpande et al., 2014; Duan-Porter et al., 2019; Eggermont et al., 2014; Fried et al., 2000; Guralnik, Ferrucci, Balfour, Volpato, & Di Iorio, 2001; Koster et al., 2005; LaCroix et al., 1993; Lang et al., 2008; Lee et al., 2005; Leveille et al., 2001, 2007; Lindberg & Tilvis, 1998; Manty et al., 2007; Nilsson et al., 2008, 2010; Pine, Gurland, & Chren, 2000, 2002; Rantakokko et al., 2016; Sauvaget, Tsuji, Aonuma, & Hisamichi, 1999; Simonsick, Schrack, et al., 2018; Thakral et al., 2014; Viljanen et al., 2012) (33/58, 57%) were rated as moderate risk of bias. Other major sources of bias related to statistical analysis and reporting, for which four studies (Crimmins & Saito, 1993; Pine et al., 2002; Simonsick, Aronson, et al., 2018; van den Brink et al., 2004) (4/58, 7%) were rated as high risk of bias, and 10 studies (Avlund, Damsgaard, & Osler, 2004; Avlund et al., 1995; Avlund, Lund, et al., 2004; Chen et al., 2012; Liljas et al., 2016; Lindberg & Tilvis, 1998; Pine et al., 2000; Sauvaget et al., 1999; Schroll et al., 1997; Viljanen et al., 2012) (10/58, 17%) were rated as moderate risk of bias.

The results of the Prediction model Risk of Bias Assessment tool appraisal for the model development studies are presented in Supplementary Table S4 (available online). All three studies (Chaves et al., 2000; Papachristou et al., 2017; Reynolds & Silverstein, 2003) were judged as having high risk of bias for the analysis domain. This was due to the absence of model performance measures, selection of predictor variables based on univariable analyses, and lack of reporting results of multivariable analyses. All three studies were judged to have both high risk of bias overall and high concern regarding applicability to the review question.

Risk Factors for Mobility Decline Identified in Etiological and Predictor Finding Studies

A synthesis of the self-reported risk factors for mobility decline examined in etiological and predictor finding studies is presented in Table 1 and Supplementary Table S5 (available online). Due to

Table 1 Overview of Risk Factors for Mobility Decline, Presented by Direction of Association, Using the Adapted GRADE (Huguet et al., 2013)

Risk factors	Number of participants ^a	Number of studies examining risk factor	Number of cohorts	Range of OR reported	Quality of evidence
Consistent evidence of increased risk of mobility decline ^b					
Older age	16,452	9 (Auais et al., 2018, 2019; Carbone et al., 2013; Crimmins & Saito, 1993; Deshpande et al., 2014; Guralnik et al., 1993, 2001; Ho et al., 1997; Sauvaget et al., 1999)	7	1.1–5.8 >75: 3.5–16.5	High
Mobility modification ^c	1,858	5 (Fried et al., 2000; Manty et al., 2009; Pine et al., 2002; Rantakokko et al., 2016; Weiss et al., 2012)	4	2.5–6.2 RR = 5.7–8.9	High
Widespread pain	1,102	3 (Eggermont et al., 2014; Leveille et al., 2001, 2007)	2	2.5–2.8 RR = 3.6	Moderate
Gender/sex (female/women)	14,528	8 (Auais et al., 2019; Ayis et al., 2007; Carbone et al., 2013; Crimmins & Saito, 1993; Deshpande et al., 2014; Guralnik et al., 2001; Sauvaget et al., 1999; Schroll et al., 1997)	8	1.2–2.6	High
Low annual income	10,047	2 (Guralnik et al., 1993; Koster et al., 2005)	2	1.3–2.6	Moderate
Low number of financial assets	6,533	5 (Deshpande et al., 2014; Jørgensen et al., 2017; Koster et al., 2005; Nilsson et al., 2010; Thorpe et al., 2011)	3	1.3–1.5	High
Low diversity in social relations	4,308	3 (Avlund, Lund, et al., 2004; Ho et al., 1997; Nilsson et al., 2010)	2	1.6–2.8	High
Low social engagement	3,292	4 (Avlund, Lund, et al., 2004; Ayis et al., 2006; Jørgensen et al., 2017; Nilsson et al., 2010)	2	1.5–2.9	High
Low physical activity	11,152	6 (Abizanda et al., 2013; Avlund et al., 2000; Carbone et al., 2013; LaCroix et al., 1993; Schroll et al., 1997; Visser et al., 2005)	5	1.5–4.3 RR = 4.3	High
Not walking each day	4,460	2 (Ho et al., 1997; Visser et al., 2005)	2	1.5–2.4	Moderate
Greater fatigue with exertion	1,514	3 (Abizanda et al., 2013; Simonsick et al., 2016; Simonsick, Schrack, et al., 2018)	2	1.1–1.5	Moderate
Greater tiredness in daily activities	1,975	5 (Avlund et al., 1995, 2002; Avlund, Pedersen, & Schroll, 2003; Simonsick et al., 2016; Simonsick, Schrack, et al., 2018)	2	1.5–10.8	Moderate
Hip pain	2,712	2 (Carbone et al., 2013; Eggermont et al., 2014)	2	1.5 RR = 4.5	Moderate
Knee pain	2,712	2 (Carbone et al., 2013; Eggermont et al., 2014)	2	2.0 RR = 2.7	Moderate
Fall/s in past year	802	2 (Carbone et al., 2013; Manty et al., 2009)	2	1.3–3.2	Moderate
Fear of falling	1,207	2 (Auais et al., 2018; Viljanen et al., 2012)	2	1.1–2.9	Moderate
Increasing no. health conditions	8,723	4 (Guralnik et al., 1993, 2001; Ho et al., 1997; Schroll et al., 1997)	3	1.4–3.4 RR = 1.3–5.3	High
Depression	4,818	4 (Ayis et al., 2006; Carbone et al., 2013; Deshpande et al., 2014; Ho et al., 1997)	4	1.1–6.5	High
Eye conditions	4,515	4 (Ayis et al., 2007; Carbone et al., 2013; Ho et al., 1997; Viljanen et al., 2012)	4	1.6–1.8	High
Heart conditions	3,537	2 (Crimmins & Saito, 1993; Ho et al., 1997)	2	1.4–1.7	High
History of stroke	10,518	4 (Crimmins & Saito, 1993; Guralnik et al., 1993, 2001; Ho et al., 1997)	3	1.4–2.7	Moderate
Sensory difficulties	2,344	2 (Crimmins & Saito, 1993; Viljanen et al., 2012)	2	1.3–2.5	Low
Poor self-rated health	7,537	4 (Avlund et al., 2000; Ayis et al., 2007; Guralnik et al., 2001; Ho et al., 1997)	4	1.3–3.3	High
Hospital stay in past year	4,963	2 (Crimmins & Saito, 1993; Duan-Porter et al., 2019)	2	1.1–1.3	Moderate

(continued)

Table 1 (continued)

Risk factors	Number of participants ^a	Number of studies examining risk factor	Number of cohorts	Range of OR reported	Quality of evidence
Consistent evidence of no association with mobility decline ^b					
Weight gain in past 12 months	4,415	2 (Ho et al., 1997; Lee et al., 2005)	2	0.9 HR = 0.88	Moderate
Cancer (previous or current)	7,213	3 (Ayis et al., 2007; Guralnik et al., 2001; Ho et al., 1997)	3	0.5–1.3	Moderate
Higher no. physical environment barriers	1,180	4 (Ayis et al., 2006; Eronen et al., 2013; Rantakokko et al., 2012, 2016)	3	0.9–1.5	Moderate
Limited evidence of increased risk of mobility decline ^d					
Housing tenure (renting)	386	1 (Avlund, Lund, et al., 2004)	1	1.9	Very low
High neighborhood deprivation	3,525	1 (Lang et al., 2008)	1	1.8	Very low
Not having health insurance	2,622	1 (Carbone et al., 2013)	1	1.3	Very low
Not driving a car	581	1 (Tsuji et al., 2018)	1	1.5	Very low
Living alone	2,825	5 (Avlund, Damsgaard, & Osler, 2004; Avlund et al., 1995; Jørgensen et al., 2017; Nilsson et al., 2008, 2010; Tsuji et al., 2018)	1	1.1–1.7	Very low
Widowhood	829	1 (van den Brink et al., 2004)	1	1.7–1.8	Very low
Currently not exercising	1,483	1 (Ho et al., 1997)	1	2.1	Very low
Low amount of exercise per day	1,483	1 (Ho et al., 1997)	1	2.2–3.4	Very low
Other functional difficulties	2,145	1 (Crimmins & Saito, 1993)	1	1.3–1.4	Very low
Low energy level	579	2 (Simonsick et al., 2016; Simonsick, Schrack, et al., 2018)	1	1.1–1.2	Very low
Low functional reserve capacity	427	1 (Ayis et al., 2006)	1	6.8	Very low
Increasing pain severity	411	1 (Eggermont et al., 2014)	1	2.0	Very low
Increasing pain interference	411	1 (Eggermont et al., 2014)	1	2.4	Very low
Back pain + other pain	412	1 (Eggermont et al., 2014)	1	3.3	Very low
Hip pain + other pain	412	1 (Eggermont et al., 2014)	1	4.8	Very low
Knee pain + other pain	413	1 (Eggermont et al., 2014)	1	3.3	Very low
Ankle/foot pain + other pain	412	1 (Eggermont et al., 2014)	1	3.7	Very low
Joint stiffness	524	1 (Thakral et al., 2014)	1	1.3–1.6	Very low
History of fracture	2,286	2 (Carbone et al., 2013; Ho et al., 1997)	1	1.1–1.6	Very low
Incontinence	5,263	1 (Guralnik et al., 2001)	1	1.1–1.2	Very low
Memory loss	1,483	1 (Ho et al., 1997)	1	1.4	Very low
Respiratory conditions	491	1 (Ayis et al., 2007)	1	1.3–3.5	Very low
Health deterioration in past year	427	1 (Ayis et al., 2006)	1	4.3	Very low
Major life changes in past 6 months	491	1 (Ayis et al., 2007)	1	1.5–2.3	Very low
Low sexual satisfaction	178	1 (Onder et al., 2003)	1	2.6	Very low

(continued)

Table 1 (continued)

Risk factors	Number of participants ^a	Number of studies examining risk factor	Number of cohorts	Range of OR reported	Quality of evidence
Weight change intention	2,907	1 (Lee et al., 2005)	1	HR: 1.6	Very low
Acetaminophen use	2,486	1 (Carbone et al., 2013)	1	1.6	Very low
Antidepressant use	4,242	1 (Carbone et al., 2013)	1	1.1	Very low
NSAID use	2,486	1 (Carbone et al., 2013)	1	1.4	Very low
Lack of resting places and long distances	266	1 (Rantakokko et al., 2012)	1	1.9–2.2	Very low
Limited evidence of no association with mobility decline ^d					
Weight change in past 12 months	2,932	1 (Lee et al., 2005)	1	HR = 0.8–1.49	Very low
Manual occupation	243	1 (Avlund et al., 2000)	1	1.1	Very low
Low satisfaction with living arrangement	1,483	1 (Ho et al., 1997)	1	1.4	Very low
Low instrumental support	156	1 (Avlund, Lund, et al., 2004)	1	2.8	Very low
Self-reported slowing	146	1 (Pine et al., 2000)	1	1.0	Very low
Changes in tiredness	136	1 (Avlund, Vass, & Hendriksen, 2003)	1	1.6	Very low
Hand/wrist pain	412	1 (Eggermont et al., 2014)	1	1.6	Very low
Hand/wrist pain + other pain	412	1 (Eggermont et al., 2014)	1	1.6	Very low
Lower limb joint pain	1,483	1 (Ho et al., 1997)	1	1.1	Very low
Ankle/foot pain	413	1 (Eggermont et al., 2014)	1	1.7	Very low
Injury caused by fall	1,483	1 (Ho et al., 1997)	1	1.3	Very low
Angina	6,981	2 (Guralnik et al., 1993, 2001)	1	1.0–1.6	Very low
Asthma	1,483	1 (Ho et al., 1997)	1	1.1	Very low
Bronchitis	1,483	1 (Ho et al., 1997)	1	1.4	Very low
Claudication	6,981	2 (Guralnik et al., 1993, 2001)	1	1.3–1.4	Very low
COPD	5,263	1 (Guralnik et al., 2001)	1	1.4–1.8	Very low
Dyspnea	6,981	2 (Guralnik et al., 1993, 2001)	1	1.2–1.5	Very low
Headache	1,483	1 (Ho et al., 1997)	1	1.3	Very low
History of heart attack	6,981	2 (Guralnik et al., 1993, 2001)	1	1.0–1.3	Very low
Lack of control over life	456	1 (Ayis et al., 2007)	1	1.0–1.9	Very low
Low satisfaction with life	1,483	1 (Ho et al., 1997)	1	1.2	Very low
Increased vulnerability to accidents	491	1 (Ayis et al., 2007)	1	1.3–2.0	Very low
Increased vulnerability to mugging, burglary	491	1 (Ayis et al., 2007)	1	0.8–1.9	Very low
Estrogen use	2,486	1 (Carbone et al., 2013)	1	0.8	Very low
Vitamin D use	2,486	1 (Carbone et al., 2013)	1	0.8	Very low
Distance to park or other green area	261	1 (Eronen et al., 2013)	1	2.3	Very low

(continued)

Table 1 (continued)

Risk factors	Number of participants ^a	Number of studies examining risk factor	Number of cohorts	Range of OR reported	Quality of evidence
Distance to outdoor recreational facilities	261	1 (Eronen et al., 2013)	1	1.7	Very low
Perceived hilliness in neighborhood	551	1 (Keskinen et al., 2020)	1	1.7	Very low
Hilly terrain and streets in poor condition	266	1 (Rantakokko et al., 2012)	1	1.1–1.4	Very low
Noisy traffic and dangerous crossroads	266	1 (Rantakokko et al., 2012)	1	1.3–1.5	Very low
Inconsistent/unclear association with mobility decline ^c					
Race (non-White)	4,784	3 (Carbone et al., 2013; Crimmins & Saito, 1993; Thorpe et al., 2011)	2	0.9–3.0	Very low
High alcohol consumption	8,464	2 (Ho et al., 1997; LaCroix et al., 1993)	2	1.0–1.2	Very low
Smoker/ex-smoker	8,707	4 (Avlund et al., 2000; Guralnik et al., 2001; Ho et al., 1997; LaCroix et al., 1993)	3	0.8–3.6	Very low
BMI outside of normal range	14,924	8 (Carbone et al., 2013; Crimmins & Saito, 1993; Deshpande et al., 2014; Guralnik et al., 2001; Ho et al., 1997; LaCroix et al., 1993; Lindberg & Tilvis, 1998; Tsuji et al., 2018)	7	0.8–2.1	Very low
Weight loss in past 12 months	5,350	3 (Abizanda et al., 2013; Ho et al., 1997; Lee et al., 2005)	3	1.0–1.6 HR = 0.8–2.0	Very low
Low level of education	12,996	11 (Avlund, Damsgaard, & Osler, 2004; Avlund et al., 1995, 2000; Ayis et al., 2007; Carbone et al., 2013; Deshpande et al., 2014; Guralnik et al., 1993, 2001; Ho et al., 1997; Koster et al., 2005; Thorpe et al., 2011)	6	0.9–2.3	Very low
Income from pension only	1,869	3 (Avlund, Damsgaard, & Osler, 2004; Avlund et al., 1995; Ho et al., 1997)	2	1.1–3.1	Very low
Low social status	2,789	2 (Avlund et al., 1995; Chen et al., 2012)	2	1.0–1.5	Very low
Low social support	6,082	5 (Avlund, Lund, et al., 2004; Ayis et al., 2006; Carbone et al., 2013; Deshpande et al., 2014; Jørgensen et al., 2017)	4	0.9–2.3	Very low
Higher no. musculoskeletal conditions	734	2 (Avlund et al., 2000; Ayis et al., 2007)	2	1.8–2.8	Very low
Back pain	1,077	2 (Eggermont et al., 2014; Simonsick, Aronson, et al., 2018)	2	1.0–1.7	Very low
Hip fracture	7,408	2 (Crimmins & Saito, 1993; Guralnik et al., 2001)	2	1.0–3.2	Very low
Anxiety	4,209	3 (Ayis et al., 2006; Brenes et al., 2005; Mehta et al., 2007)	3	1.0–1.2 HR = 1.3	Very low
Arthritis	10,609	4 (Crimmins & Saito, 1993; Guralnik et al., 1993, 2001; Ho et al., 1997)	3	1.1–1.4	Very low
Circulatory conditions	491	1 (Ayis et al., 2007)	1	1.7	Very low
Cognitive decline	6,376	3 (Ayis et al., 2007; Deshpande et al., 2014; Guralnik et al., 2001)	3	1.0–2.5	Very low
Diabetes	8,464	3 (Guralnik et al., 1993, 2001; Ho et al., 1997)	2	1.3–2.3	Very low
Dizziness/balance difficulties	1,773	2 (Ho et al., 1997; Viljanen et al., 2012)	2	1.0–1.4	Very low
Hearing problems	5,837	6 (Ayis et al., 2006, 2007; Ho et al., 1997; Liljas et al., 2016; Polku et al., 2015; Viljanen et al., 2012)	5	1.1–2.8	Very low
High blood pressure	10,609	3 (Crimmins & Saito, 1993; Guralnik et al., 1993; Ho et al., 1997)	3	1.1–1.2	Very low

Note. OR of 1.4–2.5 = small effect; 2.5–4.25 = moderate effect; ≥4.25 = large effect. The range shown is the lowest and highest OR reported by included studies. RR = risk ratio; HR = hazard ratio; GRADE = Grading of Recommendations Assessment, Development and Evaluation; OR = odds ratio; BMI = body mass index; NSAID = nonsteroidal anti-inflammatory drug; COPD = chronic obstructive pulmonary disease.

^aWhen more than one study examining the same factor was derived from the same cohort, the highest number of participants was added to the participant total. ^bTwo or more studies from two or more cohorts; >75% of studies showing the same direction of effect. ^cMobility modifications defined as any of resting during the task, using an aid, using handrails, reduced frequency, or reduced speed of performing the task. ^dSingle study or multiple studies from single cohort. ^eTwo or more studies from two or more cohorts; <75% of studies showing the same direction of effect, or single study with insufficient data to assess association.

variation in methods of measurement for risk factors and/or mobility outcomes, we were unable to formally pool the results using meta-analysis (Riley et al., 2019).

Of the 107 risk factors examined in these studies, there was consistent evidence (less than or equal to two studies from less than or equal to two cohorts) that 24 factors were associated with mobility decline. Of these 24, three were associated with a moderate/large effect and 21 associated with a small effect. There was also consistent evidence of no association with mobility decline for three factors. The results and discussion will focus on these 27 factors.

Among the remaining 80 risk factors, there was limited evidence (single study or multiple studies from a single cohort) of association with mobility decline for 30, limited evidence of no effect on mobility decline for 30, and inconsistent or unclear effects on mobility decline for 20 factors (Table 1 and Supplementary Table S5 [available online]).

Factors with consistent moderate/large effect on mobility decline. Three factors were consistently associated with mobility decline with moderate–large effect sizes. There was high quality evidence for older age, with large effect sizes reported for participants aged 75 years or older compared to those younger than 75 years. There was also high quality evidence and a dose–response effect observed for mobility modifications (resting during the task, using an aid, using handrails, reduced frequency or reduced speed of performing the task). Moderate quality evidence was found for the presence of widespread pain as a risk factor for mobility decline.

Factors with consistent small effect on mobility decline. For the remaining 21 risk factors consistently associated with mobility decline, the strength of the association represented by ORs was considered small (Table 1). There was high-quality evidence for five demographic and socioeconomic risk factors. Female gender was associated with increased risk of mobility decline, as was low

annual income; low number of financial assets; low diversity in social relations; and low social engagement.

Five risk factors were associated with physical- and fatigue-related issues. Low physical activity was associated with mobility decline with high quality evidence. There was moderate quality evidence for not walking every day, having at least one fall in the past year, higher fatigue after exertion, and greater tiredness in daily activities as risk factors for mobility decline.

Evidence for the consistent small effect of 10 health factors on mobility decline ranged from high to low quality. High quality evidence was found for increasing number of health conditions. Specific health conditions presenting as risk factors for mobility decline included depression (high quality evidence), eye conditions (high quality evidence), heart conditions (high quality evidence), history of stroke (moderate quality evidence), and sensory difficulties (low quality evidence). Other factors that showed a consistent small association with increased risk of mobility decline were hip pain (moderate quality evidence), knee pain (moderate quality evidence), poor self-rated health (high quality evidence), and reporting a hospital stay in the past year (Table 1 and Supplementary Table S5 [available online]).

Factors with consistent lack of effect on mobility decline. Moderate quality evidence was found for three factors which consistently demonstrated no association with increased risk of mobility decline: weight gain in the past 12 months, history of or current cancer diagnosis, and reporting a higher number of physical environment barriers (e.g., long distances, busy traffic, rough or hilly terrain, high crime area; Table 1 and Supplementary Table S5 [available online]).

Model Development Studies

The three model development studies are summarized in Table 2 and Supplementary Table S2 (available online). Follow-up periods

Table 2 Synthesis of Risk Factors for Mobility Decline Examined in Included Model Development Studies

Study	Outcome	Risk factor/s included in final model	Study sample size <i>n</i> (% of participants with the outcome)	Type of validation	Performance
Chaves et al. (2000)	Onset of mobility difficulty. Difficulty in at least one of the tasks: walking 0.8 km; climbing 10 steps; transferring from/into a car or bus	Task modification + walking time for 1 m at usual pace + one-leg stance balance	266 (23.9)	Bootstrap validation	Apparent performance ONLY • AUC = 0.73 • Classification and discrimination measures (correct classification ^a) were presented using multiple predicted probability thresholds
Papachristou et al. (2017)	Incident disability. Difficulty in at least one of the tasks: walking 400 yards; going up or down stairs	Age + slow walking speed + physical inactive + exhaustion	NR (15.0)	Not reported	• AUC = 0.68 (0.63–0.72)
Reynolds and Silverstein (2003)	Onset of walking disability	41 variables including demographic, medical, and behavioral characteristics ^b	3,964 (10.7)	Not reported	Apparent performance ONLY: • AUC = 0.819 • Hosmer–Lemeshow test = 14.47, <i>p</i> = .07 • Adjusted <i>R</i> ² = .1817

Note. Apparent performance is the performance observed in the development data. AUC = area under the curve; IADL = instrumental activities of daily living; NR = not reported.

^aPercentage of individuals whose predicted and observed probabilities are the same. ^bAge, sex, hypertension, diabetes, cancer, lung disease, heart condition, psychiatric problems, arthritis, stroke, married, ethnicity (Hispanic and Black), family network, asset complexity, negative affect, cognition, home modification, weight, impairment, smoke, service use, health insurance, onset of hypertension, onset of diabetes, onset of cancer, onset of lung disease, onset of heart condition, onset of psychiatric problems, onset of arthritis, onset of stroke, deterioration in affect, deterioration in cognition, new home modifications, weight loss, onset of Nagi impairments, onset of IADL impairments, smoking less, new service use, vigorous exercise, and preventive test.

varied from recruitment: 18 months (Chaves et al., 2000); 36 months (Papachristou et al., 2017); and 60 months (Reynolds & Silverstein, 2003). Overall, the three models showed a moderate discriminative ability (area under curve [AUC] = 0.68–0.81), but calibration measures were poorly reported or missing. Only one model was internally validated (Chaves et al., 2000) and none undertook external model validation.

Discussion

This systematic review provides evidence on self-reported factors which predict decline in self-reported mobility between 12 months and 5 years in older adults. Among the 107 risk factors examined in the included studies, we found that 24 had consistent moderate or high quality evidence indicating their association with mobility decline. Effect sizes were generally small, with moderate/large effect sizes observed for three factors: older age (≥ 75 years), widespread pain, and mobility modifications. Three existing model development studies were identified, all demonstrating moderate model discriminatory performance, and limited by high risk of bias because of analytical shortcomings.

To our knowledge, this is the first systematic review to comprehensively summarize available evidence about self-reported risk factors for self-reported mobility decline among older adults. Among the 107 risk factors we reviewed, only one quarter had sufficient evidence from which to draw firm conclusions regarding association with mobility decline. This does not mean that the remaining factors do not play a role in mobility decline in later life, rather that evidence was insufficient to draw such a conclusion. Significant heterogeneity across studies in definitions and assessments of risk factors and mobility decline precluded quantitative synthesis of the data, and consequently our findings must be interpreted with caution.

A 2008 narrative review concluded that the association between older age and increased mobility limitations was statistically significant over 70 years of age (Yeom et al., 2008). Age-related changes in mobility may result from physiological changes, such as decreased muscle strength and power, reduced bone mass, and decreased response to balance perturbations (Byrne, Faure, Keene, & Lamb, 2016). However, age-related changes in mobility may also be associated with, or accelerated by, lack of physical activity, leading to a negative cycle of decline (Brach et al., 2003).

The negative effects of widespread pain on physical functions, including mobility, are well established (Butera, Roff, Buford, & Cruz-Almeida, 2019). However, the pathway from widespread pain to mobility limitation is not clear. Nagi's disablement model suggests that widespread pain may lead to decreased physical activity, and in turn loss of muscle strength and the development of mobility limitations (Verbrugge & Jette, 1994). However, Leveille et al. concluded that pain is a unique domain as a cause of mobility difficulty, independent of the usual pathway via physical impairments (Leveille et al., 2007).

Mobility modifications, such as resting, using an aid, using handrails, reduced frequency, or speed of performing a task are important risk factors among older adults as they have been identified as "preclinical mobility decline" (Fried et al., 2000; Rantakokko et al., 2013). Making such modifications may compensate for the impact of underlying health changes, such as pain, reduced muscle strength, or fatigue on mobility (Fried et al., 2000). An individual's transition to mobility difficulty easily occurs when any internal or external changes overwhelm the compensations being employed. Our findings reiterate the importance of assessing the use of mobility modifications to identify people at high risk of mobility decline so the underlying causes can be addressed.

This review identifies several opportunities for targeting modifiable factors associated with mobility decline, which should be considered in management of older adults at risk of mobility decline, and intervention development. We found high quality evidence of the association between two social factors (low diversity in social relations and low social engagement) and increased risk of mobility decline. This highlights the importance of assessing social outcomes, which are often neglected, and the potential of social prescribing interventions. We also found high quality evidence for low physical activity as a risk factor for mobility decline. Many physical activity programs have been used to improve mobility in older adults, with positive effects (Rantakokko et al., 2013). Assessing and addressing fears of falling in older adults is also important, as indicated by our finding of moderate quality evidence for fear of falling as a risk factor for mobility decline.

Our findings highlight the complexity of managing older adults at risk of mobility decline. Mobility decline in older age is multifactorial, and many older adults may report multiple risk factors concurrently. Combinations of multiple risk factors may have greater impact than the sum of their individual effects, and future research should explore this. Future interventions targeting maintenance of mobility should be multidimensional, combine interventions for multiple risk factors, and control for confounding factors. Our findings also highlight the limited evidence for existing prognostic models for mobility decline. These results may guide future longitudinal studies to develop a prediction instrument for practical application.

Limitations of the Review

Despite the authors' effort to construct a sensitive search strategy, there is a chance that relevant studies were overlooked. Of the included studies, 70% (43/61) came from two countries (United States and Finland), which may impact the generalizability of findings to other populations. Only 3% of studies (2/61) included participants from low-income countries (World Bank, 2020). None of the included studies were conducted in Africa, and only two were conducted in Asia (Hong Kong and Japan). Differing lifestyles and cultures may mean that risk factors are not the same in these settings, and findings should not be generalized. Studies in more diverse international settings are warranted. Grouping together everyone with mobility decline may dilute associations linked with specific subtypes of mobility decline, such as gradual versus rapid decline. Analysis of risk factors according to definitions of mobility decline may give more insight into causal pathways. Almost all included studies (59/61, 97%) used narrow measures of mobility, focusing on ability to walk and/or climb stairs. Use of broader measures of mobility, such as life-space assessment may enable identification of additional risk factors, and/or which risk factors are associated with different aspects of mobility. Finally, by excluding studies with follow-up periods of <12 months or >5 years, there is a risk that studies showing short- or long-term results were missed.

Conclusion

In conclusion, our systematic review identified 24 wide-ranging self-reported risk factors consistently associated with self-reported mobility decline in older people. Older age beyond 75 years, the presence of widespread pain, and mobility modifications were the risk factors with the highest effect.

Acknowledgments

The authors would like to express their gratitude to those study authors who responded to study queries and data requests.

Nicolson and Sanchez-Santos are joint first authors. PN, MS, JB, SK, LW, EW, and SL made substantial contributions to the conception and design of this review; PN, MS, and SK acquired the data; PN, MS, JB, EW, and SL completed analysis and interpretation of the data. PN and MS wrote the first draft of this article, and all authors contributed to revising it critically for important intellectual content. All authors gave their final approval of the version to be published.

The authors do not have any conflicts of interest to declare. SL and EW receive funding from the Collaboration for Leadership in Applied Health Research and Care Oxford at Oxford Health NHS Foundation Trust, and SL, MS, and EW are supported by NIHR Musculoskeletal Biomedical Research Centre, Oxford. PN is funded by a Versus Arthritis Foundation Fellowship (Grant Reference Number 22428). JB is supported by NIHR Research Capability Funding via University Hospitals Coventry and Warwickshire. SK is funded by Cancer Research UK (Grant C49297/A27294). The funders had no role in the design, collection, analysis, or reporting of this review. The views expressed are those of the authors and not necessarily those of the NHS, the NIHR or the Department of Health and Social Care.

References

- Abizanda, P., Romero, L., Sánchez-Jurado, P.M., Martínez-Reig, M., Gomez-Arnedo, L., & Alfonso, S.A. (2013). Frailty and mortality, disability and mobility loss in a Spanish cohort of older adults: The FRADEA study. *Maturitas*, *74*(1), 54–60. PubMed ID: 23107816 doi:10.1016/j.maturitas.2012.09.018
- Alexander, N.B., Guire, K.E., Thelen, D.G., Ashton-Miller, J.A., Schultz, A.B., Grunawalt, J.C., & Giordani, B. (2000). Self-reported walking ability predicts functional mobility performance in frail older adults. *Journal of the American Geriatrics Society*, *48*(11), 1408–1413. PubMed ID: 11083316 doi:10.1111/j.1532-5415.2000.tb02630.x
- An, R., & Lu, L. (2016). Antidepressant use and functional limitations in U.S. older adults. *Journal of Psychosomatic Research*, *80*, 31–36. PubMed ID: 26721545 doi:10.1016/j.jpsychores.2015.11.007
- Auais, M., Ahmed, T., Alvarado, B., Phillips, S.P., Rosendaal, N., Curcio, C.L., . . . Zunzunegui, M.V. (2019). Gender differences in four-year incidence of self-reported and performance-based functional disability: The international mobility in aging study. *Archives of Gerontology and Geriatrics*, *82*, 266–272. PubMed ID: 30878824 doi:10.1016/j.archger.2019.03.002
- Auais, M., French, S., Alvarado, B., Pirkle, C., Belanger, E., & Guralnik, J. (2018). Fear of falling predicts incidence of functional disability 2 years later: A perspective from an international cohort study. *The Journals of Gerontology. Series A, Biological Sciences & Medical Sciences*, *73*(9), 1212–1215. PubMed ID: 29220420 doi:10.1093/geron/glx237
- Avlund, K., Damsgaard, M.T., & Osler, M. (2004). Social position and functional decline among non-disabled old men and women. *European Journal of Public Health*, *14*(2), 212–216. PubMed ID: 15230514 doi:10.1093/eurpub/14.2.212
- Avlund, K., Damsgaard, M.T., Sakari-Rantala, R., Laukkanen, P., & Schroll, M. (2002). Tiredness in daily activities among nondisabled old people as determinant of onset of disability. *Journal of Clinical Epidemiology*, *55*(10), 965–973. PubMed ID: 12464372 doi:10.1016/S0895-4356(02)00463-8
- Avlund, K., Davidsen, M., & Schultz-Larsen, K. (1995). Changes in functional ability from ages 70 to 75. A Danish longitudinal study. *Journal of Aging and Health*, *7*(2), 254–282. PubMed ID: 10165956 doi:10.1177/089826439500700205
- Avlund, K., Lund, R., Holstein, B.E., & Due, P. (2004). Social relations as determinant of onset of disability in aging. *Archives of Gerontology and Geriatrics*, *38*(1), 85–99. PubMed ID: 14599708 doi:10.1016/j.archger.2003.08.003
- Avlund, K., Osler, M., Damsgaard, M.T., Christensen, U., & Schroll, M. (2000). The relations between musculoskeletal diseases and mobility among old people: Are they influenced by socio-economic, psychosocial, and behavioral factors? *International Journal of Behavioral Medicine*, *7*(4), 322–339. doi:10.1207/S15327558IJBM0704_04
- Avlund, K., Pedersen, A.N., & Schroll, M. (2003). Functional decline from age 80 to 85: Influence of preceding changes in tiredness in daily activities. *Psychosomatic Medicine*, *65*(5), 771–777. PubMed ID: 14508019 doi:10.1097/01.PSY.0000082640.61645.BF
- Avlund, K., Vass, M., & Hendriksen, C. (2003). Onset of mobility disability among community-dwelling old men and women. The role of tiredness in daily activities. *Age and Ageing*, *32*(6), 579–584. PubMed ID: 14599997 doi:10.1093/ageing/afg101
- Ayis, S., Goberman-Hill, R., Bowling, A., & Ebrahim, S. (2006). Predicting catastrophic decline in mobility among older people. *Age and Ageing*, *35*(4), 382–387. PubMed ID: 16638760 doi:10.1093/ageing/afl004
- Ayis, S.A., Bowling, A., Goberman-Hill, R., & Ebrahim, S. (2007). The effect of definitions of activities of daily living on estimates of changing ability among older people. *International Journal of Rehabilitation Research*, *30*(1), 39–46. PubMed ID: 17293719 doi:10.1097/MRR.0b013e328013dafb
- Brach, J.S., FitzGerald, S., Newman, A.B., Kelsey, S., Kuller, L., VanSwearingen, J.M., & Kriska, A.M. (2003). Physical activity and functional status in community-dwelling older women: A 14-year prospective study. *Archives of Internal Medicine*, *163*(21), 2565–2571. PubMed ID: 14638556 doi:10.1001/archinte.163.21.2565
- Brenes, G.A., Guralnik, J.M., Williamson, J.D., Fried, L.P., Simpson, C., Simonsick, E.M., & Penninx, B.W. (2005). The influence of anxiety on the progression of disability. *Journal of the American Geriatrics Society*, *53*(1), 34–39. PubMed ID: 15667373 doi:10.1111/j.1532-5415.2005.53007.x
- Butera, K.A., Roff, S.R., Buford, T.W., & Cruz-Almeida, Y. (2019). The impact of multisite pain on functional outcomes in older adults: Biopsychosocial considerations. *Journal of Pain Research*, *12*, 1115–1125. PubMed ID: 30992680 doi:10.2147/JPR.S192755
- Byrne, C., Faure, C., Keene, D.J., & Lamb, S.E. (2016). Ageing, muscle power and physical function: A systematic review and implications for pragmatic training interventions. *Sports Medicine*, *46*(9), 1311–1332. PubMed ID: 26893098 doi:10.1007/s40279-016-0489-x
- Carbone, L.D., Satterfield, S., Liu, C., Kwok, K.C., Neogi, T., Tolley, E., & Nevitt, M. (2013). Assistive walking device use and knee osteoarthritis: Results from the health, aging and body composition study (health ABC study). *Archives of Physical Medicine and Rehabilitation*, *94*(2), 332–339. PubMed ID: 23041146 doi:10.1016/j.apmr.2012.09.021
- Chaves, P.H., Garrett, E.S., & Fried, L.P. (2000). Predicting the risk of mobility difficulty in older women with screening nomograms: The women's health and aging study II. *Archives of Internal Medicine*, *160*(16), 2525–2533. PubMed ID: 10979066 doi:10.1001/archinte.160.16.2525
- Chen, B., Covinsky, K.E., Stijacic Cenzer, I., Adler, N., & Williams, B.A. (2012). Subjective social status and functional decline in older adults. *Journal of General Internal Medicine*, *27*(6), 693–699. PubMed ID: 22215272 doi:10.1007/s11606-011-1963-7
- Chinn, S. (2000). A simple method for converting an odds ratio to effect size for use in meta-analysis. *Statistics in Medicine*, *19*(22), 3127–3131. PubMed ID: 11113947 doi:10.1002/1097-0258(20001130)19:22<3127::AID-SIM784>3.0.CO;2-M

- Chung, J., Demiris, G., & Thompson, H.J. (2015). Instruments to assess mobility limitation in community-dwelling older adults: A systematic review. *Journal of Aging and Physical Activity*, 23(2), 298–313. PubMed ID: 24589525 doi:10.1123/japa.2013-0181
- Crimmins, E.M., & Saito, Y. (1993). Getting better and getting worse: Transitions in functional status among older Americans. *Journal of Aging and Health*, 5(1), 3–36. doi:10.1177/089826439300500101
- Deshpande, N., Metter, J.E., Guralnik, J., Bandinelli, S., & Ferrucci, L. (2014). Sensorimotor and psychosocial determinants of 3-year incident mobility disability in middle-aged and older adults. *Age and Ageing*, 43(1), 64–69. PubMed ID: 24042004 doi:10.1093/ageing/af135
- Duan-Porter, W., Vo, T.N., Ullman, K., Langsetmo, L., Strotmeyer, E.S., Taylor, B.C., . . . Ensrud, K.E. (2019). Hospitalization-associated change in gait speed and risk of functional limitations for older adults. *The Journals of Gerontology. Series A, Biological Sciences & Medical Sciences*, 74(10), 1657–1663. PubMed ID: 30715162 doi:10.1093/gerona/glz027
- Eggermont, L.H., Leveille, S.G., Shi, L., Kiely, D.K., Shmerling, R.H., Jones, R.N., . . . Bean, J.F. (2014). Pain characteristics associated with the onset of disability in older adults: The maintenance of balance, independent living, intellect, and zest in the Elderly Boston Study. *Journal of the American Geriatric Society*, 62(6), 1007–1016. PubMed ID: 24823985 doi:10.1111/jgs.12848
- Eronen, J., von Bonsdorff, M., Rantakokko, M., & Rantanen, T. (2013). Environmental facilitators for outdoor walking and development of walking difficulty in community-dwelling older adults. *European Journal of Ageing*, 11(1), 67–75. PubMed ID: 28804315 doi:10.1007/s10433-013-0283-7
- Fried, L.P., Bandeen-Roche, K., Chaves, P.H., & Johnson, B.A. (2000). Preclinical mobility disability predicts incident mobility disability in older women. *The Journals of Gerontology. Series A, Biological Sciences & Medical Sciences*, 55(1), M43–M52. PubMed ID: 10719772 doi:10.1093/gerona/55.1.M43
- Guralnik, J.M., Ferrucci, L., Balfour, J.L., Volpato, S., & Di Iorio, A. (2001). Progressive versus catastrophic loss of the ability to walk: Implications for the prevention of mobility loss. *Journal of the American Geriatric Society*, 49(11), 1463–1470. PubMed ID: 11890584 doi:10.1046/j.1532-5415.2001.4911238.x
- Guralnik, J.M., LaCroix, A.Z., Abbott, R.D., Berkman, L.F., Satterfield, S., Evans, D.A., & Wallace, R.B. (1993). Maintaining mobility in late life. I. Demographic characteristics and chronic conditions. *American Journal of Epidemiology*, 137(8), 845–857. PubMed ID: 8484376 doi:10.1093/oxfordjournals.aje.a116746
- Hayden, J.A., Tougas, M.E., Riley, R., Iles, R., & Pincus, T. (2014). Individual recovery expectations and prognosis of outcomes in non-specific low back pain: Prognostic factor exemplar review. *Cochrane Database of Systematic Reviews*, 2014(9), CD011284. doi:10.1002/14651858.cd011284
- Hayden, J.A., van der Windt, D.A., Cartwright, J.L., Cote, P., & Bombardier, C. (2013). Assessing bias in studies of prognostic factors. *Annals of Internal Medicine*, 158(4), 280–286. PubMed ID: 23420236 doi:10.7326/0003-4819-158-4-201302190-00009
- Ho, S.C., Woo, J., Yuen, Y.K., Sham, A., & Chan, S.G. (1997). Predictors of mobility decline: The Hong Kong old-old study. *The Journals of Gerontology. Series A, Biological Sciences & Medical Sciences*, 52(6), M356–M362. PubMed ID: 9402942 doi:10.1093/gerona/52A.6.M356
- Huguet, A., Hayden, J.A., Stinson, J., McGrath, P.J., Chambers, C.T., Tougas, M.E., & Wozney, L. (2013). Judging the quality of evidence in reviews of prognostic factor research: Adapting the GRADE framework. *Systematic Reviews*, 2(1), 71. PubMed ID: 24007720 doi:10.1186/2046-4053-2-71
- Jørgensen, T.S.H., Lund, R., Siersma, V.D., & Nilsson, C.J. (2017). Interplay between financial assets and social relations on decline in physical function and mortality among older people. *European Journal of Ageing*, 15(2), 133–142. PubMed ID: 29867298 doi:10.1007/s10433-017-0437-0
- Keskinen, K.E., Rantakokko, M., Suomi, K., Rantanen, T., & Portegijs, E. (2020). Hilliness and the development of walking difficulties among community-dwelling older people. *Journal of Aging and Health*, 32(5–6), 278–284. PubMed ID: 30587067 doi:10.1177/0898264318820448
- Koster, A., Penninx, B.W., Bosma, H., Kempen, G.I., Harris, T.B., Newman, A.B., . . . Kritchevsky, S.B. (2005). Is there a biomedical explanation for socioeconomic differences in incident mobility limitation? *The Journals of Gerontology. Series A, Biological Sciences & Medical Sciences*, 60(8), 1022–1027. PubMed ID: 16127107 doi:10.1093/gerona/60.8.1022
- LaCroix, A.Z., Guralnik, J.M., Berkman, L.F., Wallace, R.B., & Satterfield, S. (1993). Maintaining mobility in late life. II. Smoking, alcohol consumption, physical activity, and body mass index. *American Journal of Epidemiology*, 137(8), 858–869. PubMed ID: 8484377 doi:10.1093/oxfordjournals.aje.a116747
- Lang, I.A., Llewellyn, D.J., Langa, K.M., Wallace, R.B., & Melzer, D. (2008). Neighbourhood deprivation and incident mobility disability in older adults. *Age and Ageing*, 37(4), 403–410. PubMed ID: 18487260 doi:10.1093/ageing/afn092
- Lee, J.S., Kritchevsky, S.B., Tyllavsky, F., Harris, T., Simonsick, E.M., Rubin, S.M., & Newman, A.B. (2005). Weight change, weight change intention, and the incidence of mobility limitation in well-functioning community-dwelling older adults. *The Journals of Gerontology. Series A, Biological Sciences & Medical Sciences*, 60(8), 1007–1012. PubMed ID: 16127104 doi:10.1093/gerona/60.8.1007
- Leveille, S.G., Bean, J., Ngo, L., McMullen, W., & Guralnik, J.M. (2007). The pathway from musculoskeletal pain to mobility difficulty in older disabled women. *Pain*, 128(1–2), 69–77. PubMed ID: 17055167 doi:10.1016/j.pain.2006.08.031
- Leveille, S.G., Ling, S., Hochberg, M.C., Resnick, H.E., Bandeen-Roche, K.J., Won, A., & Guralnik, J.M. (2001). Widespread musculoskeletal pain and the progression of disability in older disabled women. *Annals of Internal Medicine*, 135(12), 1038–1046. PubMed ID: 11747382 doi:10.7326/0003-4819-135-12-200112180-00007
- Liljas, A.E.M., Wannamethee, S.G., Whincup, P.H., Papacosta, O., Walters, K., Iliffe, S., . . . Ramsay, S.E. (2016). Hearing impairment and incident disability and all-cause mortality in older British community-dwelling men. *Age and Ageing*, 45(5), 661–666. PubMed ID: 27146303 doi:10.1093/ageing/afw080
- Lindberg, O., & Tilvis, R.S. (1998). Does the metabolic syndrome predict mobility impairment in the elderly? *Archives of Gerontology and Geriatrics*, 26(2), 131–139. PubMed ID: 18653132 doi:10.1016/S0167-4943(97)00037-X
- Mänty, M., Heinonen, A., Leinonen, R., Törmäkangas, T., Sakari-Rantala, R., Hirvensalo, M., . . . Rantanen, T. (2007). Construct and predictive validity of a self-reported measure of preclinical mobility limitation. *Archives of Physical Medicine and Rehabilitation*, 88(9), 1108–1113. PubMed ID: 17826454 doi:10.1016/j.apmr.2007.06.016
- Mänty, M., Heinonen, A., Viljanen, A., Pajala, S., Koskenvuo, M., Kaprio, J., & Rantanen, T. (2009). Outdoor and indoor falls as predictors of mobility limitation in older women. *Age and Ageing*, 38(6), 757–761. PubMed ID: 19779051 doi:10.1093/ageing/afp178
- Mehta, K.M., Yaffe, K., Brenes, G.A., Newman, A.B., Shorr, R.I., Simonsick, E.M., . . . Covinsky, K.E. (2007). Anxiety symptoms and decline in physical function over 5 years in the health, aging and body composition study. *Journal of the American Geriatrics Society*,

- 55(2), 265–270. PubMed ID: [17302665](#) doi:[10.1111/j.1532-5415.2007.01041.x](#)
- Moher, D., Liberati, A., Tetzlaff, J., & Altman, D.G. (2009). Preferred reporting items for systematic reviews and meta-analyses: The PRISMA statement. *Annals of Internal Medicine*, *151*(4), 264–269. PubMed ID: [19622511](#) doi:[10.7326/0003-4819-151-4-200908180-00135](#)
- Moons, K.G., de Groot, J.A., Bouwmeester, W., Vergouwe, Y., Mallett, S., Altman, D.G., . . . Collins, G.S. (2014). Critical appraisal and data extraction for systematic reviews of prediction modelling studies: The CHARMS checklist. *PLoS Medicine*, *11*(10), e1001744. PubMed ID: [25314315](#) doi:[10.1371/journal.pmed.1001744](#)
- Moons, K.G.M., Royston, P., Vergouwe, Y., Grobbee, D.E., & Altman, D.G. (2009). Prognosis and prognostic research: What, why, and how? *British Medical Journal*, *338*, b375. PubMed ID: [19237405](#) doi:[10.1136/bmj.b375](#)
- Nilsson, C.J., Avlund, K., & Lund, R. (2010). Social inequality in onset of mobility disability among older Danes: The mediation effect of social relations. *Journal of Aging and Health*, *22*(4), 522–541. PubMed ID: [20231729](#) doi:[10.1177/0898264309359684](#)
- Nilsson, C.J., Lund, R., & Avlund, K. (2008). Cohabitation status and onset of disability among older Danes: Is social participation a possible mediator? *Journal of Aging and Health*, *20*(2), 235–253. PubMed ID: [18089766](#) doi:[10.1177/0898264307310474](#)
- Onder, G., Penninx, B.W., Guralnik, J.M., Jones, H., Fried, L.P., Pahor, M., & Williamson, J.D. (2003). Sexual satisfaction and risk of disability in older women. *The Journal of Clinical Psychiatry*, *64*(10), 1177–1182. PubMed ID: [14658965](#) doi:[10.4088/JCP.v64n1006](#)
- Papachristou, E., Wannamethee, S.G., Lennon, L.T., Papacosta, O., Whincup, P.H., Iliffe, S., & Ramsay, S.E. (2017). Ability of self-reported frailty components to predict incident disability, falls, and all-cause mortality: Results from a population-based study of older British men. *Journal of the American Medical Directors Association*, *18*(2), 152–157. PubMed ID: [27742583](#) doi:[10.1016/j.jamda.2016.08.020](#)
- Pine, Z.M., Gurland, B., & Chren, M.M. (2000). Report of having slowed down: Evidence for the validity of a new way to inquire about mild disability in elders. *The Journals of Gerontology. Series A, Biological Sciences & Medical Sciences*, *55*(7), M378–M383. PubMed ID: [10898254](#) doi:[10.1093/gerona/55.7.M378](#)
- Pine, Z.M., Gurland, B., & Chren, M.M. (2002). Use of a cane for ambulation: Marker and mitigator of impairment in older people who report no difficulty walking. *Journal of the American Geriatrics Society*, *50*(2), 263–268. PubMed ID: [12028207](#) doi:[10.1046/j.1532-5415.2002.50057.x](#)
- Polku, H., Mikkola, T.M., Rantakokko, M., Portegijs, E., Törmäkangas, T., Rantanen, T., & Viljanen, A. (2015). Self-reported hearing difficulties and changes in life-space mobility among community-dwelling older adults: A two-year follow-up study. *BMC Geriatrics*, *15*(1), 121. PubMed ID: [26459630](#) doi:[10.1186/s12877-015-0119-8](#)
- Rantakokko, M., Iwarsson, S., Mänty, M., Leinonen, R., & Rantanen, T. (2012). Perceived barriers in the outdoor environment and development of walking difficulties in older people. *Age and Ageing*, *41*(1), 118–121. PubMed ID: [22086965](#) doi:[10.1093/ageing/afr136](#)
- Rantakokko, M., Mänty, M., & Rantanen, T. (2013). Mobility decline in old age. *Exercise and Sport Sciences Reviews*, *41*(1), 19–25. PubMed ID: [23038241](#) doi:[10.1097/JES.0b013e3182556f1e](#)
- Rantakokko, M., Portegijs, E., Viljanen, A., Iwarsson, S., & Rantanen, T. (2016). Mobility modification alleviates environmental influence on incident mobility difficulty among community-dwelling older people: A two-year follow-up study. *PLoS One*, *11*(4), e0154396. PubMed ID: [27104750](#) doi:[10.1371/journal.pone.0154396](#)
- Reynolds, S.L., & Silverstein, M. (2003). Observing the onset of disability in older adults. *Social Science & Medicine*, *57*(10), 1875–1889. PubMed ID: [14499512](#) doi:[10.1016/S0277-9536\(03\)00053-4](#)
- Riley, R.D., Moons, K.G.M., Snell, K.I.E., Ensor, J., Hooft, L., Altman, D.G., . . . Debray, T.P.A. (2019). A guide to systematic review and meta-analysis of prognostic factor studies. *BMJ*, *364*, k4597. PubMed ID: [30700442](#) doi:[10.1136/bmj.k4597](#)
- Sauvagat, C., Tsuji, I., Aonuma, T., & Hisamichi, S. (1999). Health-life expectancy according to various functional levels. *Journal of the American Geriatrics Society*, *47*(11), 1326–1331. PubMed ID: [10573441](#) doi:[10.1111/j.1532-5415.1999.tb07433.x](#)
- Schroll, M., Avlund, K., & Davidsen, M. (1997). Predictors of five-year functional ability in a longitudinal survey of men and women aged 75 to 80. The 1914-population in Glostrup, Denmark. *Aging*, *9*(1–2), 143–152. PubMed ID: [9177598](#) doi:[10.1007/bf03340140](#)
- Simonsick, E.M., Aronson, B., Schrack, J.A., Hicks, G.E., Jerome, G.J., Patel, K.V., . . . Ferrucci, L. (2018). Lumbopelvic pain and threats to walking ability in well-functioning older adults: Findings from the Baltimore Longitudinal Study of Aging. *Journal of the American Geriatrics Society*, *66*(4), 714–720. PubMed ID: [29411349](#) doi:[10.1111/jgs.15280](#)
- Simonsick, E.M., Glynn, N.W., Jerome, G.J., Shardell, M., Schrack, J.A., & Ferrucci, L. (2016). Fatigued, but not frail: Perceived fatigability as a marker of impending decline in mobility-intact older adults. *Journal of the American Geriatrics Society*, *64*(6), 1287–1292. PubMed ID: [27253228](#) doi:[10.1111/jgs.14138](#)
- Simonsick, E.M., Schrack, J.A., Santanasto, A.J., Studenski, S.A., Ferrucci, L., & Glynn, N.W. (2018). Pittsburgh Fatigability Scale: One-page predictor of mobility decline in mobility-intact older adults. *Journal of the American Geriatrics Society*, *66*(11), 2092–2096. PubMed ID: [30315707](#) doi:[10.1111/jgs.15531](#)
- Syddall, H.E., Westbury, L.D., Cooper, C., & Sayer, A.A. (2015). Self-reported walking speed: A useful marker of physical performance among community-dwelling older people? *Journal of the American Medical Directors Association*, *16*(4), 323–328. PubMed ID: [25523286](#) doi:[10.1016/j.jamda.2014.11.004](#)
- Thakral, M., Shi, L., Shmerling, R.H., Bean, J.F., & Leveille, S.G. (2014). A stiff price to pay: Does joint stiffness predict disability in an older population? *Journal of the American Geriatrics Society*, *62*(10), 1891–1899. PubMed ID: [25333527](#) doi:[10.1111/jgs.13070](#)
- Thorpe, R.J., Jr., Koster, A., Kritchevsky, S.B., Newman, A.B., Harris, T., Ayonayon, H.N., . . . Health Aging and Body Composition Study. (2011). Race, socioeconomic resources, and late-life mobility and decline: Findings from the health, aging, and body composition study. *The Journals of Gerontology. Series A, Biological Sciences and Medical Sciences*, *66*(10), 1114–1123. PubMed ID: [21743093](#) doi:[10.1093/gerona/66.10.1114](#)
- Tsuji, T., Rantakokko, M., Portegijs, E., Viljanen, A., & Rantanen, T. (2018). The effect of body mass index, lower extremity performance, and use of a private car on incident life-space restriction: A two-year follow-up study. *BMC Geriatrics*, *18*(1), 271. PubMed ID: [30409120](#) doi:[10.1186/s12877-018-0956-3](#)
- van den Berg, T., Heymans, M.W., Leone, S.S., Vergouw, D., Hayden, J.A., Verhagen, A.P., & de Vet, H.C.W. (2013). Overview of data-synthesis in systematic reviews of studies on outcome prediction models. *BMC Medical Research Methodology*, *13*(1), 42. PubMed ID: [23497181](#) doi:[10.1186/1471-2288-13-42](#)
- van den Brink, C.L., Tjihuis, M., van den Bos, G.A., Giampaoli, S., Kivinen, P., Nissinen, A., & Kromhout, D. (2004). Effect of widowhood on disability onset in elderly men from three European countries. *Journal of the American Geriatrics Society*, *52*(3), 353–358. PubMed ID: [14962148](#) doi:[10.1111/j.1532-5415.2004.52105.x](#)

- Verbrugge, L.M., & Jette, A.M. (1994). The disablement process. *Social Science & Medicine*, 38(1), 1–14. PubMed ID: 8146699 doi:10.1016/0277-9536(94)90294-1
- Viljanen, A., Kulmala, J., Rantakokko, M., Koskenvuo, M., Kaprio, J., & Rantanen, T. (2012). Fear of falling and coexisting sensory difficulties as predictors of mobility decline in older women. *The Journals of Gerontology. Series A, Biological Sciences & Medical Sciences*, 67(11), 1230–1237. PubMed ID: 22546957 doi:10.1093/gerona/gls134
- Visser, M., Simonsick, E.M., Colbert, L.H., Brach, J., Rubin, S.M., Kritchevsky, S.B., . . . Harris, T.B. (2005). Type and intensity of activity and risk of mobility limitation: The mediating role of muscle parameters. *Journal of the American Geriatrics Society*, 53(5), 762–770. PubMed ID: 15877550 doi:10.1111/j.1532-5415.2005.53257.x
- Webber, S.C., Porter, M.M., & Menec, V.H. (2010). Mobility in older adults: A comprehensive framework. *Gerontologist*, 50(4), 443–450. PubMed ID: 20145017 doi:10.1093/geront/gnq013
- Weiss, C.O., Wolff, J.L., Eggleston, B., Seplaki, C.L., & Fried, L.P. (2012). Incident preclinical mobility disability (PCMD) increases future risk of new difficulty walking and reduction in walking activity. *Archives of Gerontology and Geriatrics*, 54(3), e329–e333. PubMed ID: 21944428 doi:10.1016/j.archger.2011.08.018
- Wolff, R.F., Moons, K.G.M., Riley, R.D., Whiting, P.F., Westwood, M., Collins, G.S., . . . PROBAST Group. (2019). PROBAST: A tool to assess the risk of bias and applicability of prediction model studies. *Annals of Internal Medicine*, 170(1), 51–58. PubMed ID: 30596875 doi:10.7326/M18-1376
- World Bank. (2020). World Bank list of economies. Retrieved from <https://datahelpdesk.worldbank.org/knowledgebase/articles/906519-world-bank-country-and-lending-groups>
- Yeom, H.A., Fleury, J., & Keller, C. (2008). Risk factors for mobility limitation in community-dwelling older adults: A social ecological perspective. *Geriatric Nursing*, 29(2), 133–140. PubMed ID: 18394514 doi:10.1016/j.gerinurse.2007.07.002